

THE INVARIANT MEASURES OF SOME INFINITE INTERVAL EXCHANGE MAPS

W. PATRICK HOOPER

1. INTRODUCTION

Let $\mathbb{G}$ be an infinite group generated by the elements $\gamma_1, \dots, \gamma_n$. A *skew rotation* is transformation of $\mathbb{R}/\mathbb{Z} \times \mathbb{G}$ of the form

$$(1.1) \quad T(x, g) = (x + \alpha, \psi(x)g),$$

where $\psi(x) = \gamma_i$ iff $x \in [\frac{i-1}{n}, \frac{i}{n})$ and $\alpha \in \mathbb{R}/\mathbb{Z}$ is irrational. Skew rotations are natural examples of infinite interval exchange maps, and of deterministic random walks on groups.

In the case that $\mathbb{G}$ is the non-abelian free group generated by $\gamma_1, \dots, \gamma_n$, the system is uninteresting in the sense that it is totally dissipative. However, if $\mathbb{G}$ satisfies the relation

$$\gamma_n \gamma_{n-1} \dots \gamma_1 = e,$$

then many directions become recurrent. This collection includes all but countably many points in the limit set Λ_n of the group action of $\Gamma_n \subset PSL(2, \mathbb{Z})$ on $\mathbb{R}$ generated by

$$(1.2) \quad z \mapsto z + 1 \quad \text{and} \quad z \mapsto \frac{(n+1)z - 1}{n^2z - n + 1}.$$

When $n > 2$, the limit set Λ_n is a Cantor set, while when $n = 2$ this group is conjugate to the congruence two subgroup of $PSL(2, \mathbb{Z})$. We show that T is recurrent when

$$\alpha \in \Omega_n = \Lambda_n \setminus (\mathbb{Q} \cup \{\text{endpoints of intervals in } \mathbb{R} \setminus \Lambda_n\}).$$

Moreover, for $\alpha \in \Omega_n$, the locally finite ergodic invariant measures for the system can be classified. We will show how each positive eigenfunction for the adjacency operator of a certain Cayley graph of the group $\mathbb{G}$ gives rise to a unique locally finite invariant measure for T . Moreover, the locally finite ergodic invariant measures are in bijective correspondence with the extremal positive eigenfunctions.

A group homomorphism from $\mathbb{G}$ to the multiplicative group of positive real numbers is an example of a positive eigenfunctions for the adjacency operator. The invariant measures that arise from these group homomorphisms are the Maharam measures. When $\mathbb{G}$ is nilpotent, a theorem of Margulis [Mar66] implies that the extremal positive eigenfunctions are precisely the group homomorphisms $\mathbb{G} \rightarrow \mathbb{R}_+$. As a consequence, when $\mathbb{G}$ is nilpotent and $\alpha \in \Omega_n$, the locally finite ergodic invariant measures are the Maharam measures.

In contrast, consider the case when $\mathbb{G}$ is the non-abelian free group of rank $n - 1$,

$$\mathbb{G} = \langle \gamma_1, \dots, \gamma_n \mid \gamma_n \gamma_{n-1} \dots \gamma_1 = e \rangle.$$

Here a theorem of Picardello and Woess [PW88] implies that the space of extremal positive eigenfunctions is $\partial_{hyp}\mathbb{G} \times [0, \infty)$ where $\partial_{hyp}\mathbb{G}$ denotes the δ -hyperbolic boundary. Again $\partial_{hyp}\mathbb{G} \times [0, \infty)$ is also the space of locally finite ergodic invariant measures, when $\alpha \in \Omega_n$.

Supported by N.S.F. Postdoctoral Fellowship DMS-0803013.

These results hold in a more general setting. Namely, Thurston gave a construction which produces a finite number of translation surfaces from a connected bipartite graph $\mathcal{G}$, and a positive eigenvector for the adjacency operator [Thu88]. Thurston's used this construction to provide examples of closed surfaces admitting pseudo-Anosov automorphisms. Each such surface gives rise to a one-parameter family of interval exchange transformations. These are infinite interval exchange transformations if $\mathcal{G}$ is infinite. We are able to classify the locally finite ergodic invariant measures for many of these infinite interval exchange transformations. We explain these more general results in section 1.2. Since the point of view of Thurston's construction is more general, our main results are stated in this context. In sections 5 and 16.2-16.3, we explain several applications to skew rotations.

1.1. Historical remarks. Ergodicity of the skew rotation with $\mathbb{G} = \mathbb{Z}$ was proved for some irrational α by Schmidt [Sch78]. The same statement was shown to hold for all α by Conze and Keane [CK76]. More recently, Aaronson, Nakada, Sarig and Solomyak showed that when α is irrational, the locally finite ergodic invariant measures of T are the Maharam measures [ANSS02]. Hubert and Weiss discovered that these skew rotations with $\mathbb{G} = \mathbb{Z}$ arise as return maps of straight-line flows on the staircase surface of Figure 1 [HW08]. We utilize this observation to connect Thurston's construction to general skew rotations with $\mathbb{G}$ infinite and discrete.

The philosophy of the proofs in this paper come from Teichmüller theory. In particular, we follow the spirit of a criterion of Masur [Mas82] which guarantees unique ergodicity of an interval exchange transformation involving finitely many intervals. Masur's criterion uses the Teichmüller flow on moduli space to demonstrate unique ergodicity. Rather than using the Teichmüller flow directly, we make use of the inherent symmetries (affine automorphisms) of surfaces produced using Thurston's construction to renormalize the space of invariant measures. This iterative process was inspired by Smillie and Ulcigrai's work on the regular octagon [SU09], which produced detailed information about the trajectories of the straight-line flow.

Infinite translation surfaces arise naturally in the context of billiards in polygons with angles which are irrational multiples of π [ZK75]. In this context, infinite translation surfaces have been studied for applications to billiards. Recently however, there has been interest in infinite translation surfaces which arise from compact translation surfaces either via infinite covers or via geometric limits. See [HW08], [Hoo08b], [HS09] and [HW10] for instance. Other recent work on infinite translation surfaces has included [CGL06] and [PSV09].

1.2. Main results. A well studied construction of Thurston takes a connected, bipartite, ribbon graph $\mathcal{G}_0$ and produces a translation surface with an non-trivial automorphism group of affine symmetries, including pseudo-Anosov automorphisms [Thu88]. (McMullen introduced the ribbon graph structure to the construction in [McM06].) *Translation surfaces* are surfaces that can be constructed by gluing together edges of polygons in the plane by translations. Given a translation surface S , it is commonplace to study the natural family $\{F_\theta^t : S \rightarrow S\}_{\theta \in \mathbb{S}^1}$ of *straight-line flows*, which are parametrized by a unit vector (a direction) $\theta \in \mathbb{S}^1 \subset \mathbb{R}^2$. In local coordinates, these flows are given by

$$(1.3) \quad F_\theta^t(x, y) = (x, y) + t\theta.$$

The surfaces constructed by Thurston's method are defined by gluing rectangles with horizontal and vertical sides together. For such surfaces, it is natural to study the return maps

T_θ of the flows F_θ^t to the union of horizontal edges of the rectangles. These return maps are always interval exchange maps.

To describe the input to the modified version of Thurston's construction we need to introduce some concepts from graph theory. Let $\mathcal{G}$ be any graph and let $\mathcal{V}$ denote the collection of vertices. The *adjacency operator* is the operator $\mathbf{A} : \mathbb{R}^{\mathcal{V}} \rightarrow \mathbb{R}^{\mathcal{V}}$ defined by

$$(1.4) \quad \mathbf{A}(\mathbf{f})(v) = \sum_{w \sim v} \mathbf{f}(w),$$

where the sum is taken over edges $\overline{vw}$ with v as one endpoint. An *eigenfunction* of $\mathbf{A}$ is a function $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ which satisfies $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$ for some $\lambda \in \mathbb{R}$.

This paper observes that Thurston's construction works with a minor modification for an infinite graph. Throughout this paper $\mathcal{G}$ will denote an infinite, connected, bipartite, ribbon graph of bounded valance. See Definition 2.2. The modified construction takes as input $\mathcal{G}$ and a positive eigenfunction $\mathbf{w}$ of the adjacency operator and produces a translation surface $S(\mathcal{G}, \mathbf{w})$. This surface is always noncompact and infinite genus. These surfaces again are constructed by gluing rectangles with horizontal and vertical sides together. We will study the straight-line flows $F_\theta^t : S(\mathcal{G}, \mathbf{w}) \rightarrow S(\mathcal{G}, \mathbf{w})$. It is convenient to look at the section of these flows $H(\mathcal{G}, \mathbf{w})$ consisting of the union of all horizontal edges of rectangles making up $S(\mathcal{G}, \mathbf{w})$. The subset $H(\mathcal{G}, \mathbf{w})$ is naturally a countable union of circles, and the return map $T_\theta : H(\mathcal{G}, \mathbf{w}) \rightarrow H(\mathcal{G}, \mathbf{w})$ is an infinite interval exchange map.

Figure 1 provides an example of the construction. Let $\mathcal{G}_{\mathbb{Z}}$ denote the graph whose vertices are integers with edges joining integers which differ by ± 1 . The constant function $\mathbf{1} \in \mathbb{R}^{\mathbb{Z}}$ whose values are identically one is a positive eigenfunction for the adjacency operator. In [HW08], Hubert and Weiss recognized that the interval exchange transformations T_θ associated to this example are precisely the skew rotations studied in [ANSS02]. Figure 2 provides a second example which will be relevant later.




FIGURE 1. The graph $\mathcal{G}_{\mathbb{Z}}$ and the staircase surface $S(\mathcal{G}_{\mathbb{Z}}, \mathbf{1})$ are shown above. Opposite edges of the staircase are glued by horizontal or vertical translations.

The constructed $S(\mathcal{G}, \mathbf{w})$ are always quite symmetric in the sense that the surface is preserved under some affine changes of coordinates. These special coordinate changes are described by elements of $SL(2, \mathbb{R})$ and form a group known as the Veech group of the surface

$S(\mathcal{G}, \mathbf{w})$. See §2.2. This modified construction produces surfaces whose Veech groups always contain a free group of rank 2, which we denote by $G = \langle h, v \rangle$. If $\mathbf{A}(\mathbf{w}) = \lambda \mathbf{w}$ with $\mathbf{w}$ positive, then $\lambda \geq 2$ and the Veech group of $S(\mathcal{G}, \mathbf{w})$ contains the image of the representation $\rho_\lambda : G \rightarrow SL(2, \mathbb{Z})$ determined by

$$(1.5) \quad \rho_\lambda^h = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \rho_\lambda^v = \begin{bmatrix} 1 & 0 \\ \lambda & 1 \end{bmatrix}.$$

The purpose of this paper is to investigate the locally finite ergodic invariant measures of the flows F_θ^t on the surface $S(\mathcal{G}, \mathbf{w})$ (or equivalently of the interval exchanges T_θ). Our methods only work for particular directions θ .

Definition 1.1 (Renormalizable direction). We call a direction $\theta \in \mathbb{S}^1 \subset \mathbb{R}^2$ a λ -renormalizable direction if it satisfies the following two statements.

- (1) For all $\epsilon > 0$ there is a element g of the free group G for which $\|\rho_\lambda^g(\theta)\| < \epsilon$.
- (2) The vector θ is not an eigenvector of the matrix ρ_λ^g for any g in the conjugacy class of $hv^{-1} \in G$.

We denote the set of λ -renormalizable directions by $\mathcal{R}_\lambda$.

If $\mathbf{w}$ is a positive function satisfying $\mathbf{A}(\mathbf{w}) = \lambda \mathbf{w}$, then it must be that $\lambda \geq 2$. The 2-renormalizable directions are those vectors $\theta = (x, y) \in \mathbb{S}^1$ with $y/x \notin \mathbb{Q}$. For $\lambda > 2$, the λ -renormalizable directions form a Cantor set (the limit set of ρ_λ^G) with countably many points removed.

Our first major result is the following.

Theorem 1.2 (Topological conjugacy I). *Suppose $\mathbf{w}_1, \mathbf{w}_2 \in \mathbb{R}^\nu$ are positive functions satisfying $\mathbf{A}(\mathbf{w}_i) = \lambda_i \mathbf{w}_i$ for $i = 1, 2$. Let θ_1 be a λ_1 -renormalizable direction. Then there is a λ_2 -renormalizable direction θ_2 depending only on λ_1, λ_2 and θ_1 such that the interval exchange transformations $T_{\theta_1} : H(\mathcal{G}, \mathbf{w}_1) \rightarrow H(\mathcal{G}, \mathbf{w}_1)$ and $T_{\theta_2} : H(\mathcal{G}, \mathbf{w}_2) \rightarrow H(\mathcal{G}, \mathbf{w}_2)$ are topologically conjugate.*

We state a stronger version of this theorem in section 4. The topological conjugacy theorem allows us to construct alternate locally finite invariant measures for $T_{\theta_1} : H(\mathcal{G}, \mathbf{w}_1) \rightarrow H(\mathcal{G}, \mathbf{w}_1)$ by pulling back Lebesgue measure for $H(\mathcal{G}, \mathbf{w}_2)$ under the topological conjugacy.

Before discussing the ergodic measures, we need to further analyze the positive eigenvectors of the adjacency operator. For $\lambda > 0$, consider the set

$$E(\lambda) = \{\text{positive } \mathbf{f} \in \mathbb{R}^\nu \text{ satisfying } \mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}\}.$$

The set $E(\lambda) \cup \{\mathbf{0}\}$ is a convex closed cone in the topology of pointwise convergence. (The non-negative eigenfunctions are certainly closed. As $\mathcal{G}$ is connected, a non-negative eigenfunction must be positive.) We call a positive eigenfunction $f \in E(\lambda)$ *extremal* if $\mathbf{f} = \mathbf{f}_1 + \mathbf{f}_2$ for $\mathbf{f}_1, \mathbf{f}_2 \in E(\lambda)$ implies $\mathbf{f}_1 = c\mathbf{f}$ for some real number $c \in [0, 1]$.

The *valance* of a vertex v of a graph is the number of edges with v as an endpoint.

Theorem 1.3 (Ergodic measures I). *Assume $\mathcal{G}$, $\mathbf{w}_1$, and θ_1 are as in the Topological Conjugacy Theorem above. In addition assume that $\mathcal{G}$ has no vertices of valance one. Then the locally finite ergodic invariant measures for $T_{\theta_1} : H(\mathcal{G}, \mathbf{w}_1) \rightarrow H(\mathcal{G}, \mathbf{w}_1)$ are precisely the pullbacks of Lebesgue measure on $H(\mathcal{G}, \mathbf{w}_2)$ under the topological conjugacy given in the Topological Conjugacy Theorem, where $\mathbf{w}_2$ varies over the positive extremal eigenfunctions of $\mathbf{A}$.*

Remark 1.4 (Valance one condition). The author is willing to conjecture that the condition that $\mathcal{G}$ has no vertices of valance one can be removed from the above theorem. In fact, the arguments in this paper hold for all graphs satisfying the conditions of the Topological Conjugacy Theorem with the exception of those in section 14. In the remaining cases, this paper reduces the statements necessary to prove the same classification result to some combinatorial properties of the graph.

Remark 1.5 (Sharpness of renormalizable directions). If S were a compact translation surface with Veech group containing $\rho_\lambda^{\mathcal{G}}$, condition (1) of Definition 1.1 is sufficient to guarantee unique ergodicity of the flow in direction θ . Condition (2) is necessary for our renormalization argument, because there are surfaces for which the affine symmetry associated to the element $h\nu^{-1}$ are totally dissipative, as are the flows in the eigendirections of $\rho_\lambda^{h\nu^{-1}}$. In fact, for all $\lambda > 2$, if we choose $\mathbf{w}$ to be an extremal eigenfunction for the graph $\mathcal{G}_{\mathbb{Z}}$ with eigenvalue λ , the flow in each direction $\theta \in \mathbb{S}^1 \setminus \mathcal{R}_\lambda$ is totally dissipative.

It is worth remarking that the set of rays of extremal positive eigenfunctions with eigenvalue λ is also naturally a minimal Martin boundary of $\mathcal{G}$. See §6 of [MW89]. In particular, these objects have been well studied. We discuss the Martin boundary in §15.

Finally, we describe the case when $S(\mathcal{G}, \mathbf{w})$ has finite area. When $\mathbf{A}\mathbf{w} = \lambda\mathbf{w}$, the area of the surface $S(\mathcal{G}, \mathbf{w})$ turns out to be

$$\frac{\lambda}{2} \sum_{v \in \mathcal{V}} \mathbf{w}(v)^2.$$

So the surface has finite area if and only if $\mathbf{w} \in \ell^2(\mathcal{V})$. An example where this occurs is shown in figure 2. All the extremal positive eigenfunctions for this graph are described in example 15.10.

Similarly, the total length of $H(\mathcal{G}, \mathbf{w})$ is finite if and only if $\mathbf{w} \in \ell^1(\mathcal{V})$. We will apply the following theorem.




FIGURE 2. A surface $S(\mathcal{G}, \mathbf{w})$ with finite area. Labeled and opposite unlabeled edges are glued by horizontal or vertical translations. The graph $\mathcal{G}$ is shown with the positive eigenfunction $\mathbf{w} \in \ell^1(\mathcal{V})$ on the left.

Theorem 1.6 ([VJ68], Theorem 2.2; [Sen81], Theorem 6.4). *Let $\mathcal{G}$ be a infinite connected graph with bounded valance and suppose $\mathbf{w}$ is a positive eigenfunction in $\ell^2(\mathcal{V})$ with eigenvalue λ . Then,*

- (1) *All the other positive eigenfunction in $\ell^2(\mathcal{V})$ are scalar multiples of $\mathbf{w}$.*
- (2) *The spectral radius of $\mathbf{A}$ acting on $\ell^2(\mathcal{V})$ is λ .*

This combines with our classification of ergodic measures to yield the following.

Corollary 1.7. *If $H(\mathcal{G}, \mathbf{w})$ has finite total length, then for every λ -renormalizable direction θ , the interval exchange transformation $T_\theta : H(\mathcal{G}, \mathbf{w}) \rightarrow H(\mathcal{G}, \mathbf{w})$ is uniquely ergodic.*

This corollary follows by using the fact that $\ell^1(\mathcal{V})$ is a subset of $\ell^2(\mathcal{V})$ and by applying the two theorems above. This corollary applies to the surface in Figure 2, for instance.

1.3. Organization of Paper. In the next section, we discuss the structure of the graph $\mathcal{G}$, introduce the idea of a translation surface, and define the surface $S(\mathcal{G}, \mathbf{w})$. In section 3, we explain some of the structure of λ -renormalizable directions. In section 4, we use the notions introduced in the previous two sections to state stronger and clearer versions of the Topological Conjugacy and Ergodic Measures theorems. We discuss applications to skew rotations in section 5. We outline the proof of the Topological Conjugacy and Ergodic Measures theorems in section 6. Sections 7 through 14 contain the proofs of statements described in the outline. Section 15 describes relevant theorems about the minimal Martin boundaries (or equivalently the extremal positive eigenfunctions) of a graph $\mathcal{G}$. Finally, in section 16, we describe the interval exchange transformations that arise, and further discuss skew rotations.

2. BACKGROUND ON TRANSLATION SURFACES

2.1. Translation surfaces. A *translation surface* S is a (finite or countably infinite) collection of convex polygons $\{P_i \subset \mathbb{R}^2\}_{i \in \Lambda}$ with edges glued in pairs by translations. We call a translation surface *closed* if it is given as a finite union of polygons. Let V denote the collection of (equivalence classes of) vertices of polygons in S . Except on V , all points of S are locally isometric to $\mathbb{R}^2$. The points of V are cone points of S with cone angle $2k\pi$ for an integer $k \geq 1$, or points with infinite cone angle. We call a vertex $v \in V$ *removable* if it has cone angle 2π . Let $\Sigma \subset V$ denote the set of non-removable singularities.

Recall $F_\theta^t : S \rightarrow S$ denotes the straight-line flow in direction $\theta \in \mathbb{S}^1 \subset \mathbb{R}^2$. See equation 1.3. We let $\mathcal{F}_\theta$ denote the foliation of S by the orbits of the flow F_θ^t . The invariant measures on for the flow F_θ^t can be written as a product between a transverse measure for the foliation $\mathcal{F}_\theta$ and Lebesgue measure on the leaves. We use $\mathcal{M}_\theta$ to denote the collection of locally finite transverse measures to $\mathcal{F}_\theta$. We will be attempting to classify the ergodic measures in $\mathcal{M}_\theta$.

Remark 2.1 (Singular leaves). There is a standard method of dealing with leaves which intersect the non-removable singularities of S [KH95, §14.5]. This is to split the leaf in two, so that one of the new leaves tracks points to the right as they move around the singularity, and the other leaf tracks points to the left. We state these ideas more carefully in section 8.

2.2. The affine automorphism group. Translation surfaces inherit both a metric and a notion of direction from $\mathbb{R}^2$. The notion of direction is the pullback of the standard translation invariant fibration of the unit tangent bundle of $\mathbb{R}^2$ given by identifying $T_p^1\mathbb{R}^2$ with the unit circle $\mathbb{S}^1 \subset \mathbb{R}^2$ for each $p \in \mathbb{R}^2$. Thus, the notion of direction is a map

$dir : T^1S \rightarrow \mathbb{S}^1$. In particular, we consider two translation surfaces to be the same if there is a direction preserving isometry between them.

There is an action of $GL(2, \mathbb{R})$ on translation surfaces obtained by pulling back the action on $\mathbb{R}^2$. (This is the action on translation surfaces induced by the action of $GL(2, \mathbb{R})$ on polygons viewed as subsets of the plane.) The *Veech group* of S is the subgroup

$$\Gamma(S) = \{A \in GL(2, \mathbb{R}) : \text{there is a direction preserving isometry } \phi_A : A(S) \rightarrow S\}.$$

The *affine automorphism group* is the group

$$Aff(S) = \{\phi_A \circ A : S \rightarrow S : A \in \Gamma(S) \text{ and } \phi_A : A(S) \rightarrow S \text{ is a direction preserving isometry}\}.$$

The *derivative map* $D : Aff(S) \rightarrow \Gamma(S)$ is the map which sends $\phi_A \circ A$ to A .

2.3. Thurston's construction. Thurston used a variant of the construction we are about to describe to produce closed translation surfaces admitting pseudo-Anosov homeomorphisms [Thu88]. In fact, the closed translation surfaces admitting two non-commuting parabolic affine automorphisms can be characterized in terms of eigenvectors of graphs [Vee89]. We will follow the treatment of these ideas developed in [McM06]. See also [Hoo08a].

Definition 2.2 (Graph terminology). Throughout this paper, $\mathcal{G}$ is an infinite, connected, bipartite, ribbon graph with bounded valance. These terms are defined below.

- (1) (*Infinite*) The vertex set $\mathcal{V}$ is countably infinite.
- (2) (*Connected*) For every $v, w \in \mathcal{V}$, there is a sequence of vertices $v = v_0, v_1, \dots, v_k = w$ so that every $\overline{v_i v_{i+1}}$ lies in the edge set $\mathcal{E}$.
- (3) (*Bipartite*) The vertex set $\mathcal{V}$ decomposes into a disjoint union of two sets, $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$, and the edge set $\mathcal{E}$ consists only of edges of the form $\overline{ab}$ with $a \in \mathcal{A}$ and $b \in \mathcal{B}$. Thus we have natural maps $\alpha : \mathcal{E} \rightarrow \mathcal{A}$ and $\beta : \mathcal{E} \rightarrow \mathcal{B}$ given by the maps

$$\alpha : \overline{ab} \mapsto a \quad \text{and} \quad \beta : \overline{ab} \mapsto b.$$

- (4) (*Bounded valance*) The sets $\alpha^{-1}(a)$ and $\beta^{-1}(b)$ are finite sets whose sizes are bounded from above.
- (5) (*Ribbon structure*) For every $v \in \mathcal{V}$, the ribbon graph structure specifies a cyclic permutation P_v of the edges that contain v as an endpoint.

Definition 2.3 (East and North edge permutations). Given the above structure on $\mathcal{G}$, we define bijections $\mathcal{E}, \mathcal{N} : \mathcal{E} \rightarrow \mathcal{E}$. These are given by

$$\mathcal{E}(\overline{ab}) = P_a(\overline{ab}) \quad \text{and} \quad \mathcal{N}(\overline{ab}) = P_b(\overline{ab}).$$

Since the permutations P_v are cyclic, these maps satisfy

$$(2.1) \quad \{\mathcal{E}^k(e)\}_{k \in \mathbb{N}} = \alpha^{-1}(\alpha(e)) \quad \text{and} \quad \{\mathcal{N}^k(e)\}_{k \in \mathbb{N}} = \beta^{-1}(\beta(e)).$$

Definition 2.4 ($S(\mathcal{G}, \mathbf{w})$). Let $\mathcal{G}$ be a infinite, connected, bipartite, ribbon graph with bounded valance, and let $\mathbf{w} \in \mathbb{R}^{\mathcal{V}}$ be a positive function. Using this data, we will construct a translation surface $S(\mathcal{G}, \mathbf{w})$. This surface will be a union of rectangles R_e with $e \in \mathcal{E}$, with each R_e given by

$$R_e = [0, \mathbf{w} \circ \beta(e)] \times [0, \mathbf{w} \circ \alpha(e)].$$

We glue the rectangles so that the right (east) side of R_e is glued isometrically to the left side of $R_{\mathcal{E}(e)}$, and the top (north) side of R_e is glued isometrically to the bottom of $R_{\mathcal{N}(e)}$. (This explains the notation for the bijections $\mathcal{E}, \mathcal{N} : \mathcal{E} \rightarrow \mathcal{E}$.)

Note that this surface admits horizontal and vertical cylinder decompositions, which intersect in the given rectangles. The horizontal cylinders are in bijective correspondence with the set $\mathcal{A} \subset \mathcal{V}$. Namely, the horizontal and vertical cylinders corresponding to vertices $a \in \mathcal{A}$ and $b \in \mathcal{B}$ are the sets

$$(2.2) \quad cyl_a = \bigcup_{e \in \alpha^{-1}(a)} R_e \quad \text{and} \quad cyl_b = \bigcup_{e \in \beta^{-1}(b)} R_e.$$

The modulus of a cylinder is the ratio $\frac{\text{width}}{\text{circumference}}$. Therefore, the condition that all horizontal and vertical cylinders have equal moduli is equivalent to saying that f is an eigenfunction of the adjacency operator. In particular, by a theorem of Veech [Vee89] this guarantees the existence of two non-commuting parabolic automorphisms of our surface. Thus, we have the following.

Proposition 2.5. *Suppose w is a positive eigenfunction for the adjacency operator corresponding to an eigenvalue $\lambda > 0$. Then, there is an embedding of the free group on two generators into the affine automorphism group, $\Phi : G \rightarrow \text{Aff}(S(\mathcal{G}, \mathbf{w}))$, so that Φ^h and Φ^v act by single Dehn twists in the horizontal and vertical cylinder decompositions respectively. Their derivatives are given by*

$$D\Phi^h = \rho_\lambda^h = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad D\Phi^v = \rho_\lambda^v = \begin{bmatrix} 1 & 0 \\ \lambda & 1 \end{bmatrix}.$$

The main idea of this paper is to use this automorphism group to renormalize the straight-line flows on our surfaces.

Remark 2.6 (Dihedral group action). The dihedral group of order eight generated by

$$\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \quad \text{and} \quad \begin{bmatrix} -1 & 0 \\ 0 & 1 \end{bmatrix}$$

acts on the collection of surfaces obtainable from Thurston's construction. If the matrix A lies in this group, the surface $A(S(\mathcal{G}, \mathbf{w})) = S(\mathcal{G}', \mathbf{w})$ with $\mathcal{G}'$ arising from $\mathcal{G}$ by changing the bipartite and ribbon graph structure on $\mathcal{G}$ in a way that depends on A . Note that this matrix group also acts on directions, and on the group ρ_λ^G by conjugation. We will frequently use this dihedral group action to simplify proofs in this paper.

3. RENORMALIZABLE DIRECTIONS

In this section, we revisit the definition of a λ -renormalizable direction $\theta \in \mathbb{S}^1$. We will state results in this section without proofs. These results will be proved in section 7, unless otherwise noted.

Throughout this paper, we will be concerned with actions of the free group on two letters. As in the introduction, we denote this group by $G = \langle h, v \rangle$. We will equip G with the word metric, making G isometric to the nodes of the 4-valent tree. The following is equivalent to the metric definition of geodesic.

Definition 3.1 (Geodesics in G). An infinite (resp. finite) sequence $\langle g_0, g_1, \dots \rangle$ of elements of G is a *geodesic ray* (resp. *segment*) in G if it satisfies the following statements.

1. $g_{n+1}g_n^{-1} \in \{h, v, h^{-1}, v^{-1}\}$ for all $n \geq 0$.
2. $g_{n+2}g_n^{-1} \neq e$ for all $n \geq 0$.

In this paper, all geodesic rays and segments in G will be based at the identity, unless otherwise stated. This means that

3. $g_0 = e$, the identity.

The “symmetries” of our systems will be determined by a constant $\lambda \geq 2$. For such a λ we define the representation $\rho_\lambda : G \rightarrow SL(2, \mathbb{R})$ by

$$(3.1) \quad \rho_\lambda^h = \begin{bmatrix} 1 & \lambda \\ 0 & 1 \end{bmatrix} \quad \text{and} \quad \rho_\lambda^v = \begin{bmatrix} 1 & 0 \\ \lambda & 1 \end{bmatrix}.$$

The matrices ρ_λ^h and ρ_λ^v are parabolics with horizontal and vertical eigenvectors, respectively. Because $\lambda \geq 2$, the representations ρ_λ are always discrete. Note that ρ_2 is a bijection onto the congruence two subgroup of $SL(2, \mathbb{Z})$, which is a lattice.

Definition 3.2 (Shrinkable). We call a direction $\theta \in \mathbb{S}^1$ λ -*shrinkable* if for all $\epsilon > 0$ there is a $g \in G$ such that $\|\rho_\lambda^g(\theta)\| < \epsilon$.

Definition 3.3 (Shrinking sequence). A geodesic ray $\langle g_i \rangle$ is a λ -*shrinking sequence* of θ if for all $\epsilon > 0$ there is an $i > 0$ for which $\|\rho_\lambda^{g_i}(\theta)\| < \epsilon$. We say $\langle g_i \rangle$ is a λ -*shrinking sequence* if it is the λ -shrinking sequence of some $\theta \in \mathbb{S}^1$.

Theorem 3.4. *If $\theta \in \mathbb{S}^1$ is λ -shrinkable, then there is a unique geodesic ray $\langle g_i \rangle$ which is a λ -shrinking sequence of θ . Moreover, $\|\rho_\lambda^{g_i}(\theta)\|$ is monotonically decreasing.*

By uniqueness, we will now say $\langle g_i \rangle$ is the λ -shrinking sequence of θ .

Proposition 3.5. *If the geodesic ray $\langle g_i \rangle$ is a λ -shrinking sequence, then it is the λ -shrinking sequence of a unique pair of anti-podal vectors in $\mathbb{S}^1$.*

We define the following shift action on geodesic rays in G ,

$$S(\langle g_0, g_1, g_2, \dots \rangle) = \langle g_1 g_0^{-1}, g_2 g_1^{-1}, g_3 g_2^{-1}, \dots \rangle.$$

Note that S sends the λ -shrinking sequence of θ to the λ -shrinking sequence for $\rho_\lambda^{g_1}(\theta)$. We call two sequences of elements in G $\langle g_i \rangle$ and $\langle g'_i \rangle$ *tail equivalent* if there are non-negative integers m and n for which $S^m(\langle g_i \rangle) = S^n(\langle g'_i \rangle)$.

The following characterizes λ -shrinking sequences.

Theorem 3.6 (Characterization of shrinking sequences). *Consider the following conditions on a geodesic ray $\langle g_i \rangle$ in G .*

(1) $\langle g_i \rangle$ is not tail equivalent to any of the four sequences fixed by S ,

$$\langle e, h, h^2, \dots \rangle, \quad \langle e, h^{-1}, h^{-2}, \dots \rangle, \quad \langle e, v, v^2, \dots \rangle \quad \text{and} \quad \langle e, v^{-1}, v^{-2}, \dots \rangle.$$

(2) $\langle g_i \rangle$ is not tail equivalent to either of the following sequences of period two under S

$$\langle e, h, v^{-1}h, hv^{-1}h, v^{-1}hv^{-1}h, \dots \rangle \quad \text{and} \quad \langle e, h^{-1}, vh^{-1}, h^{-1}vh^{-1}, vh^{-1}vh^{-1}, \dots \rangle.$$

For $\lambda > 2$, a geodesic ray is a λ -shrinking sequence if and only if it satisfies condition (1). For $\lambda = 2$, a geodesic ray is a λ -shrinking sequence if and only if it satisfies conditions (1) and (2).

Definition 3.7 (Renormalizable directions). A geodesic ray $\langle g_i \rangle$ satisfying conditions (1) and (2) of Theorem 3.6 is called a *renormalizing sequence*. A direction $\theta \in \mathbb{S}^1$ is λ -*renormalizable* if it is λ -shrinkable and its λ -shrinking sequence is renormalizing. We use $\mathcal{R}_\lambda \subset \mathbb{S}^1$ to denote the set of all λ -renormalizable vectors in $\mathbb{S}^1$. If $\langle g_n \rangle$ is a renormalizing sequence and $\lambda \geq 2$,

then there is a unique pair of anti-podal vectors in $\mathcal{R}_\lambda$ whose λ -shrinking sequence is $\langle g_n \rangle$. We use $\pm \boldsymbol{\theta}(\langle g_n \rangle, \lambda) \in \mathcal{R}_\lambda$ to denote this pair.

Proposition 3.8. *If $\langle g_n \rangle$ is a renormalizing sequence, the pair of quadrants in $\mathbb{R}^2$ containing the vectors $\pm \boldsymbol{\theta}(\langle g_n \rangle, \lambda)$ is independent of the choice of $\lambda \geq 2$.*

Remark 3.9 (Equivalence of definitions). Theorem 3.6 together with Proposition 3.5 implies that the definition of λ -renormalizable directions given in the introduction is equivalent to the one given here.

Remark 3.10 (The limit set). The set of λ -renormalizable directions is closely connected to the limit set of the induced action of ρ_λ^G on $\mathbb{RP}^1$. In fact, $\mathcal{R}_\lambda$ projectivizes to the usual limit set of a group action on $\mathbb{RP}^1$ with a finite number of orbits removed. See Proposition 7.14.

4. RESTATEMENT OF MAIN RESULTS

In this section, we restate our main results from the introduction. These new statements reflect the point of view of the proofs.

If S is a translation surface and $\boldsymbol{\theta} \in \mathbb{S}^1$, then we can consider the foliation $\mathcal{F}_\boldsymbol{\theta}$ of the surface S by orbits of the straight-line flow $F_\boldsymbol{\theta}^t$ of equation 1.3. We state a stronger version of the Topological conjugacy theorem of the introduction using these foliations.

Theorem 4.1 (Topological conjugacy II). *Suppose $\mathbf{w}_1, \mathbf{w}_2 \in \mathbb{R}^\vee$ are positive functions satisfying $\mathbf{A}(\mathbf{w}_i) = \lambda_i \mathbf{w}_i$ for $i = 1, 2$. Let $\langle g_n \rangle$ be a renormalizing sequence, and let $\pm \boldsymbol{\theta}_i = \pm \boldsymbol{\theta}(\langle g_n \rangle, \lambda_i)$ be associated pairs of anti-podal λ_i -renormalizable directions. (See Definition 3.7.) Then there is a homeomorphism $\phi : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$ such that $\phi(\mathcal{F}_{\boldsymbol{\theta}_1}) = \mathcal{F}_{\boldsymbol{\theta}_2}$. Moreover, ϕ can be taken to preserve the decomposition of the surfaces into labeled rectangles as in Definition 2.4,*

$$S(\mathcal{G}, \mathbf{w}_1) = \bigcup_{e \in \mathcal{E}} R_e^1 \quad \text{and} \quad S(\mathcal{G}, \mathbf{w}_2) = \bigcup_{e \in \mathcal{E}} R_e^2,$$

and so that the restricted maps $\phi|_{R_e^1} : R_e^1 \rightarrow R_e^2$ sends the bottom (resp. top, left, right) edge of R_e^1 to the bottom (resp. top, left, right) edge of R_e^2 for all $e \in \mathcal{E}$. In this case, the restriction of ϕ to a map from $\bigcup_{e \in \mathcal{E}} \partial R_e^1$ to $\bigcup_{e \in \mathcal{E}} \partial R_e^2$ is uniquely determined.

Remark 4.2. The restriction of ϕ to the horizontal edges of rectangles gives the topological conjugacy mentioned in theorem 1.2 of the introduction.

Again let S be a translation surface and let $\boldsymbol{\theta} \in \mathbb{S}^1$. Then we can consider the space of locally finite transverse measures to the foliation $\mathcal{F}_\boldsymbol{\theta}$. A *locally finite $\mathcal{F}_\boldsymbol{\theta}$ -transverse measure* is one which assigns finite measure to every compact transversal to $\mathcal{F}_\boldsymbol{\theta}$. We also insist that these measures have no atoms supported on singular leaves. (Singular leaves are trajectories of the straight-line flow which hit singularities in forward or backward time, and therefore are not defined for all time.) An example of such a measure is the *Lebesgue $\mathcal{F}_\boldsymbol{\theta}$ -transverse measure* on S . If $\gamma : [0, 1] \rightarrow S$ is a differentiable transversal path, the Lebesgue $\mathcal{F}_\boldsymbol{\theta}$ -transverse measure satisfies

$$\gamma \mapsto \int_t |\gamma'(t) \wedge \boldsymbol{\theta}| \, dt,$$

where $\wedge$ denotes the usual wedge product between vectors in the plane;

$$(4.1) \quad \wedge : \mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R} \quad \text{defined by} \quad (a, b) \wedge (c, d) = ad - bc.$$

Theorem 4.3 (Ergodic measures II). *Assume $\mathcal{G}$, $\mathbf{w}_1$, and $\boldsymbol{\theta}_1$ are as in theorem 4.1. Additionally assume $\mathcal{G}$ has no vertices of valance one. Then the locally finite ergodic transverse measures to $\mathcal{F}_{\boldsymbol{\theta}_1}$ are the pullbacks of the Lebesgue $\mathcal{F}_{\boldsymbol{\theta}_2}$ -transverse measure on $S(\mathcal{G}, \mathbf{w}_2)$ under the homeomorphisms ϕ given in theorem 4.1, where $\mathbf{w}_2$ varies over the extremal positive eigenfunctions of $\mathbf{A}$.*

5. APPLICATIONS TO SKEW ROTATIONS

In this section, we briefly describe applications of our main results to skew rotations. Proofs and further discussion are given in section 16. Section 16 relies to a large extent on known results on the minimal Martin boundary, which we discuss in section 15.

Recall the definition of the skew rotation $T : \mathbb{R}/\mathbb{Z} \times \mathbb{G} \rightarrow \mathbb{R}/\mathbb{Z} \times \mathbb{G}$ given in equation 1.1. This transformation is determined by a choice of generators $\gamma_1, \dots, \gamma_n$ so that $\gamma_n \dots \gamma_1 = e$, and an $\alpha \in \mathbb{R}$. In section 16.2, we discuss a suspension of the skew rotation T . In fact, we give a construction of a valance n infinite bipartite ribbon graph $\mathcal{G}$ so that the skew rotation T is naturally the return map of the the straight line flow on $S(\mathcal{G}, \mathbf{w}_{\frac{1}{n}})$ in direction $\boldsymbol{\theta} = (\alpha - \frac{1}{n}, \frac{1}{n}) / \|(\alpha - \frac{1}{n}, \frac{1}{n})\|$ to the horizontal edges of the decomposition of $S(\mathcal{G}, \mathbf{w}_{\frac{1}{n}})$. Here the function $\mathbf{w}_{\frac{1}{n}} \in \mathbb{R}^{\mathcal{V}}$ is the constant function with value $\frac{1}{n}$. This satisfies $\mathbf{A}(\mathbf{w}_{\frac{1}{n}}) = n\mathbf{w}_{\frac{1}{n}}$.

Recall the definition of Ω_n given in equation 1.2.

Proposition 5.1. *$\boldsymbol{\theta}$ is n -renormalizable if and only if $\alpha \in \Omega_n$.*

The graph $\mathcal{G}$ is not a Cayley graph of $\mathbb{G}$. However, the vertices in $\mathcal{A} \subset \mathcal{V}$ are naturally labeled by the elements of the group. We construct the graph $\mathcal{G}_{\mathcal{A}}$ which has vertex set $\mathcal{A} = \mathbb{G}$ and for which an edge is joined between g and g' for any path of length two in $\mathcal{G}$ which joins $g \in \mathcal{A}$ to $g' \in \mathcal{A}$. We say a *generalized Cayley graph* is one where the list of generators is allowed to include duplicate elements and the identity element.

Proposition 5.2. *The graph $\mathcal{G}_{\mathcal{A}}$ is a generalized Cayley graph. Moreover, the restriction map $\mathbb{R}^{\mathcal{V}} \rightarrow \mathbb{R}^{\mathcal{A}}$ sends the positive eigenfunctions of the adjacency operator of $\mathcal{G}$ with eigenvalue λ to the positive eigenfunctions of the adjacency operator of $\mathcal{G}_{\mathcal{A}}$ with eigenvalue λ^2 .*

Therefore, we have the following Corollary to Theorem 4.3.

Corollary 5.3. *Suppose $\alpha \in \Omega_n$. Then if ν is a locally finite ergodic invariant measures of the skew rotation T , then the function $\mathbf{f}_{\nu} : \mathbb{G} \rightarrow \mathbb{R}$ given by*

$$\mathbf{f}_{\nu}(g) = \nu(\mathbb{R}/\mathbb{Z} \times \{g\})$$

is an extremal positive eigenfunction of the Cayley graph $\mathcal{G}_{\mathcal{A}}$. Moreover, the map $\nu \mapsto \mathbf{f}_{\nu}$ is a bijection between the space of locally finite ergodic invariant measures of T and the extremal positive eigenfunctions of $\mathcal{G}_{\mathcal{A}}$.

The positive eigenfunctions of the adjacency operator acting on a Cayley graph have been well studied. As mentioned in the introduction, we can combine the results of this paper with theorems of this nature to produce information about the spaces of locally finite ergodic invariant measures of T in special cases. From a result of Margulis [Mar66], which we restate as Theorem 15.13, we have the following.

Corollary 5.4. *Suppose $\mathcal{G}$ is nilpotent and $\alpha \in \Omega_n$. Then space of locally finite ergodic invariant measures is homeomorphic to the cone on the collection of group homomorphisms*

$\mathbb{G} \rightarrow \mathbb{R}_\times$. In particular, Lebesgue measure is the ergodic measure corresponding to the trivial group homomorphisms, up to scaling.

In contrast, suppose $n > 2$ and $\mathbb{G}$ is the free group

$$(5.1) \quad \mathbb{G} = \langle \gamma_1, \dots, \gamma_n \mid \gamma_n \dots \gamma_2 \gamma_1 = e \rangle.$$

Then $\mathcal{G}$ turns out to be the regular n -tree. This graph has spectral radius $2\sqrt{n-1}$. A theorem of Picardello and Woess [PW88], which we restate in section 15.2, gives the following.

Corollary 5.5. *Let $\alpha \in \Omega_n$ and $\mathbb{G}$ be as in equation 5.1. Then the space of locally invariant measures of T is in bijective correspondence with $[2\sqrt{n-1}, \infty) \times \partial G$, where $[2\sqrt{n-1}, \infty)$ denotes the collection of possible eigenvalues and ∂G is the space of ends of the tree. In particular, Lebesgue measure is not ergodic.*

6. OUTLINE OF ARGUMENTS

6.1. Reinterpretation of invariant measures. Let S be an infinite translation surface and let V be a non-empty set of (possibly removable) singularities of S . Let $\theta \in \mathbb{S}^1$ be a direction and $\mathcal{F}_\theta$ be the oriented foliation by orbits of the straight line flow in direction θ . Let $\mathcal{M}_\theta$ denote the set of all locally finite transverse measures for $\mathcal{F}_\theta$ with no atoms supported on singular leaves.

Let $H_1(S, V, \mathbb{R})$ denote the real homology classes of closed curves in S/V (ie. S with the points in V collapsed to a single point). We will always take the cohomology group $H^1(S, V, \mathbb{R})$ to mean the dual space to $H_1(S, V, \mathbb{R})$. That is, $H^1(S, V, \mathbb{R})$ is the collection of all linear maps from $H_1(S, V, \mathbb{R}) \rightarrow \mathbb{R}$.

Definition 6.1 (The cohomology class of a measure). We can think of a transverse measure $\mu \in \mathcal{M}_\theta$ as determining a linear map $H_1(S, V, \mathbb{Z}) \rightarrow \mathbb{R}$. This map should just be $\llbracket x \rrbracket \mapsto \mu(x)$ if x is a curve in S joining a point in V to a point in V and everywhere crossing the foliation $\mathcal{F}_\theta$ with positive algebraic sign. (To be precise, if $\mathcal{F}_\theta$ is upward pointing, transversals pointed right cross $\mathcal{F}_\theta$ with positive algebraic sign.) This map can be uniquely extended to a linear map $H_1(S, V, \mathbb{Z}) \rightarrow \mathbb{R}$, since the map is determined on a basis. We denote this linear map by

$$\Psi_\theta : \mathcal{M}_\theta \rightarrow H^1(S, V, \mathbb{R}).$$

Recall F_θ^t denoted the flow on a translation surface in direction θ . We call the flow $F_\theta^t : S \rightarrow S$ *conservative* if given any subset $A \subset S$ of positive measure and any $T > 0$, for Lebesgue a.e. $x \in S$ there is a $t > T$ for which $F_\theta^t(x) \in A$. We have the following.

Lemma 6.2. *Let S be a translation surface with non-empty singular set V . The map $\Psi_\theta : \mathcal{M}_\theta \rightarrow H^1(S, V, \mathbb{R})$ is injective if the straight-line flow F_θ^t has no periodic trajectories and is conservative.*

This statement is proved in section 8. We can apply this result because we have the following.

Theorem 6.3. *Let $S(\mathcal{G}, \mathbf{w})$ be a the translation surface constructed as in Definition 2.4, with $\mathbf{w} \in \mathbb{R}^\mathcal{V}$ a positive eigenfunction for the adjacency operator with eigenvalue λ . Then if the direction θ is λ -renormalizable, then the flow F_θ^t on $S(\mathcal{G}, \mathbf{w})$ has no periodic trajectories and is conservative.*

So, by the lemma above, we have injectivity of Ψ_θ in λ -renormalizable directions. This theorem is proved in section 9.

A *saddle connection* is an oriented geodesic segment $\sigma \subset S$ which intersects V precisely at its endpoints. The *holonomy* $hol(\sigma) \in \mathbb{R}^2$ is the vector difference between the end and starting points of a developed image of σ into the plane. The function $\text{sgn} : \mathbb{R} \rightarrow \{-1, 0, 1\}$ assigns to a real number the element of $\{-1, 0, 1\}$ with the same sign.

Lemma 6.4. *Given $m \in H^1(S, V, \mathbb{R})$, we have $m \in \Psi_\theta(\mathcal{M}_\theta)$ if and only if for all saddle connections σ with $m(\llbracket \sigma \rrbracket) \neq 0$ we have*

$$\text{sgn}(m(\llbracket \sigma \rrbracket)) = \text{sgn}(hol(\sigma) \wedge \theta).$$

The proof is in §8.3.

Remark 6.5 (Awkward inequalities). The construct “ $a \neq 0$ implies $\text{sgn}(a) = \text{sgn}(b)$ ” is equivalent to the longer statement “if $b > 0$ then $a \geq 0$ and if $b < 0$ then $a \leq 0$.” We will repeatedly use this more compact construct in this paper.

6.2. Action of the affine automorphism group. For the remainder of this section 6, we will let $S = S(\mathcal{G}, \mathbf{w})$ be a translation surface as constructed in section 2.3. Let E denote the set of horizontal and vertical edges of the rectangles making up $S(\mathcal{G}, \mathbf{w})$, and V denotes the set of vertices of rectangles. By definition, the edges in E are all saddle connections, since they join a point in V to a point in V . We orient the horizontal saddle connections in E rightward, and the vertical saddle connections upward. We have the following.

Proposition 6.6. *The homology classes of the saddle connections in E generate $H_1(S, V, \mathbb{Z})$.*

The proof is to simply note that by cutting along the horizontal and vertical saddle connections, we decompose S into rectangles. In particular, an $m \in H^1(S, V, \mathbb{R})$ is determined by what it does to E .

Let $\phi \in \text{Aff}(S)$. Then, ϕ acts on the space of transverse measures by pushing forward the measure. For all $\theta \in \mathbb{S}^1$, ϕ induces a bijection

$$\phi_* : \mathcal{M}_\theta \rightarrow \mathcal{M}_{D(\phi)(\theta)} : \mu \mapsto \mu \circ \phi^{-1}.$$

Now suppose that ϕ preserves the singular set V . (It may not, since we allow removable singularities.) Then, this action on measures is compatible with the pushforward action on $H^1(S, V, \mathbb{R})$. We abuse notation by also denoting this pushforward by $\phi_* : H^1(S, V, \mathbb{R}) \rightarrow H^1(S, V, \mathbb{R})$. For $m \in H^1(S, V, \mathbb{R})$, we define

$$(6.1) \quad (\phi_*(m))(\llbracket x \rrbracket) = m \circ \phi^{-1}(\llbracket x \rrbracket)$$

for all $\llbracket x \rrbracket \in H_1(S, V, \mathbb{Z})$. For all $\mu \in \mathcal{M}_\theta$ we have

$$\phi_* \circ \Psi_\theta(\mu) = \Psi_{D(\phi)(\theta)} \circ \phi_*(\mu).$$

We now recall proposition 2.5. If $\mathbf{w}$ is a positive eigenfunction for $\mathcal{G}$ with eigenvalue λ , we have an action Φ of the group G by affine automorphisms of $S(\mathcal{G}, \mathbf{w})$. This group action preserves the vertex set V . The action induced on homology is independent of the choice of such a positive eigenfunction $\mathbf{w}$. The following explains the action on the generating set E of $H_1(S, V, \mathbb{Z})$.

Proposition 6.7 (Action on homology). *Given any $\sigma \in E \subset H_1(S, V, \mathbb{Z})$, we have*

$$\begin{aligned} \Phi^{h^k}(\llbracket \sigma \rrbracket) &= \begin{cases} \llbracket \sigma \rrbracket & \text{if } \sigma \text{ is horizontal} \\ \llbracket \sigma \rrbracket + k \llbracket cyl_a \rrbracket & \text{if } \sigma \text{ is vertical and traverses } cyl_a \text{ with } a \in \mathcal{A} \end{cases} \\ \Phi^{v^k}(\llbracket \sigma \rrbracket) &= \begin{cases} \llbracket \sigma \rrbracket + k \llbracket cyl_b \rrbracket & \text{if } \sigma \text{ is horizontal and traverses } cyl_b \text{ with } b \in \mathcal{B} \\ \llbracket \sigma \rrbracket & \text{if } \sigma \text{ is vertical} \end{cases} \end{aligned}$$

Recall, a λ -renormalizable direction θ has a renormalizing λ -shrinking sequence of elements of G , $\langle g_i \rangle$. See definition 3.7.

Definition 6.8. We say that an $m \in H^1(S, V, \mathbb{R})$ is a (θ, n) -survivor if for all saddle connections $\sigma \in \Phi^{g_n^{-1}}(E)$ we have $m(\llbracket \sigma \rrbracket) \neq 0$ implies

$$\text{sgn}(m(\llbracket \sigma \rrbracket)) = \text{sgn}(\text{hol}(\sigma) \wedge \theta).$$

The collection of (θ, n) -survivors $m \in H^1(S, V, \mathbb{R})$ is a convex cone by remark 6.5. Note that, this condition verifies lemma 6.4 on a subset of saddle connections depending on n . We have the following motivation for our definition.

Theorem 6.9 (Survivors and measures). *Let θ be a λ -renormalizable direction. For any $m \in H^1(S, V, \mathbb{R})$, $m \in \Psi_\theta(\mathcal{M}_\theta)$ if and only if m is a (θ, n) -survivor for all $n \geq 0$.*

The proof follows from analyzing the action of the sequence $\rho_\lambda^{g_n}$ on $\mathbb{S}^1$. We show that if m is a (θ, n) -survivor for all n , then m satisfies Lemma 6.4. See the end of section 10.

We can derive an equivalent definition of being a (θ, n) -survivor by acting by Φ^{g_n} . We have the following.

Proposition 6.10. *The condition that m be a (θ, n) -survivor is equivalent to the statement that for all $\sigma \in E$, we have $(\Phi_*^{g_n}(m))(\llbracket \sigma \rrbracket) \neq 0$ implies*

$$\text{sgn}(\Phi_*^{g_n}(m))(\llbracket \sigma \rrbracket) = \text{sgn}(\text{hol}(\sigma) \wedge \rho_\lambda^{g_n}(\theta)).$$

Remark 6.11 (Renormalization argument). The interpretation of being a (θ, n) -survivor provided in by Proposition 6.10 allows for a renormalization argument in the following sense. The conditions on being a $(\theta, 0)$ -survivor are relatively weak, and depend only on the quadrant containing θ . The condition that m be a (θ, n) -survivor is equivalent to $\Phi_*^{g_n}(m)$ being a $(\rho_\lambda^{g_n}(\theta), 0)$ -survivor. By theorem 6.9, the cohomology classes associated to invariant measures are precisely those for which $\Phi_*^{g_n}(m)$ is a $(\rho_\lambda^{g_n}(\theta), 0)$ -survivor for all n . In that sense, we are utilizing the action of the shrinking sequence $\langle g_n \in G \rangle$ on the space of straight-line flows of $S(\mathcal{G}, \mathbf{w})$ to understand the invariant measures for these flows.

6.3. Operators on graphs. Recall, $\mathbb{R}^\mathcal{V}$ denotes the space of all functions $\mathcal{V} \rightarrow \mathbb{R}$. In addition to the adjacency operator defined in equation 1.4, we will be studying the following linear operators on $\mathbb{R}^\mathcal{V}$.

$$(6.2) \quad \mathbf{H}^k(\mathbf{f})(x) = \begin{cases} \mathbf{f}(x) + k \sum_{y \sim x} \mathbf{f}(y) & \text{if } x \in \mathcal{A} \\ \mathbf{f}(x) & \text{if } x \in \mathcal{B}. \end{cases}$$

$$(6.3) \quad \mathbf{V}^k(\mathbf{f})(x) = \begin{cases} \mathbf{f}(x) & \text{if } x \in \mathcal{A} \\ \mathbf{f}(x) + k \sum_{y \sim x} \mathbf{f}(y) & \text{if } x \in \mathcal{B}. \end{cases}$$

We define the group action $\Upsilon : G \times \mathbb{R}^\mathcal{V} \rightarrow \mathbb{R}^\mathcal{V}$ by extending the definition $\Upsilon^h = \mathbf{H}$ and $\Upsilon^v = \mathbf{V}$. As long as $\mathcal{G}$ is an infinite connected graph, the group action Υ is a faithful action of the free group with two generators. The operators relate to the adjacency operator, $\mathbf{A}$, by the equations

$$(6.4) \quad \mathbf{A}\mathbf{H} = \mathbf{V}\mathbf{A} \quad \text{and} \quad \mathbf{A}\mathbf{V} = \mathbf{H}\mathbf{A}.$$

There is a natural linear embedding $\Xi : \mathbb{R}^\mathcal{V} \rightarrow H^1(S, V, \mathbb{R})$. Given $\mathbf{f} \in \mathbb{R}^\mathcal{V}$, and $\llbracket x \rrbracket \in H_1(S, V, \mathbb{Z})$ we define

$$(6.5) \quad \Xi(\mathbf{f})(\llbracket x \rrbracket) = \sum_{v \in \mathcal{V}} i(\llbracket x \rrbracket, \llbracket cyl_v \rrbracket) \mathbf{f}(v).$$

Here, $i : H_1(S, V, \mathbb{Z}) \times H_1(S \setminus V, \mathbb{Z}) \rightarrow \mathbb{Z}$ denotes the usual algebraic intersection number, and $\llbracket cyl_v \rrbracket$ is the homology class of a core curve of the cylinder cyl_v defined in equation 2.2 oriented rightward or upward. This sum is well defined, because $i(\llbracket x \rrbracket, \llbracket cyl_v \rrbracket) \neq 0$ for only finitely many $v \in \mathcal{V}$.

It is not difficult to see that the image $\Xi(\mathbb{R}^\mathcal{V})$ is invariant under the action of the affine automorphism group. In fact, we have the following.

Proposition 6.12. *For all $g \in G$ we have $\Phi_*^g \circ \Xi = \Xi \circ \Upsilon^g$.*

This proposition is proved in section 11.

Suppose that $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a positive function. Then we can think of $\Xi(\mathbf{f})$ as the element of $H^1(S, V, \mathbb{R})$ arising from the Lebesgue transverse measure to the foliation in direction of angle $\frac{\pi}{4}$ on the surface $S(\mathcal{G}, \mathbf{f})$. Similar interpretations arise from considering an $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ where the signs of $\mathbf{f}$ on the subsets of vertices $\mathcal{A}$ and $\mathcal{B}$ are fixed. This idea is what gives rise to the notions of quadrants, which we are about to introduce. We will show that $\Psi_\theta(\mathcal{M}_\theta) \subset \Xi(\mathbb{R}^\mathcal{V})$. Moreover, we will show that the set of transverse measures to the foliation $\mathcal{F}_\theta$ all can be interpreted as such Lebesgue transverse measures on $S(\mathcal{G}, \mathbf{f})$ with $\mathbf{f}$ satisfying “survivor” conditions.

Definition 6.13 (Survivors in $\mathbb{R}^\mathcal{V}$). Assume $\theta \in \mathbb{S}^1$ is a λ -renormalizable direction with λ -shrinking sequence $\langle g_i \rangle$. We say that $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a (θ, n) -survivor if $\Xi(\mathbf{f})$ is a (θ, n) -survivor. We say $\mathbf{f}$ is a θ -survivor if it is a (θ, n) -survivor for all $n \geq 0$. We use $\mathcal{S}_\theta \subset \mathbb{R}^\mathcal{V}$ to denote the set of all θ -survivors.

We will introduce a more intrinsic equivalent definition. To do this, we need a few more definitions.

Definition 6.14 (Sign pairs). The set of sign pairs is the set of four elements $SP = \{(\pm 1, \pm 1)\}$. We abbreviate these elements by writing

$$++ = (1, 1) \quad +- = (1, -1) \quad -+ = (-1, 1) \quad -- = (-1, -1).$$

We will use π_1 and π_2 to denote the projection functions. For instance $\pi_1(-+) = -1$ and $\pi_2(-+) = 1$.

Definition 6.15 (Quadrants in $\mathbb{R}^2$). The four open quadrants in $\mathbb{R}^2$ are naturally in bijection to the elements $s \in SP$. We define

$$\mathcal{Q}_s = \{(x, y) \in \mathbb{R}^2 : \text{sgn } x = \pi_1(s) \text{ and } \text{sgn } y = \pi_2(s)\}.$$

We use $d(\mathcal{Q}_s)$ to denote the closure $\mathcal{Q}_s$.

Definition 6.16 (Quadrants in $\mathbb{R}^\mathcal{V}$). The four quadrants in $\mathbb{R}^\mathcal{V}$ are

$$\hat{\mathcal{Q}}_s = \left\{ \mathbf{f} \in \mathbb{R}^\mathcal{V} : \begin{array}{l} \mathbf{f}(a) = 0 \text{ or } \text{sgn } \mathbf{f}(a) = \pi_1(s) \text{ for all } a \in \mathcal{A}. \\ \text{And, } \mathbf{f}(b) = 0 \text{ or } \text{sgn } \mathbf{f}(b) = \pi_2(s) \text{ for all } b \in \mathcal{B} \end{array} \right\}.$$

Recall that the orbit under ρ_λ^G of a λ -renormalizable direction does not include the horizontal or vertical directions.

Definition 6.17 (Sign sequence of $\boldsymbol{\theta}$). Suppose $\boldsymbol{\theta}$ has λ -shrinking sequence $\langle g_i \rangle$. We define the corresponding (λ) -sign sequence of $\boldsymbol{\theta}$ to be the sequence $\langle s_0, s_1, s_2, \dots \rangle$ with $s_i \in SP$ so that $\rho_\lambda^{g_i}(\boldsymbol{\theta}) \in \mathcal{Q}_{s_i}$ for all i .

The λ -sign and λ -shrinking sequences turn out to uniquely determine $\boldsymbol{\theta}$.

Proposition 6.18 (Quadrant sequences). Let $\boldsymbol{\theta} \in \mathcal{R}_\lambda$ with λ -shrinking sequence $\langle g_i \rangle$ and sign sequence $\langle s_i \rangle$. The only $\mathbf{v} \in \mathbb{S}^1$ for which $\rho_\lambda^{g_i}(\mathbf{v}) \in \overline{\mathcal{Q}_{s_i}}$ for all i is $\mathbf{v} = \boldsymbol{\theta}$.

This proposition is proved in section 10.4.

Recall from definition 3.7 that an $\boldsymbol{\theta} \in \mathbb{R}_\lambda$ can be written in the form $\boldsymbol{\theta} = \boldsymbol{\theta}(\langle g_n \rangle, \lambda)$ up to a choice of sign. If we fix another $\lambda' \geq 2$. The pair of antipodal vectors $\pm \boldsymbol{\theta}(\langle g_n \rangle, \lambda')$ lie in the same quadrants as $\pm \boldsymbol{\theta}$. (See Proposition 3.8.) We choose $\boldsymbol{\theta}(\langle g_n \rangle, \lambda')$ so that it lies in the same quadrant as $\boldsymbol{\theta}$. We will need the following in the proof of the Topological conjugacy theorem. A stronger statement that implies this Proposition is provided by Proposition 10.6.

Proposition 6.19. The λ -sign sequence of $\boldsymbol{\theta}$ is the same as the λ' -sign sequence of $\boldsymbol{\theta}(\langle g_n \rangle, \lambda')$.

The sign sequence gives us a more natural definition of being a $(\boldsymbol{\theta}, n)$ -survivor.

Proposition 6.20 (Equivalent notion of survivors). $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a $(\boldsymbol{\theta}, n)$ -survivor if and only if $\Upsilon^{g_n}(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_n}$.

Proof. By proposition 6.12, $\Phi_*^{g_n} \circ \Xi(\mathbf{f}) = \Xi \circ \Upsilon^{g_n}(\mathbf{f})$. Showing that $\mathbf{f}$ is a $(\boldsymbol{\theta}, n)$ -survivor is equivalent to showing that $\Xi \circ \Upsilon^{g_n}(\mathbf{f})$ is a $(\rho_\lambda^{g_n}(\boldsymbol{\theta}), 0)$ -survivor. Therefore, it is enough to consider the case when $n = 0$.

Write $\boldsymbol{\theta} = (x, y)$ and set $s_x = \text{sgn } x$ and $s_y = \text{sgn } y$. As $\boldsymbol{\theta}$ is λ -shrinkable, $s_x, s_y \in \{-1, 1\}$. If $\sigma_v \in E$ is a vertical saddle connection, then $\text{sgn}(\text{hol}(\sigma_v) \wedge \boldsymbol{\theta}) = -s_x$. If $\sigma_v \in E$ is horizontal, then $\text{sgn}(\text{hol}(\sigma_h) \wedge \boldsymbol{\theta}) = s_y$. Assuming σ_v crosses the horizontal cylinder $a \in \mathcal{A}$ and σ_h crosses the vertical cylinder $b \in \mathcal{B}$, we have

$$\Xi(\mathbf{f})(\sigma_v) = -\mathbf{f}(a) \quad \text{and} \quad \Xi(\mathbf{f})(\sigma_h) = \mathbf{f}(b).$$

Therefore, $\Xi(\mathbf{f})$ is a $(\boldsymbol{\theta}, 0)$ -survivor if and only if $\text{sgn } \mathbf{f}(a) \in \{0, s_x\}$ and $\text{sgn } \mathbf{f}(b) \in \{0, s_y\}$ for all $a \in \mathcal{A}$ and $b \in \mathcal{B}$. The conclusion of the proposition follows. $\square$

6.4. The topological conjugacy. In this section we prove Theorem 4.1. Throughout this section $\mathbf{w}_1, \mathbf{w}_2 \in \mathbb{R}^\mathcal{V}$ will be positive functions satisfying $\mathbf{A}(\mathbf{w}_i) = \lambda_i \mathbf{w}_i$. The sequence $\langle g_n \rangle$ will be a renormalizing sequence, and $\boldsymbol{\theta}_i = \pm \boldsymbol{\theta}(\langle g_n \rangle, \lambda_i) \in \mathbb{S}^1$ will be chosen from the pairs of antipodal λ_i -renormalizable directions with λ_i -shrinking sequence $\langle g_n \rangle$ so that $\boldsymbol{\theta}_1$ and $\boldsymbol{\theta}_2$ lie in the same quadrant. We write $\boldsymbol{\theta}_i = (x_i, y_i)$.

For any positive $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ with $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$, there is a naturally related parametrized plane in $\mathbb{R}^\mathcal{V}$.

$$(6.6) \quad P_{\mathbf{f}} : \mathbb{R}^2 \rightarrow \mathbb{R}^\mathcal{V}; \quad P_{\mathbf{f}}(x, y)(\mathbf{v}) = \begin{cases} x\mathbf{f}(\mathbf{v}) & \text{if } \mathbf{v} \in \mathcal{A} \\ y\mathbf{f}(\mathbf{v}) & \text{if } \mathbf{v} \in \mathcal{B}. \end{cases}$$

This plane is invariant under the action of $\mathbf{H}$ and $\mathbf{V}$:

Proposition 6.21 (Invariant planes). *Let $\mathbf{v} \in \mathbb{R}^2$, then for all $g \in G$ we have $\Upsilon^g(P_{\mathbf{f}}(\mathbf{v})) = P_{\mathbf{f}}(\rho_{\lambda_i}^g(\mathbf{v}))$.*

The proof is a simple calculation.

We will be considering the parametrized planes $P_{\mathbf{w}_1}$ and $P_{\mathbf{w}_2}$. For $i = 1, 2$, let μ_i be the Lebesgue $\mathcal{F}_{\theta_i}$ -transverse measures on $S(\mathcal{G}, \mathbf{w}_i)$. These measures are closely connected to the planes constructed above.

Proposition 6.22. *For $i = 1, 2$, we have $\Psi_{\theta_i}(\mu_i) = \Xi(P_{\mathbf{w}_i}(\theta_i))$.*

Proof. Let $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$. Let $\llbracket \sigma_v \rrbracket$ be a vertical saddle connection crossing the horizontal cylinder $cyl_{\mathbf{a}}$ and oriented upward. Let $\llbracket \sigma_h \rrbracket$ be a rightward oriented horizontal saddle connection crossing the vertical cylinder $cyl_{\mathbf{b}}$. By definition of $\Psi_{\theta_i}(\mu_i)$ and the Lebesgue transverse measure, we have the following.

$$\Psi_{\theta_i}(\mu_i)(\llbracket \sigma_v \rrbracket) = (0, \mathbf{w}_i(\mathbf{a})) \wedge \theta_i = -x_i \mathbf{w}_i(\mathbf{a}).$$

$$\Psi_{\theta_i}(\mu_i)(\llbracket \sigma_h \rrbracket) = (\mathbf{w}_i(\mathbf{b}), 0) \wedge \theta_i = y_i \mathbf{w}_i(\mathbf{b}).$$

Similarly, by definition of Ξ we have the following.

$$\Xi(P_{\mathbf{w}_i}(\theta_i))(\llbracket \sigma_v \rrbracket) = i(\llbracket \sigma_v \rrbracket, \llbracket cyl_{\mathbf{a}} \rrbracket) P_{\mathbf{w}_i}(\theta_i)(\mathbf{a}) = -x_i \mathbf{w}_i(\mathbf{a}).$$

$$\Xi(P_{\mathbf{w}_i}(\theta_i))(\llbracket \sigma_h \rrbracket) = i(\llbracket \sigma_h \rrbracket, \llbracket cyl_{\mathbf{b}} \rrbracket) P_{\mathbf{w}_i}(\theta_i)(\mathbf{b}) = y_i \mathbf{w}_i(\mathbf{b}).$$

□

Proposition 6.23. *$P_{\mathbf{w}_2}(\theta_2)$ is a θ_1 -survivor. Moreover $\mathbf{v} = \theta_2$ is the unique $\mathbf{v} \in \mathbb{S}^1$ for which $P_{\mathbf{w}_2}(\mathbf{v})$ is a θ_1 -survivor.*

Proof. We show that $P_{\mathbf{w}_2}(\theta_2)$ is a (θ_1, n) -survivor for all n . Let s_n be the n -th entry in the sign sequences of θ_1 and θ_2 . (These are the same by Proposition 6.19.) By Proposition 6.21, we have that $\Upsilon^{g_n}(P_{\mathbf{w}_2}(\theta_2)) = P_{\mathbf{w}_2}(\rho_{\lambda_2}^{g_n}(\theta_2))$. We know $\rho_{\lambda_2}^{g_n}(\theta_2) \in \mathcal{Q}_{s_n}$. So by definition of $P_{\mathbf{w}_2}$, we have $P_{\mathbf{w}_2}(\rho_{\lambda_2}^{g_n}(\theta_2)) \in \hat{\mathcal{Q}}_{s_n}$ and we may apply Proposition 6.20.

Uniqueness of the choice θ_1 -survivor follows by combining the above argument with Proposition 6.18. □

Let $\mathcal{M}_i$ denote the locally finite $\mathcal{F}_{\theta_i}$ -transverse measures on $S(\mathcal{G}, \mathbf{w}_i)$. Then the following follows from Theorem 6.9 and Definition 6.13.

Corollary 6.24. *There is an unique measure $\mu'_2 \in \mathcal{M}_1$ with $\Psi_{\theta_1}(\mu'_2) = \Xi(P_{\mathbf{w}_2}(\theta_2))$.*

This measure μ'_2 is our candidate pull-back measure. It remains to build our homomorphism. We begin by building a continuous map from $\phi_1 : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$. For $e \in \mathcal{E}$, let R_e^i be the rectangles of $S(\mathcal{G}, \mathbf{w}_i)$ associated to the edge of the graph $e = \overline{ab}$. We define the map on these rectangles. Recall we may view each R_e^i as the rectangular subset of the plane, $[0, \mathbf{w}_i(\mathbf{b})] \times [0, \mathbf{w}_i(\mathbf{a})]$. (See definition 2.4.) We define the restriction map $\phi_1|_{\partial R_e^1} : \partial R_e^1 \rightarrow \partial R_e^2$ along the edges by integrating the measure μ'_2 and rescaling appropriately. For instance along the bottom edge $B \subset \mathbb{R}_e^1$, which is identified with the interval $[0, \mathbf{w}(\mathbf{b})] \times \{0\}$, we use the formula

$$\phi_1|_B(t, 0) = \left(\frac{1}{|y_2|} \mu'_2([0, t] \times \{0\}), 0 \right) \in R_e^2.$$

(Recall y_2 is the y -coordinate of θ_2 .) We must check that this map sends the bottom edge of R_e^1 to the bottom edge of R_e^2 . This follows from the fact that $\Psi_{\theta_1}(\mu'_2) = \Xi(P_{\mathbf{w}_2}(\theta_2))$, because the bottom edge of R_e^1 is a saddle connection, $\sigma_h = [0, \mathbf{w}_1(b)] \times \{0\}$. We evaluate the x -coordinate of the lower right endpoint as

$$(6.7) \quad \begin{aligned} \phi_1|_B(\mathbf{w}_i(b), 0) &= \frac{1}{|y_2|} \mu'_2(\sigma_h) = \frac{1}{|y_2|} |\Xi(P_{\mathbf{w}_2}(\theta_2))(\llbracket \sigma_h \rrbracket)| \\ &= \frac{1}{|y_2|} |P_{\mathbf{w}_2}(\theta_2)(b)| = \frac{1}{|y_2|} |y_2 \mathbf{w}_2(b)| = \mathbf{w}_2(b). \end{aligned}$$

This shows that $\phi_1|_L$ maps to the lower edge, but it may not be surjective or continuous if μ' contains atoms. For the left, top and right edges (L , T and R respectively), we use the following formulas.

$$(6.8) \quad \begin{aligned} \phi_1|_L(0, t) &= \left(0, \frac{1}{|x_2|} \mu'_2(\{0\} \times [0, t])\right), \quad \phi_1|_T(t, \mathbf{w}_1(a)) = \left(\frac{1}{|y_2|} \mu'_2([0, t] \times \{0\}), \mathbf{w}_2(a)\right), \\ \text{and } \phi_1|_R(\mathbf{w}_1(b), t) &= \left(\mathbf{w}_2(b), \frac{1}{|x_2|} \mu'_2(\{w_1(b)\} \times [0, t])\right). \end{aligned}$$

Together these define the map $\phi_1|_{\partial R_e^1} : \partial R_e^1 \rightarrow \partial R_e^2$. A similar check for each edge shows that $\phi_1|_{\partial R_e^1}$ sends the respective vertices of R_e^1 to the respective vertices of R_e^2 .

Proposition 6.25. *The map $\phi_1|_{\partial R_e^1}$ sends points on the same connected component of a leaf of $\mathcal{F}_{\theta_1}$ inside R_e^1 to points on the same connected component of a leaf of $\mathcal{F}_{\theta_2}$ inside R_e^2 .*

Proof. We can assume without loss of generality that θ_1 and θ_2 are in the first quadrant. We must check that if the points $P, Q \in \partial R_e^1$ satisfy $\overrightarrow{PQ} \parallel \theta_1$, then $\overrightarrow{P'Q'} \parallel \theta_2$ where $P' = \phi_1|_{\partial R_e^1}(P)$ and $Q' = \phi_1|_{\partial R_e^1}(Q)$. We will check this in case P lies on the left edge L and Q lies on the top edge T . We leave the remaining cases to the reader. In this case via the identification of R_e^1 with the rectangle $[0, \mathbf{w}_i(b)] \times [0, \mathbf{w}_i(a)]$, there is a $r > 0$ so that $P = (0, \mathbf{w}_1(a) - ry_1)$ and $Q = (rx_1, \mathbf{w}_1(a))$. Let $O = (0, \mathbf{w}_1(a))$. We note that the transversals $\overline{OP}$ and $\overline{OQ}$ cross the same collection of leaves. Let $s = \mu'_2(\overline{OP})$. Then by this observation $s = \mu'_2(\overline{OQ})$. We compute

$$\begin{aligned} P' &= \left(0, \frac{1}{x_2} \mu'_2(\{0\} \times [0, \mathbf{w}_1(a) - ry_1])\right) = \left(0, \frac{1}{x_2} (\mu'_2(\{0\} \times [0, \mathbf{w}_1(a)]) - \mu'_2(\overline{OP}))\right) \\ &= \left(0, \frac{1}{x_2} (x_2 \mathbf{w}_2(a) - s)\right) = \left(0, \mathbf{w}_2(a) - \frac{s}{x_2}\right), \text{ and} \\ Q' &= \left(\frac{1}{y_2} \mu'_2(\overline{OQ}), \mathbf{w}_2(a)\right) = \left(\frac{s}{y_2}, \mathbf{w}_2(a)\right). \end{aligned}$$

Indeed, $\overrightarrow{P'Q'} = \frac{s}{x_2 y_2} \theta_2$, with $\theta_2 = (x_2, y_2)$. □

By the proposition, we can extend definition of $\phi_1|_{\partial R_e^1}$ to a map $\phi_1|_{R_e^1} : R_e^1 \rightarrow R_e^2$. If $\{P, Q\}$ is the boundary of a connected component of a leaf of $\mathcal{F}_{\theta_1}$ inside R_e^1 , then by the proposition we know $\overrightarrow{P'Q'}$ is a connected component of a leaf of $\mathcal{F}_{\theta_2}$ inside R_e^2 , where $P' = \phi_1|_{\partial R_e^1}(P)$ and $Q' = \phi_1|_{\partial R_e^1}(Q)$. Then we can define ϕ_1 onto $\overrightarrow{PQ}$ by defining $\overrightarrow{PQ} \rightarrow \overrightarrow{P'Q'}$ affinely (so it scales distance linearly). Thus we have defined $\phi_1|_{R_e^1} : R_e^1 \rightarrow R_e^2$.

We similarly define $\phi_1|_{R_e^1} : R_e^1 \rightarrow R_e^2$ for all $e \in \mathcal{E}$. These maps agree on the boundaries of rectangles, so we have defined the map $\phi_1 : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$. We must show that ϕ_1 is continuous. Discontinuities of ϕ_1 can only arise from atoms of μ'_2 . For each atom A of μ'_2 there is a strip in $S(\mathcal{G}, \mathbf{w}_2) \setminus \phi_1(S(\mathcal{G}, \mathbf{w}_1))$ whose width is the measure of the atom.

Note that each vertex of each rectangles R_e^2 is in $\phi_1(S(\mathcal{G}, \mathbf{w}_1))$. Therefore for each atom A we have an isometrically embedded strip $\epsilon : (0, \mu'_2(\{A\})) \times \mathbb{R} \rightarrow S(\mathcal{G}, \mathbf{w}_2)$ which sends the vertical foliation of the strip to the foliation $\mathcal{F}_{\theta_2}$. But, the vertical flow in the strip is not conservative, and so the existence of this strip contradicts the fact that the straight-line flow in direction θ_2 on $S(\mathcal{G}, \mathbf{w}_2)$ is conservative. See Theorem 6.3. Thus $\phi_1 : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$ is indeed continuous and surjective.

It remains to show that ϕ_1 is invertible. We can construct a map $\phi_2 : S(\mathcal{G}, \mathbf{w}_2) \rightarrow S(\mathcal{G}, \mathbf{w}_1)$ by switching the rolls of 1 and 2. The composition $\phi_2 \circ \phi_1 : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_1)$ preserves the Lebesgue measures of transversals to $\mathcal{F}_{\theta_1}$, and it preserves all vertices of rectangles R_e^1 . Therefore, the composition must act trivially on the boundaries of rectangles R_e^1 . Moreover, the composition acts affinely on connected components of leaves intersected with R_e^1 , and so must act trivially on each such connected component of a leaf. Therefore, $\phi_2 \circ \phi_1$ is the identity map.

Finally, we consider the uniqueness statement. Now suppose there were $\mu'_2, \mu''_2 \in \mathcal{M}_1$ so that there are homeomorphisms $\phi'_1, \phi''_1 : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$ which send R_e^1 to R_e^2 and respect the names of the boundary edges of the rectangles (bottom, top, left and right) and which push the measures μ'_2, μ''_2 forward to the Lebesgue $\mathcal{F}_{\theta_2}$ -transverse measure on $S(\mathcal{G}, \mathbf{w}_2)$. Then we notice that the cohomology classes $\Psi_{\theta_1}(\mu'_2)$ and $\Psi_{\theta_1}(\mu''_2)$ must be the same. Therefore, $\mu'_2 = \mu''_2$ by Lemma 6.2 and Theorem 6.3. Finally, we see that the restrictions of ϕ'_1 and ϕ''_1 to the boundaries of rectangles are determined by μ'_2 and μ''_2 . (In fact they must be determined as in equations 6.7 and 6.8.) Thus, the restrictions of ϕ'_1 and ϕ''_1 to the boundaries of rectangles must be the same. This concludes the proof of Theorem 4.1.

We have the following corollary to the topological conjugacy theorem, which will be useful later. It is stated in the context of Theorem 4.1.

Corollary 6.26. *The pullback under ϕ of the Lebesgue $\mathcal{F}_{\theta_2}$ -transverse measure on $S(\mathcal{G}, \mathbf{w}_2)$ is $\Psi_{\theta_1}^{-1} \circ \Xi(P_{\mathbf{w}_2}(\theta_2))$.*

6.5. Surviving functions and measures. We will now continue the discussion from Section 6.3. Recall, we were able to define invariant measures in $\mathcal{M}_{\theta}$ by showing that $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ was a θ -survivor. Then we obtain an invariant measure $\Psi_{\theta}^{-1} \circ \Xi(\mathbf{f})$ by Theorem 6.9 and Definition 6.13. It turns out that under a “reasonable” assumption all measures arise in this way. Recall that $\mathcal{S}_{\theta}$ denotes the set of all θ -survivors in $\mathbb{R}^{\mathcal{V}}$.

Definition 6.27 (Subsequence decay property). We say that $S(\mathcal{G}, \mathbf{w})$ has the *subsequence decay property*, if for any $\theta \in \mathcal{R}_{\lambda}$ and any $\mathbf{f} \in \mathcal{S}_{\theta}$ there is a subsequence $\langle g_{n_i} \rangle$ of the λ -shrinking sequence of θ for which

$$\lim_{i \rightarrow \infty} \Upsilon^{g_{n_i}}(\mathbf{f})(v) = 0 \quad \text{for all } v \in \mathcal{V}.$$

Theorem 6.28. *Suppose $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property, and let $\theta \in \mathcal{R}_{\lambda}$. If $m \in H^1(S, V, \mathbb{R})$ is a θ -survivor, then $m = \Xi(\mathbf{f})$ where $\mathbf{f} \in \mathcal{S}_{\theta}$.*

We prove Theorem 6.28 in section 11. In section 14, we will show that if $\mathcal{G}$ has no nodes of valance one, then $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property. (In fact, we prove that $S(\mathcal{G}, \mathbf{w})$ has a stronger property. See Definition 6.37 and Theorem 6.39, below.)

Remark 6.29. The author believes that the subsequence decay property should hold whenever $\mathcal{G}$ is an infinite connected graph.

By combining Theorems 6.28 and 6.9, we have the following.

Corollary 6.30. *If $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property, then $\Psi_{\boldsymbol{\theta}}(\mathcal{M}_{\boldsymbol{\theta}}) = \Xi(\mathcal{S}_{\boldsymbol{\theta}})$. Moreover, both $\Psi_{\boldsymbol{\theta}}$ and Ξ are injective, so this yields a linear isomorphism between $\mathcal{M}_{\boldsymbol{\theta}}$ and $\mathcal{S}_{\boldsymbol{\theta}}$.*

The following is equivalent to the corollary above, but sounds more impressive.

Corollary 6.31 (Semi-conjugacy). *Assume $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property. Given any $\mu \in \mathcal{M}_{\boldsymbol{\theta}}$, there is a function $\mathbf{f} \in \mathbb{R}_{\geq 0}^{\mathcal{V}}$ (with $\Psi_{\boldsymbol{\theta}}(\mu) = \Xi(\mathbf{f})$) and a continuous map $h : S(\mathcal{G}, \sqrt{2} \cdot \mathbf{f}) \rightarrow S(\mathcal{G}, \mathbf{w})$ which sends the foliation by lines of slope ± 1 (with sign depending on the sign of the slope of $\boldsymbol{\theta}$) to the foliation $\mathcal{F}_{\boldsymbol{\theta}}$ of $S(\mathcal{G}, \mathbf{w})$ such that μ is the pushforward of the Lebesgue transverse measure on $S(\mathcal{G}, \sqrt{2} \cdot \mathbf{f})$ which is transverse to the foliation by lines of slope ± 1 .*

We will not describe the proof of this fact, because it the construction of the semi-conjugacy is very similar to a portion of the proof of the Topological Conjugacy Theorem given in the previous subsection. The statement of this corollary is also irrelevant to our later arguments.

6.6. The action of the adjacency operator. In this section, we introduce the adjacency operator $\mathbf{A}$ to our arguments, and observe how $\mathbf{A}$ interacts with the operators $\mathbf{H}$ and $\mathbf{V}$.

Let $\bar{\cdot} : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ denote the involution $\overline{(x, y)} = (y, x)$. The action of $\bar{\cdot}$ permutes the quadrants. We define the action of $\bar{\cdot}$ on pairs of signs so that $\overline{\mathcal{Q}_s} = \mathcal{Q}_{\bar{s}}$. The action of $\bar{\cdot}$ conjugates ρ_{λ}^h to ρ_{λ}^v and vice versa. Thus we also define the group homomorphism $\bar{\cdot} : G \rightarrow G$ by extending the action on generators $\bar{h} = v$ and $\bar{v} = h$. Thus, for all $g \in G$ and all $\mathbf{v} \in \mathbb{R}^2$ we have

$$\rho_{\lambda}^{\bar{g}}(\mathbf{v}) = \overline{\rho_{\lambda}^g(\bar{\mathbf{v}})}.$$

Recall that $\mathbf{A}\mathbf{H} = \mathbf{V}\mathbf{A}$ and $\mathbf{A}\mathbf{V} = \mathbf{H}\mathbf{A}$. Thus, for all $g \in G$ we have

$$(6.9) \quad \mathbf{A}\Upsilon^g = \Upsilon^{\bar{g}}\mathbf{A}.$$

The previous section showed that if $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property, then for a λ -renormalizable direction $\boldsymbol{\theta} \in \mathcal{R}_{\lambda}$ the composition $\Xi^{-1} \circ \Psi_{\boldsymbol{\theta}}$ is a bijective correspondence between $\mathcal{M}_{\boldsymbol{\theta}}$, the space of transverse measures for the foliation in direction $\boldsymbol{\theta}$, and the space $\mathcal{S}_{\boldsymbol{\theta}}$ of $\boldsymbol{\theta}$ -survivors in $\mathbb{R}^{\mathcal{V}}$. We have the following.

Proposition 6.32. *If $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ is a $\boldsymbol{\theta}$ -survivor then $\mathbf{A}(\mathbf{f})$ is a $\bar{\boldsymbol{\theta}}$ -survivor.*

Proof. Let $\langle g_n \rangle$ and $\langle s_n \rangle$ denote the shrinking and sign sequences of $\boldsymbol{\theta}$, respectively. Then the shrinking sequence of $\bar{\boldsymbol{\theta}}$ is $\langle \bar{g}_n \rangle$ and the sign sequence is $\langle \bar{s}_n \rangle$. By the notion of survivors given in Proposition 6.20, as $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor, we have $\Upsilon^{g_n}(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_n}$ for all n . By equation 6.9, we know

$$\Upsilon^{\bar{g}_n} \circ \mathbf{A}(\mathbf{f}) = \mathbf{A} \circ \Upsilon^{g_n}(\mathbf{f}) \in \mathbf{A}(\hat{\mathcal{Q}}_{s_n}) \subset \hat{\mathcal{Q}}_{\bar{s}_n}$$

for all n . Hence, $\mathbf{A}(\mathbf{f})$ is a $\bar{\boldsymbol{\theta}}$ -survivor by Proposition 6.20. $\square$

We will show that $\mathbf{A}$ is a bijection from the cone of $\boldsymbol{\theta}$ -survivors to the cone of $\bar{\boldsymbol{\theta}}$ -survivors. First we will discuss injectivity. Note that $\mathbf{A}$ is not injective when considered on $\mathbb{R}^{\mathcal{V}}$. We have the following description of the the group action Υ on $\ker \mathbf{A} = \{\mathbf{k} \in \mathbb{R}^{\mathcal{V}} : \mathbf{A}\mathbf{k} = \mathbf{0}\}$. For any subset $U \subset \mathcal{V}$ we define $\pi_U : \mathbb{R}^{\mathcal{V}} \rightarrow \mathbb{R}^{\mathcal{V}}$ by

$$(6.10) \quad \pi_U(\mathbf{f})(v) = \begin{cases} \mathbf{f}(v) & \text{if } v \in U \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 6.33 (Kernel of $\mathbf{A}$). *For all $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ and all $k \in \mathbb{Z}$, we have*

$$\mathbf{H}^k(\mathbf{f}) = \mathbf{f} + k\pi_B \circ \mathbf{A}(\mathbf{f}) \quad \text{and} \quad \mathbf{V}^k(\mathbf{f}) = \mathbf{f} + k\pi_A \circ \mathbf{A}(\mathbf{f}).$$

In particular, if $\mathbf{k} \in \ker \mathbf{A}$ then $\Upsilon^g(\mathbf{k}) = \mathbf{k}$ for all $g \in G$.

The proof is trivial, and follows from comparing the definitions of $\mathbf{A}$, $\mathbf{H}$ and $\mathbf{V}$. (See equations 1.4, 6.2 and 6.3, respectively.)

Proposition 6.34 (Injectivity of $\mathbf{A}$). *Suppose $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property. Then, the restriction of $\mathbf{A}$ to the cone of all $\boldsymbol{\theta}$ -survivors is injective.*

Proof. Suppose that $\mathbf{f}_1, \mathbf{f}_2$ are both $\boldsymbol{\theta}$ -survivors and that $\mathbf{A}(\mathbf{f}_1) = \mathbf{A}(\mathbf{f}_2)$. Then $\mathbf{f}_2 = \mathbf{f}_1 + \mathbf{k}$ with $\mathbf{k} \in \ker \mathbf{A}$. We will show $\mathbf{k}(\mathbf{x}) = 0$ for all $\mathbf{x} \in \mathcal{V}$. By Proposition 6.33, for all $g \in G$ we have

$$\mathbf{k} = \Upsilon^g(\mathbf{k}) = \Upsilon^g(\mathbf{f}_2) - \Upsilon^g(\mathbf{f}_1)$$

By the decay lemma, we know that there is a subsequence g_{n_i} of the λ -shrinking sequence for $\boldsymbol{\theta}$ so that $\lim_{i \rightarrow \infty} \Upsilon^{g_i}(\mathbf{f}_j) = \mathbf{0}$ coordinatewise for $j = 1, 2$. For all $\mathbf{x} \in \mathcal{V}$,

$$\mathbf{k}(\mathbf{x}) = \lim_{i \rightarrow \infty} \Upsilon^{g_{n_i}}(\mathbf{k})(\mathbf{x}) = \lim_{i \rightarrow \infty} (\Upsilon^{g_{n_i}}(\mathbf{f}_2) - \Upsilon^{g_{n_i}}(\mathbf{f}_1))(\mathbf{x}) = 0.$$

□

We will now discuss why $\mathbf{A}$ surjectively maps the $\boldsymbol{\theta}$ -survivors onto the space of $\bar{\boldsymbol{\theta}}$ -survivors. To do this we need to be able to find inverses under the adjacency operator.

We use $\mathbb{R}_c^{\mathcal{V}}$ to denote compactly supported functions $\mathcal{V} \rightarrow \mathbb{R}$. That is, a $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$ takes non-zero values only on finitely many $\mathbf{v} \in \mathcal{V}$. There is a natural bilinear pairing $\langle \cdot, \cdot \rangle : \mathbb{R}^{\mathcal{V}} \times \mathbb{R}_c^{\mathcal{V}} \rightarrow \mathbb{R}$ given by

$$(6.11) \quad \langle \mathbf{f}, \mathbf{x} \rangle = \sum_{\mathbf{v} \in \mathcal{V}} \mathbf{x}(\mathbf{v}) \mathbf{f}(\mathbf{v}).$$

This sum is well defined because $\mathbf{x}$ is only non-zero at finitely many $\mathbf{v} \in \mathcal{V}$. The operators $\mathbf{A}$, $\mathbf{H}$ and $\mathbf{V}$ restrict to actions on $\mathbb{R}_c^{\mathcal{V}}$. Furthermore,

$$(6.12) \quad \langle \mathbf{A}\mathbf{f}, \mathbf{x} \rangle = \langle \mathbf{f}, \mathbf{A}\mathbf{x} \rangle, \quad \langle \mathbf{H}\mathbf{f}, \mathbf{x} \rangle = \langle \mathbf{f}, \mathbf{V}\mathbf{x} \rangle, \quad \text{and} \quad \langle \mathbf{V}\mathbf{f}, \mathbf{x} \rangle = \langle \mathbf{f}, \mathbf{H}\mathbf{x} \rangle$$

for all $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ and all $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$. We define the group automorphism $\gamma : G \rightarrow G$ by extending the definition on generators

$$(6.13) \quad \gamma(h) = v^{-1} \quad \text{and} \quad \gamma(v) = h^{-1}.$$

Note γ^2 is trivial. The natural extension of equation 6.12 to all $g \in G$ is

$$(6.14) \quad \langle \Upsilon^g \mathbf{f}, \Upsilon^{\gamma(g)} \mathbf{x} \rangle = \langle \mathbf{f}, \mathbf{x} \rangle$$

We have the following variation of a theorem of Farkas [Far02]. (See Corollary 3.46 of [AT07], for a more modern treatment.)

Lemma 6.35 (Farkas' theorem for the adjacency operator). *Let $\mathbf{f} \in \hat{\mathcal{Q}}_{++}$. The following two statements are equivalent.*

- (1) *There is a $\mathbf{g} \in \hat{\mathcal{Q}}_{++}$ with $\mathbf{A}\mathbf{g} = \mathbf{f}$.*
- (2) *For all $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$, if $\mathbf{A}\mathbf{x} \in \hat{\mathcal{Q}}_{++}$ then $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$.*

This lemma is proved in section 13. We will use criterion (2) to check for surjectivity. We will simplify this criterion. First, we introduce a more strict version of the subsequence decay property. In order to define this property, we need the following useful fact about renormalizable directions.

Lemma 6.36 (Critical times). *Suppose θ is a λ -renormalizable direction. Then there are infinitely many $n \in \mathbb{N}$ such that if $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ is a (θ, n) -survivor, then $\mathbf{f}$ is a (θ, m) -survivor for all $m \leq n$.*

This lemma is a direct consequence of Corollaries 10.9 and 11.5. We prove this lemma, by showing that there a sequence of values of n which are guaranteed to have the property above. These values of n satisfy a simple combinatorial criterion related to the shrinking sequence, and we call them *critical times*. See Definition 10.7.

Definition 6.37 (Critical decay property). $S(\mathcal{G}, \mathbf{w})$ has the *critical decay property*, if for any $\theta \in \mathcal{R}_\lambda$ and any $\mathbf{f} \in \mathcal{S}_\theta$ the sequence of critical times, $\langle g_{n_i} \rangle$, of the λ -shrinking sequence of θ satisfies

$$\lim_{i \rightarrow \infty} \Upsilon^{g_{n_i}}(\mathbf{f})(\mathbf{v}) = 0 \quad \text{for all } \mathbf{v} \in \mathcal{V}.$$

The critical decay property implies the subsequence decay property of definition 6.27.

Definition 6.38 (Adjacency sign property). The graph $\mathcal{G}$ has the *adjacency sign property*, if for any $\theta \in \mathcal{Q}_{++} \cap \mathcal{R}_\lambda$, any $\mathbf{f} \in \mathcal{S}_\theta$, any $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$ such that $\mathbf{A}\mathbf{x} \in \hat{\mathcal{Q}}_{++}$ and any critical time t for the λ -shrinking sequence $\langle g_n \rangle$ we have

$$\langle \Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{x}) - \mathbf{x} \rangle \geq 0.$$

In section 14 we prove the following.

Theorem 6.39 (Graphs without vertices of valance one). *If $\mathcal{G}$ is an infinite connected bipartite graph with no vertices of valance one and $\mathbf{w}$ is a positive eigenfunction for the adjacency operator, then $S(\mathcal{G}, \mathbf{w})$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property.*

Proposition 6.40. *Assume $S(\mathcal{G}, \mathbf{w})$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Let $\theta \in \mathcal{R}_\lambda$ and $\mathbf{g} \in \mathcal{S}_\theta$. Then there is a $(\theta, 0)$ -survivor, $\mathbf{f}$, with $\mathbf{A}(\mathbf{f}) = \mathbf{g}$.*

Proof. Without loss of generality, we may assume $\mathbf{g} \in \hat{\mathcal{Q}}_{++}$. The statement that $\mathbf{f}$ is a $(\theta, 0)$ -survivor is equivalent to saying $\mathbf{f} \in \hat{\mathcal{Q}}_{++}$. We show the existence of such an $\mathbf{f}$ using Farkas' theorem for the adjacency operator. Let $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$ with $\mathbf{A}\mathbf{x} \in \mathcal{Q}_{++}$. By equation 6.14, for any $g \in G$,

$$\langle \mathbf{f}, \mathbf{x} \rangle = \langle \Upsilon^g(\mathbf{f}), \Upsilon^{\gamma(g)}(\mathbf{x}) - \mathbf{x} \rangle + \langle \Upsilon^g(\mathbf{f}), \mathbf{x} \rangle.$$

Let $\langle g_{n_i} \rangle$ be the critical subsequence of the λ -shrinking sequence of θ . By the critical decay property, we see $\langle \Upsilon^{g_{n_i}}(\mathbf{f}), \mathbf{x} \rangle \rightarrow 0$ as $i \rightarrow \infty$. By the adjacency sign property, we have $\langle \Upsilon^{g_{n_i}}(\mathbf{f}), \Upsilon^{\gamma(g_{n_i})}(\mathbf{x}) - \mathbf{x} \rangle$ is always non-negative. Thus,

$$\langle \mathbf{f}, \mathbf{x} \rangle = \lim_{i \rightarrow \infty} \langle \Upsilon^{g_{n_i}}(\mathbf{f}), \Upsilon^{\gamma(g_{n_i})}(\mathbf{x}) - \mathbf{x} \rangle + \langle \Upsilon^{g_{n_i}}(\mathbf{f}), \mathbf{x} \rangle \geq 0$$

as needed to apply Farkas' Theorem (Lemma 6.35). □

Given proposition 6.40, the proof that there is a θ -survivor, $\mathbf{g}$, with $\mathbf{A}\mathbf{g} = \mathbf{f}$ is not difficult. Roughly, the proof carefully combines this Proposition with the Critical Times Lemma. Let n be a critical time. Then, we can find a $(\rho_\lambda^{g_n}(\theta), 0)$ -survivor, $\mathbf{g}_n$, such that $\mathbf{A}(\mathbf{g}_n) = \Upsilon^{g_n}(\mathbf{f})$. Let $\mathbf{g} = \Upsilon^{\bar{g}_n^{-1}}(\mathbf{g}_n)$. By Lemma 6.36, $\mathbf{g}$ is a $(\bar{\theta}, m)$ -survivor for all $m = 0, \dots, n$. Thus, the main technical difficulty is showing that the sets of all such candidates of the form $\Upsilon^{\bar{g}_n^{-1}}(\mathbf{g}_n)$ for each critical time n have a non-trivial intersection. This argument is fully explained in section 13 in the proof of the following theorem.

Lemma 6.41 (Surjectivity). *Assume $S(\mathcal{G}, \mathbf{w})$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Let $\theta \in \mathcal{R}_\lambda$ and $\mathbf{g} \in \mathcal{S}_\theta$. Then there is a $\mathbf{f} \in \mathcal{S}_\theta$ with $\mathbf{A}(\mathbf{f}) = \mathbf{g}$.*

Combining Lemma 6.41 with Proposition 6.34, we observe the following.

Theorem 6.42. *Assume $S(\mathcal{G}, \mathbf{w})$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. For θ a λ -renormalizable direction, the adjacency operator restricts to bijection between the set of θ -survivors and the set of $\bar{\theta}$ -survivors.*

We have the following trivial proposition.

Proposition 6.43. *Suppose $\mathbf{f} \in \mathcal{S}_\theta$ then $\mathbf{A}^2(\mathbf{f}) - \mathbf{f} \in \mathcal{S}_\theta$.*

Proof. We consider the operator $\mathbf{A}^2 - \mathbf{I}$. First observe that for all $s \in SP$, we have $(\mathbf{A}^2 - \mathbf{I})(\hat{\mathcal{Q}}_s) \subset \hat{\mathcal{Q}}_s$. To see this, we may take $s = ++$ without loss of generality. Then, if $\mathbf{v} \in \mathcal{V}$ we can find an $\mathbf{w} \sim \mathbf{v}$. Then by definition of $\mathbf{A}$ we have

$$\mathbf{A}^2(\mathbf{f})(\mathbf{v}) \geq \mathbf{A}(\mathbf{f})(\mathbf{w}) \geq \mathbf{f}(\mathbf{v}),$$

and therefore $\mathbf{A}^2(\mathbf{f})(\mathbf{v}) - \mathbf{f}(\mathbf{v}) \geq 0$ as desired.

Second, we may observe that $\mathbf{A}^2 - \mathbf{I}$ commutes with Υ^g for all $g \in G$. This follows from linearity and equation 6.9. That is,

$$\Upsilon^{g_n} \circ (\mathbf{A}^2 - \mathbf{I})(\mathbf{f}) = (\mathbf{A}^2 - \mathbf{I}) \circ \Upsilon^{g_n}(\mathbf{f}).$$

Since we assumed $\mathbf{f}$ was a (θ, n) survivor, we know $\Upsilon^{g_n}(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_n}$. Therefore, by our first observation applied to the right hand side above, we see $\Upsilon^{g_n} \circ (\mathbf{A}^2 - \mathbf{I})(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_n}$ as well. Thus, $(\mathbf{A}^2 - \mathbf{I})(\mathbf{f})$ is a (θ, n) -survivor for all n . $\square$

Our main theorem is really a corollary of Theorem 6.42. The Ergodic measures theorem (Theorem 4.3) follows by combining the theorem below with Theorem 6.39 and some Martin boundary theory as described in the next subsection.

Theorem 6.44. *Assume $S(\mathcal{G}, \mathbf{w})$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. If $\mathbf{f} \in \mathcal{S}_\theta$ is extremal, then $\mathbf{A}^2(\mathbf{f}) = \lambda_2^2(\mathbf{f})$ for some positive $\lambda_2 \in \mathbb{R}$.*

Proof. If $\mathbf{f}$ is extremal, then $\mathbf{A}^2(\mathbf{f})$ must be extremal, since $\mathbf{A}^2 : \mathcal{S}_\theta \rightarrow \mathcal{S}_\theta$ is a linear bijection. But, $\mathbf{A}^2(\mathbf{f}) = \mathbf{f} + (\mathbf{A}^2(\mathbf{f}) - \mathbf{f})$ and both $\mathbf{f}, \mathbf{A}^2(\mathbf{f}) - \mathbf{f} \in \mathcal{S}_\theta$. These two functions are linearly independent unless $\mathbf{A}^2(\mathbf{f}) = \lambda_2^2(\mathbf{f})$ for some positive $\lambda_2 \in \mathbb{R}$. $\square$

6.7. Martin boundary theory. In this subsection, we use Martin boundary theory to finish the proof of the ergodic measures theorem. We will not define the Martin boundary here. Rather we will only introduce the facts we need about it. We will define the Martin boundary in section 15. We refer the reader to [MW89] and [Woe00] for more information.

Choose any vertex $o \in \mathcal{V}$ called the root. Given λ such that positive eigenfunctions of $\mathbf{A}$ exist with eigenvalue λ , the Martin compactification $\mathcal{V}_\lambda$ is a compactification of the vertex

set $\mathcal{V}$. The Martin boundary is the set $\mathcal{M}_\lambda = \mathcal{V}_\lambda \setminus \mathcal{V}$. Points $\zeta \in \mathcal{M}_\lambda$ correspond to positive functions $\mathbf{k}_\zeta \in \mathbb{R}^\mathcal{V}$ so that $\mathbf{A}(\mathbf{k}_\zeta) = \lambda \mathbf{k}_\zeta$ and $\mathbf{k}_\zeta(o) = 1$. The function $\zeta \mapsto \mathbf{k}_\zeta$ is continuous when $\mathbb{R}^\mathcal{V}$ is given the topology of pointwise convergence. The subset

$$\mathcal{M}_\lambda^{\min} = \{\zeta \in \mathcal{M}_\lambda : \mathbf{k}_\zeta \text{ is an extremal eigenfunction}\}.$$

is called the minimal Martin boundary, and is a Borel subset of $\mathcal{M}_\lambda$.

Theorem 6.45 (Poisson-Martin representation theorem). *For each positive $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ with $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$, there is a unique Borel measure $\nu_{\mathbf{f}}$ on $\mathcal{M}_\lambda$ with $\nu_{\mathbf{f}}(\mathcal{M}_\lambda \setminus \mathcal{M}_\lambda^{\min}) = 0$ and*

$$\mathbf{f}(v) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(v) d\nu_{\mathbf{f}}(\zeta),$$

for all $v \in \mathcal{V}$.

Now we will study those $\mathbf{f} \in \hat{\mathcal{Q}}_s$ with s a sign pair and for which $\mathbf{A}^2(\mathbf{f}) = \lambda^2(\mathbf{f})$. We let $\mathcal{G}^2$ denote the graph with vertex set $\mathcal{V}$ and edge set $\mathcal{E}^2$, where each path of length two in $\mathcal{E}$ corresponds to an edge of $\mathcal{E}^2$. We allow loops in the graph $\mathcal{G}^2$, which arise from moving forward along an edge in $\mathcal{E}$ and then back to the starting point along the same edge or another edge with the same endpoints. Since $\mathcal{G}$ is bipartite, the graph $\mathcal{G}^2$ consists of two connected components with vertex sets $\mathcal{A}$ and $\mathcal{B}$. We denote these two components by $\mathcal{G}_{\mathcal{A}}$ and $\mathcal{G}_{\mathcal{B}}$. We will let $\mathbf{A}_{\mathcal{A}}$ and $\mathbf{A}_{\mathcal{B}}$ be the adjacency operators on these two graphs and they satisfy the equations

$$\mathbf{A}^2(\mathbf{f})(a) = \mathbf{A}_{\mathcal{A}}(\mathbf{f}|_{\mathcal{A}})(a) \quad \text{and} \quad \mathbf{A}^2(\mathbf{f})(b) = \mathbf{A}_{\mathcal{B}}(\mathbf{f}|_{\mathcal{B}})(b)$$

for all $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ and all $a \in \mathcal{A}$ and $b \in \mathcal{B}$.

Let E be the collection of positive $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ such that $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$. For $s \in SP$, define $\hat{E}_s = \{\mathbf{f} \in \hat{\mathcal{Q}}_s : \mathbf{A}^2(\mathbf{f}) = \lambda^2(\mathbf{f})\}$. Note that $E \subset \hat{E}_{++}$. For $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ we define the functions $\mathbf{f}_{\mathcal{A}}, \mathbf{f}_{\mathcal{B}} \in \mathbb{R}^\mathcal{V}$ according to the rule

$$(6.15) \quad \mathbf{f}_{\mathcal{A}}(v) = \begin{cases} \mathbf{f}(v) & \text{if } v \in \mathcal{A} \\ \frac{1}{\lambda} \mathbf{A}(\mathbf{f})(v) & \text{if } v \in \mathcal{B} \end{cases} \quad \text{and} \quad \mathbf{f}_{\mathcal{B}}(v) = \begin{cases} \frac{1}{\lambda} \mathbf{A}(\mathbf{f})(v) & \text{if } v \in \mathcal{A} \\ \mathbf{f}(v) & \text{if } v \in \mathcal{B}. \end{cases}$$

Note that if $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a positive function satisfying $\mathbf{A}^2(\mathbf{f}) = \lambda^2 \mathbf{f}$ then we have $\mathbf{A}(\mathbf{f}_{\mathcal{A}}) = \lambda \mathbf{f}_{\mathcal{A}}$ and $\mathbf{A}(\mathbf{f}_{\mathcal{B}}) = \lambda \mathbf{f}_{\mathcal{B}}$. In fact,

Proposition 6.46. *For all $s = (s_1, s_2) \in SP$, the function $\mathbf{f} \mapsto (\mathbf{f}_{\mathcal{A}}, \mathbf{f}_{\mathcal{B}})$ restricts to a linear bijection $\mathbf{D} : \hat{E}_s \rightarrow s_1 E \times s_2 E$. Here, $s_i E$ denotes positive eigenfunctions with eigenvalue λ when $s_i = 1$ and the negative eigenfunctions when $s_i = -1$. The inverse map is given by $\mathbf{D}^{-1}(\mathbf{f}_{\mathcal{A}}, \mathbf{f}_{\mathcal{B}}) = \pi_{\mathcal{A}}(\mathbf{f}_{\mathcal{A}}) + \pi_{\mathcal{B}}(\mathbf{f}_{\mathcal{B}})$, with π_* defined as in equation 6.10.*

This means that we can use the Poisson-Martin representation theorem to express those $\mathbf{f} \in \hat{E}_s$ for $s \in SP$. For this it is natural to consider signed measures on the Martin boundary. For ν a Borel measure on $\mathcal{M}_\lambda$, we let ν^+ and ν^- be the mutually singular unsigned Borel measures satisfying $\nu = \nu^+ - \nu^-$ obtained via the Hahn decomposition theorem. We let M be the space of signed Borel measures ν on $\mathcal{M}_\lambda$ such that

- (1) $\nu(A) = 0$ for all measurable $A \subset \mathcal{M}_\lambda \setminus \mathcal{M}_\lambda^{\min}$, and
- (2) $\int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(v) d\nu(\zeta)$ is defined and finite for all $v \in \mathcal{V}$.

Note the space M is a real vector space. We also define the subsets

$$M^+ = \{\nu \in M : \nu^-(\mathcal{M}_\lambda) = 0\} \quad \text{and} \quad M^- = \{\nu \in M : \nu^+(\mathcal{M}_\lambda) = 0\}.$$

We have the following corollary to the Poisson-Martin representation theorem.

Corollary 6.47. *Let $s = (s_1, s_2) \in SP$ and $\mathbf{f} \in \widehat{E}_s$. Then there is a unique $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}}) \in M^2 = M \times M$ for which*

$$\mathbf{f}(\mathbf{a}) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{a}) d\nu_{\mathcal{A}}(\zeta) \quad \text{and} \quad \mathbf{f}(\mathbf{b}) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{b}) d\nu_{\mathcal{B}}(\zeta)$$

for all $\mathbf{a} \in \mathcal{A}$ and all $\mathbf{b} \in \mathcal{B}$. Moreover, $\nu_{\mathcal{A}} \in M^{s_1}$ and $\nu_{\mathcal{B}} \in M^{s_2}$.

Proof. We first prove existence. Given $\mathbf{f}$, we may consider $\mathbf{f}_{\mathcal{A}}, \mathbf{f}_{\mathcal{B}}$ defined as in equation 6.15. We have that $s_1 \mathbf{f}_{\mathcal{A}}, s_2 \mathbf{f}_{\mathcal{B}} \in E$. Therefore using the notation from the Poisson-Martin representation theorem, we may set $\nu_{\mathcal{A}} = s_1 \nu_{s_1 \mathbf{f}_{\mathcal{A}}}$ and $\nu_{\mathcal{B}} = s_2 \nu_{s_2 \mathbf{f}_{\mathcal{B}}}$. The pair $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}})$ satisfy the equations of the corollary and we have $\nu_{\mathcal{A}} \in M^{s_1}$ and $\nu_{\mathcal{B}} \in M^{s_2}$.

Now assume that $(\mu_{\mathcal{A}}, \mu_{\mathcal{B}}) \in M^2$ is a second pair of measures satisfying the statement of the corollary. We will prove that $\mu_{\mathcal{A}} = \nu_{\mathcal{A}}$. Define the unsigned measures $\sigma_1 = \nu_{\mathcal{A}}^+ + \mu_{\mathcal{A}}^-$ and $\sigma_2 = \nu_{\mathcal{A}}^- + \mu_{\mathcal{A}}^+$. Define $\mathbf{g}_1, \mathbf{g}_2 \in \mathbb{R}^{\mathcal{V}}$ by

$$\mathbf{g}_i(\mathbf{v}) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{v}) d\sigma_i(\zeta) \quad \text{for all } \mathbf{v} \in \mathcal{V}.$$

Note that for all $\mathbf{a} \in \mathcal{A}$ we have $\mathbf{g}_1(\mathbf{a}) = \mathbf{g}_2(\mathbf{a})$ since

$$\mathbf{f}(\mathbf{a}) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{a}) d\nu_{\mathcal{A}}(\zeta) = \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{a}) d\mu_{\mathcal{A}}(\zeta).$$

So that $\mathbf{g}_1(\mathbf{a}) = \mathbf{g}_2(\mathbf{a}) = \mathbf{f}(\mathbf{a}) + \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{a}) d\nu_{\mathcal{A}}^-(\zeta) + \int_{\mathcal{M}_\lambda} \mathbf{k}_\zeta(\mathbf{a}) d\mu_{\mathcal{A}}^-(\zeta)$. Now observe that both $\mathbf{g}_1$ and $\mathbf{g}_2$ satisfy $\mathbf{A}(\mathbf{g}_i) = \lambda \mathbf{g}_i$, since each $\mathbf{k}_\zeta$ is such an eigenfunction. It follows that for all $\mathbf{b} \in \mathcal{B}$,

$$\mathbf{g}_i(\mathbf{b}) = \frac{1}{\lambda} \mathbf{A}(\mathbf{g}_i)(\mathbf{b}) = \frac{1}{\lambda} \sum_{\mathbf{a} \sim \mathbf{b}} \mathbf{g}_i(\mathbf{a}).$$

Since the expression on the right only depends on vertices $\mathbf{a} \in \mathcal{A}$, we see $\mathbf{g}_1 = \mathbf{g}_2$. Therefore, the uniqueness part of the Poisson-Martin representation theorem implies $\sigma_1 = \sigma_2$. Finally, observe that if $s_1 = +1$ then $\nu_{\mathcal{A}}^- \equiv 0$. Therefore, $\nu_{\mathcal{A}}^+ + \mu_{\mathcal{A}}^- = \mu_{\mathcal{A}}^+$. But, the fact that $\mu_{\mathcal{A}}^- \perp \mu_{\mathcal{A}}^+$ implies that $\mu_{\mathcal{A}}^- \equiv 0$ and $\nu_{\mathcal{A}}^+ = \mu_{\mathcal{A}}^+$. Similarly, if $s_1 = -1$ then $\nu_{\mathcal{A}}^+ \equiv 0$, and so $\mu_{\mathcal{A}}^+ \equiv 0$ and $\nu_{\mathcal{A}}^- = \mu_{\mathcal{A}}^-$. In either case, $\mu_{\mathcal{A}} = \nu_{\mathcal{A}}$. An identical argument shows $\mu_{\mathcal{B}} = \nu_{\mathcal{B}}$, concluding the uniqueness part of the proof. $\square$

Corollary 6.47 defines a map $\mathcal{N} : \bigcup_{s \in SP} \widehat{E}_s \rightarrow M^2$ according to the rule $\mathcal{N}(\mathbf{f}) = (\nu_{\mathcal{A}}, \nu_{\mathcal{B}})$ with $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}})$ the unique pair of signed measures guaranteed to exist by the corollary.

The group of matrices $GL(2, \mathbb{R})$ acts on M^2 by matrix multiplication. Namely,

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \begin{bmatrix} \nu_{\mathcal{A}} \\ \nu_{\mathcal{B}} \end{bmatrix} = \begin{bmatrix} a\nu_{\mathcal{A}} + b\nu_{\mathcal{B}} \\ c\nu_{\mathcal{A}} + d\nu_{\mathcal{B}} \end{bmatrix}.$$

Therefore, our representation $\rho_\lambda : G \rightarrow SL(2, \mathbb{R})$ induces an action on M^2 .

Proposition 6.48 (Adjacency operation on measures). *Let $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ and suppose there exists $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}}) \in M^2$ so that*

$$\mathbf{f}(\mathbf{a}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{a}) \, d\nu_{\mathcal{A}}(\zeta) \quad \text{and} \quad \mathbf{f}(\mathbf{b}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{b}) \, d\nu_{\mathcal{B}}(\zeta)$$

for all $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$. Then

$$\mathbf{A}(\mathbf{f})(\mathbf{a}) = \lambda \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{a}) \, d\nu_{\mathcal{B}}(\zeta) \quad \text{and} \quad \mathbf{A}(\mathbf{f})(\mathbf{b}) = \lambda \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{b}) \, d\nu_{\mathcal{A}}(\zeta).$$

Proof. Define $\mathbf{f}_{\mathcal{A}}, \mathbf{f}_{\mathcal{B}} \in \mathbb{R}^{\mathcal{V}}$ according to the rule that

$$\mathbf{f}_{*}(\mathbf{v}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{v}) \, d\nu_{*}(\zeta) \quad \text{for all } \mathbf{v} \in \mathcal{V} \text{ and } * \in \{\mathcal{A}, \mathcal{B}\}.$$

Then $\mathbf{A}(\mathbf{f}_{*}) = \lambda \mathbf{f}_{*}$ since each $\mathbf{k}_{\zeta}$ satisfies $\mathbf{A}(\mathbf{k}_{\zeta}) = \lambda \mathbf{k}_{\zeta}$. Then we have $\mathbf{f} = \pi_{\mathcal{A}}(\mathbf{f}_{\mathcal{A}}) + \pi_{\mathcal{B}}(\mathbf{f}_{\mathcal{B}})$. It follows that

$$\mathbf{A}(\mathbf{f}) = \mathbf{A}(\pi_{\mathcal{A}}(\mathbf{f}_{\mathcal{A}}) + \pi_{\mathcal{B}}(\mathbf{f}_{\mathcal{B}})) = \pi_{\mathcal{B}} \circ \mathbf{A}(\mathbf{f}_{\mathcal{A}}) + \pi_{\mathcal{A}} \circ \mathbf{A}(\mathbf{f}_{\mathcal{B}}) = \pi_{\mathcal{B}}(\lambda \mathbf{f}_{\mathcal{A}}) + \pi_{\mathcal{A}}(\lambda \mathbf{f}_{\mathcal{B}}).$$

Therefore, the conclusion follows. $\square$

Proposition 6.49 (Group operation on measures). *Suppose $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ satisfies $\mathbf{A}^2(\mathbf{f}) = \lambda^2 \mathbf{f}$ and there exist $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}}) \in M^2$ so that*

$$\mathbf{f}(\mathbf{a}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{a}) \, d\nu_{\mathcal{A}}(\zeta) \quad \text{and} \quad \mathbf{f}(\mathbf{b}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{b}) \, d\nu_{\mathcal{B}}(\zeta)$$

for all $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$. Let $g \in G$, and set $(\mu_{\mathcal{A}}, \mu_{\mathcal{B}}) = \rho_{\lambda}^g \cdot (\nu_{\mathcal{A}}, \nu_{\mathcal{B}})$ and $\hat{\mathbf{f}} = \Upsilon^g(\mathbf{f})$. Then $\mathbf{A}^2(\hat{\mathbf{f}}) = \lambda^2 \hat{\mathbf{f}}$ and

$$\hat{\mathbf{f}}(\mathbf{a}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{a}) \, d\mu_{\mathcal{A}}(\zeta) \quad \text{and} \quad \hat{\mathbf{f}}(\mathbf{b}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{b}) \, d\mu_{\mathcal{B}}(\zeta)$$

for all $\mathbf{a} \in \mathcal{A}$ and $\mathbf{b} \in \mathcal{B}$.

Proof. First, the statement that $\mathbf{A}^2(\hat{\mathbf{f}}) = \lambda^2 \hat{\mathbf{f}}$ follows from equation 6.9. Namely,

$$\mathbf{A}^2(\hat{\mathbf{f}}) = \Upsilon^g \circ \mathbf{A}^2(\mathbf{f}) = \lambda^2 \Upsilon^g(\mathbf{f}) = \lambda^2 \hat{\mathbf{f}}.$$

We prove the remainder of the statement by induction. In fact, it is sufficient to prove it for g the identity and a collection of generators. The case of the identity is trivial. We consider the case of $g = h^k$ and $k \in \mathbb{Z}$. Then $(\mu_{\mathcal{A}}, \mu_{\mathcal{B}}) = (\nu_{\mathcal{A}} + k\nu_{\mathcal{B}}, \nu_{\mathcal{B}})$. We know that

$$\hat{\mathbf{f}} = \mathbf{H}^k(\mathbf{f}) = \mathbf{f} + k\pi_{\mathcal{A}} \circ \mathbf{A}(\mathbf{f}) = \mathbf{f} + k\pi_{\mathcal{A}} \circ \mathbf{A}(\mathbf{f}).$$

In this case, the conclusion follows from Proposition 6.48. The case of $g = v^K$ is similar. $\square$

Corollary 6.50. *Let $\mathbf{f} \in \hat{E}_s$ for some $s \in SP$. Suppose for some $g \in G$ we have $\Upsilon^g(\mathbf{f}) \in \hat{\mathcal{Q}}_{s'}$ for some $s' \in SP$. Then, $\Upsilon^g(\mathbf{f}) \in \hat{E}_{s'}$ and*

$$\mathcal{N}(\Upsilon^g(\mathbf{f})) = \rho_{\lambda}^g \cdot \mathcal{N}(\mathbf{f}).$$

Proof. Let $\hat{\mathbf{f}} = \Upsilon^g(\mathbf{f})$. Proposition 6.49 implies that $\mathbf{A}^2(\hat{\mathbf{f}}) = \lambda^2 \hat{\mathbf{f}}$. Therefore, $\hat{\mathbf{f}} \in \hat{E}_{s'}$. Set $(\mu_{\mathcal{A}}, \mu_{\mathcal{B}}) = \rho_{\lambda}^g \cdot \mathcal{N}(\mathbf{f}) \in M^2$. Proposition 6.49 indicates that

$$\hat{\mathbf{f}}(\mathbf{a}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{a}) \, d\mu_{\mathcal{A}}(\zeta) \quad \text{and} \quad \hat{\mathbf{f}}(\mathbf{b}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{b}) \, d\mu_{\mathcal{B}}(\zeta)$$

for all $a \in \mathcal{A}$ and $b \in \mathcal{B}$. By Corollary 6.47, the pair of measures that satisfy this statement are uniquely determined by $\hat{\mathbf{f}}$. Therefore, it must be that $\mathcal{N}(\hat{\mathbf{f}}) = (\mu_{\mathcal{A}}, \mu_{\mathcal{B}})$ as desired. $\square$

This corollary allows us to prove the following important lemma.

Lemma 6.51. *Suppose $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ satisfies $\mathbf{A}^2(\mathbf{f}) = \lambda^2 \mathbf{f}$ and is a $\boldsymbol{\theta}$ -survivor with $\boldsymbol{\theta} = (x, y) \in \mathcal{R}_{\lambda}$. Let $(\nu_{\mathcal{A}}, \nu_{\mathcal{B}}) = \mathcal{N}(\mathbf{f})$. Then, $\frac{1}{x}\nu_{\mathcal{A}} = \frac{1}{y}\nu_{\mathcal{B}}$.*

Proof. Let $\langle g_n \rangle$ be the λ -shrinking sequence of $\boldsymbol{\theta}$, and let $\langle s_n \rangle$ be the shrinking sequence. Since $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor, $\Upsilon^{g_n}(\mathbf{f}) \in \widehat{\mathcal{Q}}_{s_n}$. Moreover, by equation 6.9, $\mathbf{A}^2(\Upsilon^{g_n}(\mathbf{f})) = \lambda^2 \Upsilon^{g_n}(\mathbf{f})$. Therefore, $\Upsilon^{g_n}(\mathbf{f}) \in \widehat{E}_{s_n}$ for all n . Let $(\nu_{\mathcal{A},n}, \nu_{\mathcal{B},n}) = \mathcal{N} \circ \Upsilon^{g_n}(\mathbf{f})$. Write $s_n = (s_{n,1}, s_{n,2})$. By corollary 6.47, we have $\nu_{\mathcal{A},n} \in M^{s_{n,1}}$ and $\nu_{\mathcal{B},n} \in M^{s_{n,2}}$. Then by corollary 6.50, we have

$$(\nu_{\mathcal{A},n}, \nu_{\mathcal{B},n}) = \rho_{\lambda}^{g_n} \cdot (\nu_{\mathcal{A},0}, \nu_{\mathcal{B},0}).$$

Now let $A \subset \mathcal{M}_{\lambda}$ be any measurable subset. Let $\mathbf{v} = (\nu_{\mathcal{A},0}(A), \nu_{\mathcal{B},0}(B))$. Then we know $(\nu_{\mathcal{A},n}(A), \nu_{\mathcal{B},n}(A)) = \rho_{\lambda}^{g_n}(\mathbf{v})$ for all n . Moreover, since $\nu_{\mathcal{A},n} \in M^{s_{n,1}}$ and $\nu_{\mathcal{B},n} \in M^{s_{n,2}}$, we have $\rho_{\lambda}^{g_n}(\mathbf{v}) \in cl(Q_{s_n})$ for all n . Since $\langle g_n \rangle$ and $\langle s_n \rangle$ are λ -shrinking and sign sequences for a λ -renormalizable $\boldsymbol{\theta}$, by Proposition 6.18, we see that $\mathbf{v}$ must be a non-negative scalar multiple of $\boldsymbol{\theta}$. Equivalently, $\frac{1}{x}\nu_{\mathcal{A}}(A) = \frac{1}{y}\nu_{\mathcal{B}}(A)$. Since this is true for all measurable sets A , it must be that $\frac{1}{x}\nu_{\mathcal{A}} = \frac{1}{y}\nu_{\mathcal{B}}$. $\square$

A converse to Lemma 6.51 follows from Proposition 6.23. If ν is any Borel measure on $\mathcal{M}_{\lambda}$ with $\int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{v}) d\nu(\zeta)$ defined and finite for all $\mathbf{v} \in \mathcal{V}$, then the pair $(x\nu, y\nu) = \mathcal{N}(\mathbf{f})$ for a $\boldsymbol{\theta}$ -survivor $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$. Indeed, if we define $\mathbf{g} \in \mathbb{R}^{\mathcal{V}}$ according to the rule that $\mathbf{g}(\mathbf{v}) = \int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{v}) d\nu(\zeta)$ for all $\mathbf{v} \in \mathcal{V}$, then we see that $\mathbf{f} = P_{\mathbf{g}}(\boldsymbol{\theta})$ where $P_{\mathbf{g}}$ denotes the parametrized plane defined in section 6.4. Therefore, we have the following two results.

Corollary 6.52 (Survivors and the Martin boundary). *The linear map $\mathcal{N}$ restricts to a bijection from the space of $\boldsymbol{\theta}$ -survivors $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ satisfying $\mathbf{A}^2(\mathbf{f}) = \lambda^2 \mathbf{f}$ to pairs of measures of the form $(x\nu, y\nu)$ with ν an unsigned Borel measure on $\mathcal{M}_{\lambda}$ satisfying $\nu(\mathcal{M}_{\lambda} \setminus \mathcal{M}_{\lambda}^{min}) = 0$ and so that $\int_{\mathcal{M}_{\lambda}} \mathbf{k}_{\zeta}(\mathbf{v}) d\nu(\zeta)$ is defined and finite for all $\mathbf{v} \in \mathcal{V}$.*

We have been working in the specific case when $\boldsymbol{\theta}$ is a λ -renormalizable direction and $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$. But, the statements of our main results in section 4 concern two eigenfunctions with possibly different eigenvalues. In the context of the statements of these theorems we have the following.

Corollary 6.53 (Extremal survivors). *Suppose $\mathbf{w}_1 \in \mathbb{R}^{\mathcal{V}}$ satisfies $\mathbf{A}(\mathbf{w}_1) = \lambda_1 \mathbf{w}_1$ and assume $S(\mathcal{G}, \mathbf{w}_1)$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Let $\boldsymbol{\theta}_1$ be a λ_1 -renormalizable direction with λ_1 -shrinking sequence $\langle g_n \rangle$. Suppose $\mathbf{f} \in \mathcal{S}_{\boldsymbol{\theta}_1}$ is extremal. By Theorem 6.44, there is a λ_2 so that $\mathbf{A}(\mathbf{f}) = \lambda_2^2 \mathbf{f}$. Set $\boldsymbol{\theta}_2 = (x_2, y_2) = \boldsymbol{\theta}(\langle g_n \rangle, \lambda_2)$ as in Definition 3.7 and define $\mathbf{w}_2 \in \mathbb{R}^{\mathcal{V}}$ by*

$$\mathbf{w}_2(a) = \frac{1}{x_2} \mathbf{f}(a) \quad \text{and} \quad \mathbf{w}_2(b) = \frac{1}{y_2} \mathbf{f}(b) \quad \text{for } a \in \mathcal{A} \text{ and } b \in \mathcal{B}.$$

Then $\mathbf{w}_2$ is an extremal positive eigenfunction.

Note that $\mathbf{f} = P_{\mathbf{w}_2}(\boldsymbol{\theta}_2)$ by definition of $P_{\mathbf{w}_2}$ in section 6.4. This is the connection with the topological conjugacy construction.

Proof. Let $\mathbf{f} \in \mathcal{S}_{\theta_1}$ be extremal. The statement of the corollary defines $\mathbf{w}_2$. We only need to prove that $\mathbf{w}_2$ is an extremal positive eigenfunction. From the previous corollary with $\lambda = \lambda_2$ and $\theta = \theta_2$, we know that $\mathcal{N}(\mathbf{f}) = (x_2\nu, y_2\nu)$ for some Borel measure ν on $\mathcal{M}_\lambda$ satisfying $\nu(\mathcal{M}_\lambda \setminus \mathcal{M}_\lambda^{\min}) = 0$. Moreover, since $\mathcal{N}$ is a linear bijection, we know that since $\mathbf{f}$ is extremal, ν a Dirac measure supported on a single point $\zeta \in \mathcal{M}_\lambda$. Let $c = \nu(\{\zeta\})$. Then, since $\mathcal{N}(\mathbf{f}) = (x_2\nu, y_2\nu)$,

$$\mathbf{f}(\mathbf{a}) = cx_2\mathbf{k}_\zeta(\mathbf{a}) \quad \text{and} \quad \mathbf{f}(\mathbf{b}) = cy_2\mathbf{k}_\zeta(\mathbf{b}) \quad \text{for } \mathbf{a} \in \mathcal{A} \text{ and } \mathbf{b} \in \mathcal{B}.$$

Therefore by definition of $\mathbf{w}_2$, we have $\mathbf{w}_2 = c\mathbf{k}_\zeta$. Thus, $\mathbf{w}_2$ is an extremal positive eigenfunction. $\square$

We need to show that each measure associated to an extremal positive eigenfunction $\mathbf{w}_2$ is ergodic. This is essentially a converse of the above Corollary.

Lemma 6.54. *Assume $S(\mathcal{G}, \mathbf{w}_1)$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Let θ_1 be a λ_1 -renormalizable direction with λ_1 shrinking sequence $\langle g_n \rangle$. Suppose $\mathbf{w}_2$ is an extremal positive eigenfunction with eigenvalue λ_2 , and let $\theta_2 = \theta(\langle g_n \rangle, \lambda_2)$. Then the transverse measure $\mu_2 = \Psi_{\theta_1}^{-1} \circ \Xi(P_{\mathbf{w}_2}(\theta_2))$ is ergodic.*

Proof. By Proposition 6.23, the function $\mathbf{f} = P_{\mathbf{w}_2}(\theta_2)$ is a θ_1 -survivor. Therefore, by Lemma 6.2 and Theorem 6.3 there is a unique transverse measure $\mu_2 = \Psi_{\theta_1}^{-1} \circ \Xi(\mathbf{f})$. By Corollary 6.26, we know that μ_2 arises from the pullback of a homomorphism $\phi : S(\mathcal{G}, \mathbf{w}_1) \rightarrow S(\mathcal{G}, \mathbf{w}_2)$ as described in the Topological Conjugacy Theorem 4.1. In particular, Theorem 6.3 implies that the system consisting of the surface $S(\mathcal{G}, \mathbf{w}_1)$ and measure μ_2 is conservative. Hence the measure μ_2 has an ergodic decomposition. See Theorem 2.2.9 of [Aar97]. Recall that $\mathcal{M}_{\theta_1}$ denotes the collection of all locally finite transverse measures to the straight-line foliation of $S(\mathcal{G}, \mathbf{w}_1)$ in direction θ_1 . The ergodic decomposition yields a space Ω equipped with a probability measure m and a collection of ergodic transverse measures

$$\{\mu_\omega \in \mathcal{M}_{\theta_1} : \omega \in \Omega\}$$

(all depending on μ_2) so that for any measurable transversal $\tau \subset S(\mathcal{G}, \mathbf{w}_1)$ the map $\omega \mapsto \mu_\omega(\tau)$ is measurable and

$$\mu_2(\tau) = \int_{\Omega} \mu_\omega(\tau) \, dm(\omega).$$

For each $\mathbf{a} \in \mathcal{A}$, let $\sigma_{\mathbf{a}}$ be a vertical saddle connection crossing $\text{cyl}_{\mathbf{a}}$. Similarly for each $\mathbf{b} \in \mathcal{B}$, let $\sigma_{\mathbf{b}}$ be a horizontal saddle connection crossing the vertical cylinder $\text{cyl}_{\mathbf{b}}$. Note that since $\Psi_{\theta_1}(\mu_2) = \Xi(\mathbf{f})$ we have $|\mathbf{f}(\mathbf{a})| = \mu_2(\sigma_{\mathbf{a}})$ and $|\mathbf{f}(\mathbf{b})| = \mu_2(\sigma_{\mathbf{b}})$, with a sign depending only on the quadrant containing θ . In particular,

$$|\mathbf{f}(\mathbf{a})| = \int_{\Omega} \mu_\omega(\sigma_{\mathbf{a}}) \, dm(\omega) \quad \text{and} \quad |\mathbf{f}(\mathbf{b})| = \int_{\Omega} \mu_\omega(\sigma_{\mathbf{b}}) \, dm(\omega).$$

For each $\omega \in \Omega$ set $\mathbf{g}_\omega = \Xi^{-1} \circ \Psi_{\theta_1}(\mu_\omega)$. Each $\mathbf{g}_\omega$ is well defined and is a θ_1 -survivor by Corollary 6.30. Then $|\mathbf{g}_\omega(\mathbf{a})| = \mu_\omega(\sigma_{\mathbf{a}})$ and $|\mathbf{g}_\omega(\mathbf{b})| = \mu_\omega(\sigma_{\mathbf{b}})$, with the same sign considerations as before. Therefore, we have

$$\mathbf{f}(\mathbf{v}) = \int_{\Omega} \mathbf{g}_\omega(\mathbf{v}) \, dm(\omega) \quad \text{for all } \mathbf{v} \in \mathcal{V}.$$

The map $\Xi^{-1} \circ \Psi_{\theta_1}$ is a linear bijection $\mathcal{M}_{\theta_1} \rightarrow \mathcal{S}_{\theta_1}$. Therefore, each $\mathbf{g}_\omega$ is extremal in $\mathcal{S}_{\theta_1}$, because μ_ω is ergodic. From Theorem 6.44, it follows that there is an $\lambda_\omega > 0$ depending

on ω for which $\mathbf{A}^2(\mathbf{g}_\omega) = \lambda_\omega^2 \mathbf{g}_\omega$. Moreover the map $\omega \mapsto \lambda_\omega$ is measurable, since it can be computed as a ratio of sums of measures of saddle connections. Then for any integer k we have

$$(6.16) \quad \lambda_2^{2k} \mathbf{f}(\mathbf{v}) = \mathbf{A}^{2k}(\mathbf{f})(\mathbf{v}) = \int_{\Omega} \lambda_\omega^{2k} \mathbf{g}_\omega(\mathbf{v}) \, dm(\omega) \quad \text{for all } \mathbf{v} \in \mathcal{V},$$

where we make sense of negative of $\mathbf{A}$ powers by noting that $\mathbf{A}^2$ restricts to a bijection $\mathcal{S}_{\theta_1} \rightarrow \mathcal{S}_{\theta_1}$. See Theorem 6.42. Now suppose that

$$m(\{\omega : \lambda_\omega > \lambda_2\}) > 0.$$

Then we observe from the right side of equation 6.16 that

$$\lim_{k \rightarrow +\infty} (\mathbf{A}^{2k}(\mathbf{f})(\mathbf{v}))^{\frac{1}{2k}} > \lambda_2,$$

but this contradicts the left side of the equation which says that this limit must be λ_2 . Similarly, if

$$m(\{\omega : \lambda_\omega < \lambda_2\}) < 0 \quad \text{then} \quad \lim_{k \rightarrow -\infty} (\mathbf{A}^{2k}(\mathbf{f})(\mathbf{v}))^{\frac{1}{2k}} < \lambda_2,$$

which again is a contradiction. Thus m -almost everywhere we have $\lambda_\omega = \lambda_2$. Then as in corollary 6.53, we may set

$$\mathbf{w}_\omega(\mathbf{a}) = \frac{1}{x_2} \mathbf{f}(\mathbf{a}) \quad \text{and} \quad \mathbf{w}_\omega(\mathbf{b}) = \frac{1}{y_2} \mathbf{f}(\mathbf{b}) \quad \text{for all } \mathbf{a} \in \mathcal{A} \text{ and } \mathbf{b} \in \mathcal{B}.$$

Then by this corollary, $\mathbf{w}_\omega$ is an extremal positive eigenfunction with eigenvalue λ_2 m -a.e.. Since $\mathbf{f} = P_{\mathbf{w}_2}(\theta_2)$, we know that

$$\mathbf{w}_2(\mathbf{v}) = \int_{\Omega} \mathbf{w}_\omega(\mathbf{v}) \, dm(\omega) \quad \text{for all } \mathbf{v} \in \mathcal{V}.$$

But since $\mathbf{w}_2$ is an extremal positive eigenfunction and almost every $\mathbf{w}_\omega$ is an eigenfunction with the same eigenvalue, we have that $\mathbf{w}_\omega$ is a scalar multiple of $\mathbf{w}_2$ m -almost everywhere. Therefore m -a.e., $\mathbf{g}_\omega$ is a scalar multiple of $\mathbf{f}$. Since $\mu_\omega = \Psi_{\theta_1}^{-1} \circ \Xi(\mathbf{g}_\omega)$ and $\mu_2 = \Psi_{\theta_1}^{-1} \circ \Xi(\mathbf{f})$, we know m -a.e. that μ_ω is a scalar multiple of μ_2 . In particular, each μ_ω was known to be ergodic, so the measure μ_2 must be ergodic as well. $\square$

By combining the corollary and lemma above, we have the following.

Theorem 6.55. *Assume $S(\mathcal{G}, \mathbf{w}_1)$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Let θ_1 be a λ_1 -renormalizable direction with λ_1 shrinking sequence $\langle g_n \rangle$. Then the collection of locally finite ergodic invariant measures is given by*

$$\mu_2 = \Psi_{\theta_1}^{-1} \circ \Xi(P_{\mathbf{w}_2}(\theta_2))$$

where $\mathbf{w}_2$ varies over the extremal positive eigenfunctions of $\mathcal{G}$, $\theta_2 = \theta(\langle g_n \rangle, \lambda_2)$ and λ_2 is the eigenvalue of $\mathbf{w}_2$.

Proof. All locally finite ergodic invariant measures arise in this way by Corollary 6.53 and Corollary 6.30. Conversely, every such measure is ergodic by Lemma 6.54. $\square$

Finally, we can prove the ergodic measure classification theorem.

Proof of the Ergodic Measures Theorem II (Theorem 4.3). Since $\mathcal{G}$ has no vertices of valance one, by Theorem 6.39 we know that $S(\mathcal{G}, \mathbf{w}_1)$ has the critical decay property and $\mathcal{G}$ has the adjacency sign property. Therefore by Theorem 6.55 each ergodic measure is of the form $\mu_2 = \Psi_{\theta_1}^{-1} \circ \Xi(P_{\mathbf{w}_2}(\theta_2))$, where $\mathbf{w}_2$ is an extremal positive eigenfunction. Then it follows from Corollary 6.26 that μ_2 arises from a pullback construction as described in the statement of the theorem. $\square$

7. RENORMALIZABLE DIRECTIONS AND THE LIMIT SET

In this section, we will make concrete and prove some of the statements provided in section 3. We will also describe the connection between λ -renormalizable directions and the limit set.

Recall from Definition 3.1, given $g \in G$ there is a unique geodesic segment (in the word metric) $\langle g_0, \dots, g_n \rangle$ in G which satisfies $g_0 = e$ and $g_n = g$.

Definition 7.1 (Shrinking and expanding sets). For $g \in G$ define $Shr_\lambda(g)$ and $Exp_\lambda(g)$ to be the sets

$$\begin{aligned} Shr_\lambda(g) &= \{\theta \in \mathbb{S}^1 : \|\rho_\lambda^{g_n}(\theta)\| < \|\rho_\lambda^{g_{n-1}}(\theta)\| < \dots < \|\rho_\lambda^{g_1}(\theta)\| < \|\theta\| = 1\}. \\ Exp_\lambda(g) &= \{\theta \in \mathbb{S}^1 : \|\rho_\lambda^{g_n}(\theta)\| > \|\rho_\lambda^{g_{n-1}}(\theta)\| > \dots > \|\rho_\lambda^{g_1}(\theta)\| > \|\theta\| = 1\}. \end{aligned}$$

The next proposition follows from the observation that these two definitions are related by switching the directions of the inequalities. We define the projection map $\pi_{\mathbb{S}^1} : \mathbb{R}^2 \setminus \{\mathbf{0}\} \rightarrow \mathbb{S}^1$ to be the map $\pi_{\mathbb{S}^1}(\mathbf{v}) = \frac{\mathbf{v}}{\|\mathbf{v}\|}$.

Proposition 7.2. For all $g \in G$, $Shr_\lambda(g) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g^{-1}}(Exp_\lambda(g^{-1}))$.

Proof. The geodesic segment joining e to g^{-1} is the sequence $\langle g'_i = g_{n-i}g^{-1} \rangle$. Thus,

$$\begin{aligned} Exp_\lambda(g^{-1}) &= \{\theta \in \mathbb{S}^1 : \|\rho_\lambda^{g^{-1}}(\theta)\| > \|\rho_\lambda^{g_1 g^{-1}}(\theta)\| > \dots > \|\rho_\lambda^{g_{n-1} g^{-1}}(\theta)\| > \|\theta\|\} \\ &= \pi_{\mathbb{S}^1} \circ \rho_\lambda^g(\{\theta' \in \mathbb{S}^1 : \|\theta'\| > \|\rho_\lambda^{g_1}(\theta')\| > \dots > \|\rho_\lambda^{g_{n-1}}(\theta')\| > \|\rho_\lambda^{g_n}(\theta')\|\}) \\ &= \pi_{\mathbb{S}^1} \circ \rho_\lambda^g(Shr_\lambda(g)) \end{aligned}$$

This is equivalent to the version stated in the proposition. $\square$

Proposition 7.3. We have the following descriptions of the shrinking sets of generators.

$$\begin{aligned} Shr_\lambda(h) &= \{(x, y) \in \mathbb{S}^1 : \frac{-2}{\lambda} < \frac{y}{x} < 0\}. & Shr_\lambda(h^{-1}) &= \{(x, y) \in \mathbb{S}^1 : 0 < \frac{y}{x} < \frac{2}{\lambda}\}. \\ Shr_\lambda(v) &= \{(x, y) \in \mathbb{S}^1 : -\infty < \frac{y}{x} < \frac{-\lambda}{2}\}. & Shr_\lambda(v^{-1}) &= \{(x, y) \in \mathbb{S}^1 : \frac{\lambda}{2} < \frac{y}{x} < \infty\}. \end{aligned}$$

When $g \in \{h, v, h^{-1}, v^{-1}\}$ we have $Exp_\lambda(g) = \mathbb{S}^1 \setminus cl(Shr_\lambda(g))$, where cl denotes the closure.

We will not prove this proposition as it is a simple computation.

Recall the Definitions 6.14 and 6.15 of sign pairs and the corresponding open quadrants in $\mathbb{R}^2$. The following two corollaries can be verified by inspection of the above proposition. They will be relevant in later sections.

Corollary 7.4. Let $g \in \{h, h^{-1}, v, v^{-1}\}$ and $s \in SP$. Then the following statements are equivalent.

- (1) $\rho_\lambda^g(\mathcal{Q}_s) \subset \mathcal{Q}_s$.
- (2) The pair (g, s) belongs to the set

$$\{(h, ++), (h, --), (v, ++), (v, --), (h^{-1}, +-), (h^{-1}, -+), (v^{-1}, +-), (v^{-1}, -+)\}.$$

- (3) There is a $\mathbf{v} \in \mathcal{Q}_s$ for which $\|\rho_\lambda^{g^{-1}}(\mathbf{v})\| < \|\mathbf{v}\|$.
 (4) $\mathcal{Q}_s \subset \text{Exp}_\lambda(g)$.

Recall that $\gamma : G \rightarrow G$ is the group automorphism so that $\gamma(h) = v^{-1}$ and $\gamma(v) = h^{-1}$.

Corollary 7.5. *Let $g \in \{h, v, h^{-1}, v^{-1}\}$ and let $s \in SP$. If $\text{Shr}_\lambda(g) \cap \mathcal{Q}_s \neq \emptyset$ then $\mathcal{Q}_s \subset \text{Exp}_\lambda(\gamma(g))$.*

Corollary 7.6 (Unique shrinking generator). *Given any $\lambda \geq 2$ and $\boldsymbol{\theta} \in \mathbb{R}^2 \setminus \{\mathbf{0}\}$, there is at most one element $g \in \{h, h^{-1}, v, v^{-1}\}$ so that $\|\rho_\lambda^g(\boldsymbol{\theta})\| < \|\boldsymbol{\theta}\|$. Moreover, if $g_1 \in \{h, h^{-1}, v, v^{-1}\}$ satisfies $|\rho_\lambda^{g_1}(\boldsymbol{\theta})| < |\boldsymbol{\theta}|$ (resp. $|\rho_\lambda^{g_1}(\boldsymbol{\theta})| \leq |\boldsymbol{\theta}|$), then every $g_2 \in \{h, h^{-1}, v, v^{-1}\}$ with $g_1 \neq g_2$ satisfies $|\rho_\lambda^{g_2}(\boldsymbol{\theta})| > |\boldsymbol{\theta}|$ (resp. $|\rho_\lambda^{g_2}(\boldsymbol{\theta})| \geq |\boldsymbol{\theta}|$).*

Proof of Corollary 7.6. The proof follows from the fact that if g_1 and g_2 are distinct elements of $\{h, h^{-1}, v, v^{-1}\}$ then $\text{Shr}_\lambda(g_1) \subset \text{Exp}_\lambda(g_2)$ (resp. $\text{cl}(\text{Shr}_\lambda(g_1)) \subset \text{cl}(\text{Exp}_\lambda(g_2))$). This can be derived directly from Proposition 7.3. $\square$

Corollary 7.7 (Unique shrinking sequence). *Let $\boldsymbol{\theta} \in \mathbb{S}^1$. Then, there is at most one geodesic ray $\langle g_i \rangle$ for which the sequence $\|\rho_\lambda^{g_i}(\boldsymbol{\theta})\|$ is strictly monotone decreasing. Similarly, for all $n \geq 1$, there is at most one geodesic segment $\langle g_0, \dots, g_n \rangle$ for which $\|\rho_\lambda^{g_i}(\boldsymbol{\theta})\|$ is strictly monotone decreasing.*

Sketch of proof. Define the sequence inductively, so that $\gamma_i = g_i g_{i-1}^{-1} \in \{h, v, h^{-1}, v^{-1}\}$ is the unique generator for which $\|\rho_\lambda^{\gamma_i}(\rho_\lambda^{g_{i-1}}(\boldsymbol{\theta}))\| < \|\rho_\lambda^{g_{i-1}}(\boldsymbol{\theta})\|$. $\square$

Proposition 7.8. *Let $\boldsymbol{\theta} \in \mathbb{S}^1$. Let $\langle g_i \rangle$ be a geodesic segment or ray in G for which $\|\rho_\lambda^{g_{n+1}}(\boldsymbol{\theta})\| > \|\rho_\lambda^{g_n}(\boldsymbol{\theta})\|$ (resp. $\|\rho_\lambda^{g_{n+1}}(\boldsymbol{\theta})\| \geq \|\rho_\lambda^{g_n}(\boldsymbol{\theta})\|$) for some $n \geq 0$. Then, the subsequence $\|\rho_\lambda^{g_{n+j}}(\boldsymbol{\theta})\|$ for $j \geq 0$ is a strictly increasing (resp. nonstrictly increasing) sequence.*

Proof. We will prove the strictly increasing case. The nonstrict case follows similarly. The proof is by induction. Suppose $\|\rho_\lambda^{g_{i+1}}(\boldsymbol{\theta})\| > \|\rho_\lambda^{g_i}(\boldsymbol{\theta})\|$. Then $\pi_{\mathbb{S}^1}(\rho_\lambda^{g_{i+1}}(\boldsymbol{\theta})) \in \text{Shr}_\lambda(g_i g_{i+1}^{-1})$. Since $\langle g_i \rangle$ is a geodesic, $g_{i+2} g_{i+1}^{-1} \neq g_i g_{i+1}^{-1}$. By corollary 7.6, $\pi_{\mathbb{S}^1}(\rho_\lambda^{g_{i+1}}(\boldsymbol{\theta})) \in \text{cl}(\text{Exp}(g_{i+2} g_{i+1}^{-1}))$. In other words, $\|\rho_\lambda^{g_{i+2}}(\boldsymbol{\theta})\| \geq \|\rho_\lambda^{g_{i+1}}(\boldsymbol{\theta})\|$. $\square$

Corollary 7.9. *If $\boldsymbol{\theta} \in \mathbb{S}^1 \setminus (\text{Shr}_\lambda(h) \cup \text{Shr}_\lambda(h^{-1}) \cup \text{Shr}_\lambda(v) \cup \text{Shr}_\lambda(v^{-1}))$, then for all $g \in G$ we have $\|\boldsymbol{\theta}\| \leq \|\rho_\lambda^g(\boldsymbol{\theta})\|$.*

Proof. Apply proposition 7.8 to the geodesic segment joining e to g . $\square$

Extending Proposition 7.2, we have a description of the remaining shrinking and expanding sets.

Corollary 7.10. *Let $g \in G$ and let $\langle g_0 = e, \dots, g_n = g \rangle$ be the corresponding geodesic segment. For $\lambda \geq 2$, we have $\text{Exp}_\lambda(g) = \text{Exp}_\lambda(g_1)$ and*

$$\text{Shr}_\lambda(g) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g^{-1}}(\text{Exp}_\lambda(g_{n-1} g^{-1})) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g_{n-1}^{-1}}(\text{Shr}_\lambda(g g_{n-1}^{-1})).$$

Proof. The statement $\text{Exp}_\lambda(g_1) = \text{Exp}_\lambda(g)$ follows directly from Proposition 7.8. To prove the identity for $\text{Shr}_\lambda(g)$, we apply the equation for $\text{Exp}_\lambda(g)$ and Proposition 7.2:

$$\text{Shr}_\lambda(g) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g^{-1}}(\text{Exp}_\lambda(g^{-1})) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g_{n-1}^{-1}}(\text{Exp}_\lambda(g_{n-1} g^{-1})).$$

Finally by Proposition 7.2, we see $\text{Exp}_\lambda(g_{n-1} g^{-1}) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g g_{n-1}^{-1}}(\text{Shr}_\lambda(g g_{n-1}^{-1}))$. $\square$

We now recall some definitions from section 3. A $\theta \in \mathbb{S}^1$ is called λ -*shrinkable* if for all $\epsilon > 0$ there is a $g \in G$ such that $\|\rho_\lambda^g(\theta)\| < \epsilon$. A λ -*shrinking sequence* for θ is a geodesic ray $\langle g_i \rangle$ such that for each $\epsilon > 0$ there is an i such that $\|\rho_\lambda^{g_i}(\theta)\| < \epsilon$. Theorem 3.4 stated that λ -shrinking sequences of λ -shrinkable directions θ exist and are unique, and the associated sequence $\|\rho_\lambda^{g_i}(\theta)\|$ is strictly monotone decreasing.

Proof of Theorem 3.4. Let $\langle g_i \rangle$ be a λ -shrinking sequence for θ . We will first prove that $\|\rho_\lambda^{g_i}(\theta)\|$ is strictly monotone decreasing. Suppose not. Then, there is an n for which $\|\rho_\lambda^{g_{n+1}}(\theta)\| \geq \|\rho_\lambda^{g_n}(\theta)\|$. Thus, Proposition 7.8 implies that $\inf_i \|\rho_\lambda^{g_i}(\theta)\| > 0$, contradicting the definition of a λ -shrinking sequence. Thus, $\|\rho_\lambda^{g_i}(\theta)\|$ is strictly monotone decreasing. By Corollary 7.7, we know $\langle g_i \rangle$ is unique. $\square$

We now prove Proposition 3.5, which says that if $\langle g_i \rangle$ be a λ -shrinking sequence for θ , then the only distinct $\theta' \in \mathbb{S}^1$ for which $\langle g_i \rangle$ is also a λ -shrinking sequence for θ' is $\theta' = -\theta$.

Proof of Proposition 3.5. Suppose $\langle g_i \rangle$ is a λ -shrinking sequence for both θ and θ' and that their wedge product satisfies $\theta \wedge \theta' > 0$. For all i we have

$$(7.1) \quad 0 < \theta \wedge \theta' = \rho_\lambda^{g_i}(\theta) \wedge \rho_\lambda^{g_i}(\theta') \leq \|\rho_\lambda^{g_i}(\theta)\| \|\rho_\lambda^{g_i}(\theta')\|.$$

By definition, for all $\epsilon > 0$, there is an i and a j for which $\|\rho_\lambda^{g_i}(\theta)\| < \epsilon$ and $\|\rho_\lambda^{g_j}(\theta')\| < \epsilon$. By Theorem 3.4, both the sequences $\|\rho_\lambda^{g_i}(\theta)\|$ and $\|\rho_\lambda^{g_j}(\theta')\|$ are monotone decreasing. Therefore, if $k = \max(i, j)$ then $\|\rho_\lambda^{g_k}(\theta)\| \|\rho_\lambda^{g_k}(\theta')\| < \epsilon^2$. Therefore, $\|\rho_\lambda^{g_i}(\theta)\| \|\rho_\lambda^{g_i}(\theta')\| \rightarrow 0$ as $i \rightarrow \infty$. This contradicts equation 7.1. $\square$

We will now introduce some hyperbolic geometry and the limit set. For the necessary background, we refer the reader to the book of Matsuzaki and Taniguchi [MT98]. Recall the *hyperbolic plane* is $\mathbb{H}^2 = SO(2) \backslash SL(2, \mathbb{R})$. The boundary of the hyperbolic plane is naturally defined to be $\partial\mathbb{H}^2 = \mathbb{RP}^1 = \mathbb{R}^2 \setminus \{0\} / \mathbb{R}$. For $M \in SL(2, \mathbb{R})$ and $\mathbf{v} \in \mathbb{R}^2$, we will use $[M] \in \mathbb{H}^2$ and $[\mathbf{v}] \in \mathbb{RP}^1$ to denote the corresponding equivalence classes. To make $\mathbb{RP}^1$ the boundary of the hyperbolic plane, we say a sequence $\langle [M_n] \in \mathbb{H}^2 \rangle$ converges to $[\mathbf{v}] \in \mathbb{RP}^1$ if for any $[\mathbf{w}] \neq [\mathbf{v}] \in \mathbb{RP}^1$,

$$\frac{\|M_n \mathbf{v}\|}{\|M_n \mathbf{w}\|} \rightarrow 0 \text{ as } n \rightarrow \infty.$$

The *limit set*, $\Lambda(\Gamma) \subset \mathbb{RP}^1$ of a discrete group $\Gamma \subset SL(2, \mathbb{R})$ is the set of all limit points of sequences in $SO(2) \backslash \Gamma \subset \mathbb{H}^2$. The limit set $\Lambda(\Gamma) \subset \mathbb{RP}^1$ is the smallest non-empty closed invariant subset of $\mathbb{RP}^1$. An *open horodisk* in $\mathbb{H}^2$ at $[\mathbf{v}] \in \mathbb{RP}^1$ is a set of the form

$$\{[M] \in \mathbb{H}^2 : \|M \mathbf{v}\| < \epsilon\}$$

for some $\epsilon > 0$. The only accumulation point in $\mathbb{RP}^1$ of the horodisk defined above is $[\mathbf{v}]$. The *horospherical limit set* $\Lambda_h(\Gamma) \subset \mathbb{RP}^1$ is the set of all $[\mathbf{v}] \in \Lambda(\Gamma)$ for which any horodisk at $[\mathbf{v}]$ contains points in the orbit $[\Gamma] = \{[M] : M \in \Gamma\}$.

By recalling the definition of λ -shrinkable vectors given in section 3, we have the following tautology.

Proposition 7.11. *The direction $\theta \in \mathbb{S}^1$ is λ -shrinkable if and only if $[\theta] \in \Lambda_h(\rho_\lambda^G)$.*

We now recall a theorem of Beardon and Maskit [BM74, Theorem 2]. If $\Gamma \subset SL(2, \mathbb{R})$ is geometrically finite, then $\Lambda(\Gamma) \setminus \Lambda_h(\Gamma)$ is the collection of fixed points of parabolics.

Corollary 7.12. *If $[\theta] \in \Lambda(\rho_\lambda^G)$ then either θ is fixed by a parabolic in ρ_λ^G or θ is λ -shrinkable.*

We will now turn our attention to proving Theorem 3.6 which characterizes the set of all λ -shrinking sequences.

Proof of theorem 3.6. Powers of a parabolic in $SL(2, \mathbb{R})$ can not make any vector arbitrarily short. This explains why condition (1) is necessary when $\lambda \geq 2$ and why condition (2) is necessary when $\lambda = 2$.

Now consider the converse. Let $\langle g_i \rangle$ be any geodesic ray satisfying condition (1) and condition (2) when $\lambda = 2$. We must show that there is a $\theta \in \mathbb{S}^1$ such that $\langle g_i \rangle$ is a λ -shrinking sequence for θ . Note that the sequence of sets $Shr_\lambda(g_i)$ projectivize to a nested sequence of open intervals by Corollary 7.10 and

We break into two cases. First suppose we can find a $\theta \in \bigcap_i Shr_\lambda(g_i)$. Let $D \subset \mathbb{H}^2$ be the horodisk $\{[M] \in \mathbb{H}^2 : \|M\theta\| < \frac{11}{10}\}$. Note that $\|\rho_\lambda^{g_0}(\theta)\| = 1$, and recall that $\|\rho_\lambda^{g_i}(\theta)\|$ is monotone decreasing. Therefore, $[\rho_\lambda^{g_i}] \in D$ for all i . Thus by discreteness, $[\rho_\lambda^{g_i}] \rightarrow [\theta]$ in $\partial\mathbb{H}^2$. We conclude $[\theta] \in \Lambda(\rho_\lambda^G)$. We claim that $[\theta]$ is not fixed by any parabolic. Then the conclusion will follow by Corollary 7.12 and Proposition 7.3.

Suppose there is a $g \in G$ with $\rho_\lambda^g(\theta) = \pm\theta$. Any such $g \in G$ is conjugate to a power p of h or v (or hv^{-1} in the case $\lambda = 2$). Thus we can find a $\gamma \in G$ so that $g = \gamma p \gamma^{-1}$. Then, $\theta = \rho_\lambda^\gamma(\mathbf{v})$ with $\mathbf{v} = (x, 0)$ or $\mathbf{v} = (0, y)$ (or $\mathbf{v} = (x, x)$ in case $\lambda = 2$). Note that

$$\mathbf{v} \notin Shr_\lambda(h) \cup Shr_\lambda(h^{-1}) \cup Shr_\lambda(v) \cup Shr_\lambda(v^{-1}).$$

Therefore by Corollary 7.9, $\inf_{g \in G} \|\rho_\lambda^g(\theta)\| = \|\mathbf{v}\|$. Choose a $g' \in G$ of minimal distance from e and such that $\|\rho_\lambda^{g'}(\theta)\| = \|\mathbf{v}\|$. As the sequence $\|\rho_\lambda^{g_n}(\theta)\|$ was assumed to be decreasing, we see

$$\|\mathbf{v}\| = \|\rho_\lambda^{g'}(\theta)\| < \|\rho_\lambda^{g_n}(\theta)\| \text{ for all } n.$$

In particular, g' does not lie on the geodesic ray $\langle g_i \rangle$. Let $\langle g'_0 = e, \dots, g'_k = g' \rangle$ be the geodesic segment joining e to g' . Proposition 7.8 implies that $\|\rho_\lambda^{g'_j}(\theta)\|$ is strictly decreasing. The existence of the two distinct strictly monotone decreasing sequences $\langle g_i \rangle$ and $\langle g'_j \rangle$ contradicts Corollary 7.7.

In the second case, suppose that $\bigcap_i Shr_\lambda(g_i) = \emptyset$. A nested sequence of open intervals $\langle I_i \rangle$ has a non-empty intersection unless there is an N for which I_n shares a common endpoint for all $n > N$. So, we may suppose there is an N such that for all $n > N$ the intervals of the projectivization $[Shr_\lambda(g_n)]$ share a common endpoint. By applying Corollary 7.10 we see

$$[Shr_\lambda(g_n)] = [\rho_\lambda^{g_n^{-1}}(Shr_\lambda(g_n g_n^{-1}))].$$

Setting $i = n - N - 1$ we see $[Shr_\lambda(g_n)] = [\rho_\lambda^{g_N^{-1}} \circ \rho_\lambda^{g_N g_{N+1}^{-1}}(Shr_\lambda(g_n g_n^{-1}))]$. Thus, the intervals $J_i = [\rho_\lambda^{g_N g_{N+1}^{-1}}(Shr_\lambda(g_n g_n^{-1}))]$ all must share a common endpoint. Consider the set of endpoints

$$E = \partial[Shr_\lambda(h)] \cup \partial[Shr_\lambda(h^{-1})] \cup \partial[Shr_\lambda(v)] \cup \partial[Shr_\lambda(v^{-1})].$$

We see $J_0 = [Shr_\lambda(g_{N+1} g_N^{-1})]$. Thus, the common endpoint of the intervals J_i is some $e \in E$. Now consider that e must be an endpoint of $J_i = [\rho_\lambda^{g_N g_{N+1}^{-1}}(Shr_\lambda(g_n g_n^{-1}))]$. We conclude that $\rho_\lambda^{g_N g_{N+1}^{-1}}(e) \in E$. Let $g'_i = g_{N+i} g_N^{-1}$. In summary, $\langle g'_i \rangle$ is a geodesic ray with the property

that there is an $e \in E$ for which $\rho_\lambda^{g'_i}(e) \in E$ for all i . By inspection, it can be observed that

$$E \cap (\rho_\lambda^h(E) \cup \rho_\lambda^{h^{-1}}(E) \cup \rho_\lambda^v(E) \cup \rho_\lambda^{v^{-1}}(E)) = \begin{cases} \{(1, 0), (0, 1)\} & \text{if } \lambda > 2 \\ \{(1, 0), (0, 1), (1, 1), (-1, 1)\} & \text{if } \lambda = 2. \end{cases}$$

Moreover, the choice of an endpoint $e \in E$ nearly determines the geodesic ray $\langle g'_i \rangle$. For $\lambda \geq 2$ we have the following implications:

$$e = [(1, 0)] \text{ implies } g'_i = h^i \text{ or } g'_i = h^{-i}.$$

$$e = [(0, 1)] \text{ implies } g'_i = v^i \text{ or } g'_i = v^{-i}.$$

In other words, $\langle g_n \rangle$ is tail equivalent to one of the words in statement (1) of the theorem. When $\lambda = 2$ we also have the following four possible rules which determine $\langle g'_i \rangle$:

$$e = [(1, 1)] \text{ implies } g'_{i+1}g_i'^{-1} = \begin{cases} h & \text{if } i \text{ odd} \\ v^{-1} & \text{if } i \text{ even} \end{cases} \text{ or } g'_{i+1}g_i'^{-1} = \begin{cases} v & \text{if } i \text{ odd} \\ h^{-1} & \text{if } i \text{ even}. \end{cases}$$

$$e = [(-1, 1)] \text{ implies } g'_{i+1}g_i'^{-1} = \begin{cases} h^{-1} & \text{if } i \text{ odd} \\ v & \text{if } i \text{ even} \end{cases} \text{ or } g'_{i+1}g_i'^{-1} = \begin{cases} v^{-1} & \text{if } i \text{ odd} \\ h & \text{if } i \text{ even}. \end{cases}$$

In other words, $\langle g_n \rangle$ is tail equivalent to one of the geodesic rays in statement (2) of the theorem. $\square$

Proposition 7.13. *For all $\lambda \geq 2$, we have the following statements.*

$$Shr_\lambda(h^{-1}) \cap \mathcal{R}_\lambda \subset \{\theta = (x, y) \in \mathbb{S}^1 : 0 < \frac{y}{x} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\}.$$

$$Shr_\lambda(v^{-1}) \cap \mathcal{R}_\lambda \subset \{\theta = (x, y) \in \mathbb{S}^1 : \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} < \frac{y}{x} < \infty\}.$$

$$Shr_\lambda(h) \cap \mathcal{R}_\lambda \subset \{\theta = (x, y) \in \mathbb{S}^1 : \frac{-\lambda + \sqrt{\lambda^2 - 4}}{2} < \frac{y}{x} < 0\}.$$

$$Shr_\lambda(v) \cap \mathcal{R}_\lambda \subset \{\theta = (x, y) \in \mathbb{S}^1 : -\infty < \frac{y}{x} < \frac{-\lambda - \sqrt{\lambda^2 - 4}}{2}\}.$$

Proof. Note that when $\lambda = 2$, this statement is directly implied by Proposition 7.3. So, we will assume $\lambda > 2$. The quotient $\mathbb{H}^2/\rho_\lambda^G$ is homeomorphic to a three punctured sphere. Recall that the homotopy classes of loops on $\mathbb{H}^2/\rho_\lambda^G$ are in bijective correspondence with conjugacy classes in G . The conjugacy classes of h , v , and vh^{-1} correspond to simple loops traveling around each of the three punctures. As $\lambda > 2$, $\rho_\lambda^{vh^{-1}}$ is a hyperbolic isometry of $\mathbb{H}^2$ whose axis is the hyperbolic geodesic with whose endpoints are the projectivizations of the eigendirections of $\rho^{vh^{-1}}$. The endpoints of this geodesic are the points of $\mathbb{RP}^1$,

$$e_1 = [(2, \lambda - \sqrt{\lambda^2 - 4})] \quad \text{and} \quad e_2 = [(2, \lambda + \sqrt{\lambda^2 - 4})].$$

In particular, the geodesic joining e_1 to e_2 in $\mathbb{H}^2$ is the boundary of the convex core of the group ρ_λ^G , and the interval

$$I_1 = \{[(2, y)] : \lambda - \sqrt{\lambda^2 - 4} < y < \lambda + \sqrt{\lambda^2 - 4}\} \subset \mathbb{RP}^1$$

is a complimentary interval of $\mathbb{S}^1 \setminus \Lambda(\rho_\lambda^G)$. The set of such complimentary intervals is ρ_λ^G invariant, thus

$$I_2 = \rho_\lambda^{h^{-1}}(I_1) = \{[(2, y)] : \lambda - \sqrt{\lambda^2 - 4} < -y < \lambda + \sqrt{\lambda^2 - 4}\}$$

is also a complimentary interval. Now recall that $\mathcal{R}_\lambda \subset \Lambda(\rho_\lambda^G)$ by Proposition 7.11. We conclude for instance

$$\text{Shr}_\lambda(h) \cap \mathcal{R}_\lambda \subset \text{Shr}_\lambda(h) \setminus (\mathbb{S}^1 \setminus \tilde{\Lambda}(\rho_\lambda^G)) \subset \text{Shr}_\lambda(h) \setminus (\tilde{I}_1 \cup \tilde{I}_2),$$

where $\tilde{\Lambda}(\rho_\lambda^G)$, $\tilde{I}_1$ and $\tilde{I}_2$ denote the preimages of these subsets of $\mathbb{RP}^1$ under the projectivization $\mathbb{R}^2 \setminus \{0\} \rightarrow \mathbb{RP}^1$. Finally note that

$$\text{Shr}_\lambda(h) \setminus (\tilde{I}_1 \cup \tilde{I}_2) = \left\{ \theta = (x, y) \in \mathbb{S}^1 : 0 < \frac{-y}{x} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2} \right\}.$$

Similar statements hold for the other generators. $\square$

The following summarizes the connection between the limit set and the λ -renormalizable directions.

Proposition 7.14 ($\mathcal{R}_\lambda$ and the limit set). *Let $[\mathcal{R}_\lambda]$ denote the projectivization of the set of all λ -renormalizable directions.*

- In the case $\lambda = 2$, we have $\Lambda(\rho_2^G) = \mathbb{RP}^1$ and

$$[\mathcal{R}_2] = \mathbb{S}^1 \setminus \rho_2^G(\{[(1, 0)], [(0, 1)], [(1, 1)]\}).$$

- If $\lambda > 2$, the limit set $\Lambda(\rho_\lambda^G)$ is a Cantor set and

$$[\mathcal{R}_\lambda] = \Lambda(\rho_\lambda^G) \setminus \rho_\lambda^G(\{[(1, 0)], [(0, 1)], [(2, \lambda - \sqrt{\lambda^2 - 4})], [(2, \lambda + \sqrt{\lambda^2 - 4})]\}).$$

The orbits $\rho_\lambda^G(\{[(2, \lambda - \sqrt{\lambda^2 - 4})], [(2, \lambda + \sqrt{\lambda^2 - 4})]\})$ form the collection of endpoints of intervals in the complement of the Cantor set $\Lambda(\rho_\lambda^G)$.

8. INVARIANT MEASURES AND CODING

In this section, we rehash some of the arguments used to understand invariant measures of (finite) interval exchange maps. See [KH95, §14.5], for instance.

8.1. Coding trajectories. Let $S = \bigsqcup_{i \in \Lambda} P_i / \sim$ be a translation surface written as a union of polygons with edge identifications. Let E denote the set of all identified pairs of edges in $\partial P_i \subset S$ for $i \in \Lambda$. Choose θ so that it is not parallel to any edge in E .

The union of edges, $\bigcup_{e \in E} e$ is a section for the straight line flow in direction θ . We use $T_\theta : \bigcup_{e \in E} e \rightarrow \bigcup_{e \in E} e$ to denote the return map to this section. When equipped with the measure on $\bigcup_{e \in E} e$ induced by the Lebesgue transverse measure, this return map is essentially an interval exchange involving infinitely many intervals.

As with interval exchange transformations, we run into a problem concerning orbits which visit the endpoints of an edge $e \in E$. We resolve this by splitting all orbits which hit a singularity in two. Namely, if the forward or backward orbit of a point p hits an endpoint of an edge, we replace p with two new points p^- and p^+ . The orbit of p^- tracks the points to the left of p , and the orbit of p^+ tracks points to the right. (Left and right make sense, if we rotate the picture so that θ is vertical.) By *track* we mean to follow points infinitesimally nearby. We keep track of both the points location, and the edge it lies on. For $e \in E$, let $\hat{e}$ denote e with all points with singular orbits split as described above. Let l and r be the left and right endpoints of $e \in E$, respectively. We also replace l with l^+ and r with r^- in $\hat{e}$. We call $\hat{e}$ a *split edge*.

For each $e \in E$, there is a natural “unsplitting” map $\pi : \hat{e} \rightarrow e$. This map is surjective, and one-to-one except at countably many points, where it is two-to-one. We will be considering

the disjoint union of all split edges, $\bigsqcup_{e \in E} \hat{e}$. The map $\pi : \bigsqcup_{e \in E} \hat{e} \rightarrow \bigcup_{e \in E} \hat{e}$ may be countable-to-one at the vertices.

The return map T_θ has a natural lift to the map $\tilde{T}_\theta : \bigsqcup_{e \in E} \hat{e} \rightarrow \bigsqcup_{e \in E} \hat{e}$. In this system, $\tilde{T}_\theta^n(p)$ well defined for all $p \in \bigsqcup_{e \in E} \hat{e}$ and all $n \in \mathbb{Z}$. Let $edge : \bigsqcup_{e \in E} \hat{e} \rightarrow E$ denote the map which recovers the edge a point lies on. Now consider the coding map

$$(8.1) \quad code : \bigsqcup_{e \in E} \hat{e} \rightarrow E^{\mathbb{Z}} : x \mapsto \langle edge \circ \tilde{T}_\theta^n(x) \rangle_{n \in \mathbb{Z}}.$$

The image of this map is a shift space Ω on the countable alphabet E . Ω has the property that each symbol $e \in E$ is only followed by finitely many other symbols.

Proposition 8.1. *The coding map is injective if the straight line flow in direction θ is conservative and has no periodic trajectories.*

Proof. Suppose the straight line flow is conservative and has no periodic trajectories. We will show that distinct trajectories have distinct codes. Suppose we have a pair of distinct non-periodic trajectories. Then, there is an interval I of non-periodic trajectories with the same coding. Note that a trajectory's coding is periodic if and only if the trajectory is periodic. Since the straight line flow is conservative, almost all (and hence all) trajectories in I return infinitely often to some edge $e \in E$. But each intersection of I with e must be disjoint and of equal length. This contradicts the finite length of e . $\square$

8.2. Spaces of invariant measures. Given the conditions of proposition 8.1, the coding map given in equation 8.1 is a measure theoretic isomorphism from the space $\bigsqcup_{e \in E} \hat{e}$ to the shift space Ω . However, we would like to understand measures on $\bigcup_{e \in E} e$. These measure spaces are essentially the same.

Let $X \subset \bigcup_{e \in E} e$ be the set of points with more than one preimage under the unsplitting map π . We have the following.

Proposition 8.2. *Suppose the coding map is injective. Then, the map $\pi \circ code^{-1} : \Omega \rightarrow \bigcup_{e \in E} e$ induces an isomorphism between the following spaces of measures.*

- (1) *The shift invariant locally finite measures on Ω which assign zero measure to $code \circ \pi^{-1}(X)$.*
- (2) *The locally finite measures on $\bigcup_{e \in E} e$ which assign zero measure to X and which are invariant under T_θ .*

Note that the measure space in item (2) is what we want. It is the space of invariant measures that give no mass to singular trajectories. This space of measures is isomorphic to the space of transverse measures $\mathcal{M}_\theta$ defined in section 6.1. Namely, a measure in the space mentioned in item (2) is the restriction of a measure in $\mathcal{M}_\theta$ to the space of transversals contained in $\bigcup_{e \in E} e$.

Remark 8.3. Note that since X is countable, the spaces of non-atomic invariant measures on each space are isomorphic. But, the proposition even allows us to handle atomic measures, so long as the atoms are not contained in X .

8.3. Interaction with homology. In this section, we will consider the edges of the set E to be oriented. The homology classes of oriented edges generate $H_1(S, V, \mathbb{Z})$.

We recall some notation from section 6.1. The cohomology space $H^1(S, V, \mathbb{R})$ is the space of all linear maps $H_1(S, V, \mathbb{R}) \rightarrow \mathbb{R}$. Recall we defined a map $\Psi_\theta : \mathcal{M}_\theta \rightarrow H^1(S, V, \mathbb{R})$ in definition 6.1.

Lemma 8.4. *Suppose the coding map given in equation 8.1 is injective. Then the map Ψ_θ is also injective.*

Proof. We will utilize proposition 8.2 to translate the question to the shift space Ω . Note that the definition of the space $\mathcal{M}_\theta$, given in §2.1, allows us to apply the proposition without worrying about the space X .

By Carathéodory's extension theorem, a Borel measure on a shift space is determined by the measures of the cylinder sets. Let A be a cylinder set. Then, $\pi \circ \text{code}^{-1}(A)$ is a strip of trajectories which hit a sequence of specified edges of E . Such a strip of trajectories is precisely the collection of trajectories which cross some specific saddle connection σ on S . (Namely, the left and right sides of the strip must hit at least one vertex of an edge. Let P be a vertex on the left side and Q be a vertex on the right. Then $\sigma = \overline{PQ}$ is a saddle connection contained in the strip.) See figure 3. Thus, if μ is a shift invariant measure on Ω which assigns zero mass to $\text{code} \circ \pi^{-1}(X)$, and ν is the transverse measure on S corresponding to $\pi \circ \text{code}^{-1}(\mu)$, then $\mu(A) = \nu(\sigma)$.




FIGURE 3. This figure indicates how the measure of a cylinder set of Ω is determined by the measure of a saddle connection. Here the cylinder set is the set of trajectories which cross the sequence of edges $e_0, \dots, e_4$. The saddle connection σ is chosen so that it spans the width of the strip of trajectories which cross this sequence of edges.

Observe that for all saddle connections σ and all $\mu \in \mathcal{M}_\theta$, we have that

$$\mu(\sigma) = |\Psi_\theta(\mu)([\![\sigma]\!])|.$$

Therefore, we can recover the transverse measures of saddle connections from the image under Ψ_θ . Therefore the conclusion follows from Carathéodory's extension theorem and proposition 8.2. $\square$

Proof of lemma 6.2. If the straight line flow has no periodic trajectories and is conservative, the coding map is injective by proposition 8.1. By lemma 8.4, we know Ψ_θ is also injective. $\square$

Proof of lemma 6.4. The “only if” part of the lemma follows directly from Definition 6.1. The “if” part follows from Carathéodory’s extension theorem again. Assume $m \in H^1(S, V, \mathbb{R})$ satisfies

$$\operatorname{sgn}(m(\llbracket \sigma \rrbracket)) = \operatorname{sgn}(\operatorname{hol}(\sigma) \wedge \boldsymbol{\theta}) \quad \text{whenever } m(\llbracket \sigma \rrbracket) \neq 0.$$

By Carathéodory’s extension theorem, we can define a unique measure μ on Ω by determining its value on cylinder sets. Define f on the semi-ring of transversals generated by the saddle connections. We define $f(\sigma) = |m(\llbracket \sigma \rrbracket)|$. This function is finitely additive because of the sign condition. Thus it extends uniquely to a measure on Ω . The measure is invariant by the uniqueness provided by Carathéodory’s extension theorem. We can use proposition 8.2 to pull this measure back to a transverse invariant measure on S . (In general, we can find a non-unique pull back even if the coding map is not injective, but we would need to collect more detailed information about how the coding map fails to be injective in order to do this.) $\square$

9. CONSERVATIVITY AND APERIODICITY

In this section, we prove the straight-line flow in λ -renormalizable directions on surfaces $S(\mathcal{G}, \mathbf{w})$ is conservative and has no periodic trajectories when $\mathbf{w}$ is a positive function satisfying $\mathbf{A}(\mathbf{w}) = \lambda \mathbf{w}$.

The following two lemmas describe the technique we use to prove conservativity and the absence of periodic trajectories. For these two lemmas, let S be a translation surface and $\boldsymbol{\theta} \in S^1$ be a direction. Let μ be the Lebesgue transverse measure to the foliation in direction $\boldsymbol{\theta}$.

Lemma 9.1 (Criterion for conservativity). *Suppose that for all compact subsets $K \subset S$ and all $\epsilon > 0$, there is a compact subset $U \subset S$ such that $K \subset U$ and $\mu(\partial U) < \epsilon$. Then the straight line flow in direction $\boldsymbol{\theta}$ is conservative.*

The proof of this lemma was given in [Hoo08b]. The proof used ideas from [Tro04].

Lemma 9.2 (Criterion for aperiodicity). *Let V be a discrete set containing the singularities of S . Suppose that for all compact subsets $K \subset S$ and all $\epsilon > 0$, there is a compact subset $U \subset S$ such that $K \subset U$ and U has a decomposition into polygons $U = \bigcup_{m=1}^M Q_m$ such that each vertex of each Q_m lies in V and each edge σ satisfies $\mu(\sigma) < \epsilon$. Then the straight line flow in direction $\boldsymbol{\theta}$ has no periodic trajectories.*

Proof. Suppose there is a periodic trajectory. Then, there is a periodic cylinder. Let K be a cylinder which does not contain points of V in its interior. Let ϵ be the width of this cylinder. Then, we are given a U containing K and a decomposition into polygons $U = \bigcup_{i=1}^n Q_i$. Then K passes through some edge e of some Q_i . Thus, $\mu(e) \geq \epsilon$. This is a contradiction. $\square$

Theorem 9.3. *Let S be an infinite graph surface and let $\boldsymbol{\theta} \in S^1$ be a λ -renormalizable direction. The straight line flow in direction $\boldsymbol{\theta}$ is conservative, but has no periodic trajectories.*

We will find a nested sequence of compact sets $U_1 \subset U_2 \subset \dots \subset S$ to which we can apply the above two lemmas. We have the following.

- (1) If $\liminf_{n \rightarrow \infty} \mu(\partial U_n) = 0$ then the straight-line flow in direction $\boldsymbol{\theta}$ is conservative.
- (2) Each U_n admits a decomposition $U_n = \bigcup_{m=1}^M Q_{n,m}$ into polygons whose edges are saddle connections. Consider the sequence $e_n = \max_{\sigma} \mu(\sigma)$ where σ ranges over all edges of all $Q_{n,m}$ for $m = 1, \dots, M$. If $\liminf_{n \rightarrow \infty} e_n = 0$ then the straight line flow in direction $\boldsymbol{\theta}$ has no periodic trajectories.

We have $S = S(\mathcal{G}, \mathbf{w})$ where $\mathbf{w}$ is an eigenfunction of the adjacency operator with eigenvalue λ . By definition of a λ -renormalizable direction, we have a λ -renormalizable shrinking sequence $\langle g_0, g_1, \dots \rangle$ for $\boldsymbol{\theta}$ satisfying the five conditions of Theorem 3.6 as described in section 3. By an action of the dihedral group, we may assume that $g_1 = h$. See Remark 2.6. Our sequence of sets U_i will be defined using a natural subsequence $\gamma_n = g_{i(n)}$ of the shrinking sequence. We will inductively define this subsequence. We define $i(0) = 0$ so that $\gamma_0 = e$, the identity. Recall that $g_i \circ g_{i-1}^{-1} \in \{h, h^{-1}, v, v^{-1}\}$ for all $i \geq 1$. For $n > 0$, inductively define

$$i(n) = \min\{j > i(n-1) : g_j \circ g_{j-1}^{-1} \neq g_{j+1} \circ g_j^{-1}\}.$$

Such an $i(n)$ exists by statement (4) of Theorem 3.6. For example if

$$\langle g_i \rangle = \langle e, h, h^2, vh^2, v^2h^2, v^3h^2, h^{-1}v^3h^2, \dots \rangle \quad \text{then} \quad \langle \gamma_n \rangle = \langle e, h^2, v^3h^2, \dots \rangle.$$

In particular, $\gamma_n \circ \gamma_{n-1}^{-1}$ is a non-zero power of h when n is odd and a non-zero power of v when n is even.

For a vertex $v \in \mathcal{V}$, let $cyl_v \subset S$ denote the cylinder associated to v . Recall cyl_a is horizontal for $a \in \mathcal{A}$ and cyl_b is vertical for $b \in \mathcal{B}$. We will now define subsets $V_n \subset \mathcal{V}$. Recall $K \subset S$ is compact. Define

$$V_0 = \{a \in \mathcal{A} : cyl_a \cap K \neq \emptyset\}.$$

For $i > 0$ inductively define

$$V_n = \{v \in \mathcal{V} : v \sim w \text{ for some } w \in V_{n-1}\}.$$

Note that $V_n \subset \mathcal{A}$ for n even, and $V_n \subset \mathcal{B}$ for n odd. It can be observed inductively that each V_n is finite because K is compact.

Using the V_n we define subsets of $U_n \subset S$. Let $X_n = \bigcup_{v \in V_n} cyl_v$. Note that $X_{n-1} \subset X_n$ for all $n \geq 1$. We define

$$U_n = \Phi^{\gamma_n^{-1}}(X_n).$$

We can see that these sets are nested by noting that

$$U_n = \Phi^{\gamma_{n-1}^{-1}} \circ \Phi^{(\gamma_n \gamma_{n-1}^{-1})^{-1}}(X_n) \supset \Phi^{\gamma_{n-1}^{-1}} \circ \Phi^{(\gamma_n \gamma_{n-1}^{-1})^{-1}}(X_{n-1}) = U_{n-1}.$$

The last equality follows from two statements. First, X_{n-1} is a union of horizontal cylinders when n is odd, and a union of vertical cylinders when n is even. And second, $\gamma_n \gamma_{n-1}^{-1}$ is a power of h when n is odd, and a power of v when n is even. Thus, $\Phi^{(\gamma_n \gamma_{n-1}^{-1})^{-1}}(X_{n-1}) = X_{n-1}$ for all n .

The subsurface X_n comes with a natural decomposition into rectangles obtained by intersecting horizontal and vertical cylinders. Set

$$(9.1) \quad R_n = \{\text{rectangles } cyl_a \cap cyl_b : a \in \mathcal{A}, b \in \mathcal{B} \text{ and } cyl_a \cap cyl_b \subset X_n\}.$$

It follows that we get a corresponding decomposition of U_n into parallelograms. Define

$$(9.2) \quad Q_n = \{\Phi^{\gamma_n^{-1}}(cyl_a \cap cyl_b) : cyl_a \cap cyl_b \in R_n\}.$$

The U_n are getting larger with much longer boundary measured using Euclidean length. Similarly, the side lengths of the decomposition Q_n of U_n into parallelograms are very long. But, both ∂U_n and the sides of these parallelograms are getting closer to pointing in the direction $\boldsymbol{\theta}$. We will show that this convergence in direction is happening so fast that the Lebesgue transverse measure in direction $\boldsymbol{\theta}$ of ∂U_n decays to zero, as does the maximal measure of an edge in the decomposition Q_n .

Proposition 9.4 (Boundary growth). *Let ℓ_n denote the Euclidean length of ∂X_n . If $\lambda > 2$ then there is a constant c such that*

$$\ell_n \leq c \left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2} \right)^n$$

for all n . If $\lambda = 2$ then there is a c for which $\ell_n < cn$ for all n .

Claim 9.5. *Let $\lambda \geq 2$ and assume that $\langle a_0, a_1, \dots \rangle$ is a sequence of non-negative real numbers satisfying $a_{n+1} \leq \lambda a_n - a_{n-1}$. If $\lambda > 2$ then there is a constant c such that*

$$a_n \leq c \left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2} \right)^n$$

for all n . If $\lambda = 2$ then there is a c for which $a_n < cn$ for all n .

Proof. We consider the linear map $\phi(x, y) = (y, \lambda y - x)$. We note that (a_n, a_{n+1}) has the same x coordinate and non-strictly smaller y coordinate than $\phi(a_{n-1}, a_n)$. Let $\omega = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2}$. The eigenvectors of ϕ are $\mathbf{v}_1 = (1, \omega)$ and $\mathbf{v}_2 = (\omega, 1)$. They satisfy $\phi(\mathbf{v}_1) = \omega \mathbf{v}_1$ and $\phi(\mathbf{v}_2) = \omega^{-1} \mathbf{v}_2$. We first note that if $\frac{a_n}{a_{n-1}} < \omega^{-1}$ for some n , then there is no way to infinitely continue the sequence so that $a_n \geq 0$ for all n . This because for every point (x, y) with $\frac{y}{x} < \omega^{-1}$ is eventually ejected from the positive quadrant by a power of ϕ . This is illustrated in the left half of figure 4. Moreover, lowering the y -coordinate along the way, will only result in the vector being ejected from the quadrant faster.

Now assume $\lambda > 2$. By the above argument, we know that $\frac{a_n}{a_{n-1}} \geq \omega^{-1}$ for all n . Let P denote the closed parallelogram constructed from the convex hull of the points $(0, 0)$, $\omega^{-1} \mathbf{v}_2 = (1, \omega^{-1})$, $\mathbf{v}_1$ and $(0, \omega - 1)$. We note that $\phi(P) \subset \omega P$. See the right half of figure 4. Moreover

$$\{(x, y) : \exists (x_0, y_0) \in P \text{ such that } x = y_0, y < \lambda y_0 - x_0 \text{ and } \frac{y}{x} \geq \omega^{-1}\} \subset \omega(P).$$

We may assume that $(a_0, a_1) \in cP$ for some $c > 0$. By induction, we conclude that $(a_n, a_{n+1}) \in c\omega^n P$ for all n . The conclusion for $\lambda > 2$ follows.

When $\lambda = 2$, we set P_k to be the convex hull of the points $(0, 0)$, (k, k) , $(k, k + 1)$, and $(0, 1)$. We may check that $\phi(P_k) \subset P_{k+1}$ for all $k > 0$. Moreover,

$$\{(x, y) : \exists (x_0, y_0) \in P_k \text{ such that } x = y_0, y < 2y_0 - x_0 \text{ and } \frac{y}{x} \geq 1\} \subset P_{k+1}.$$

Assuming $(a_0, a_1) \in cP_0$, we have $(a_n, a_{n+1}) \in cP_n$ for all n , and the conclusion follows. $\square$

Proof of Proposition 9.4. We will show that the numbers ℓ_n are related by the inequality

$$(9.3) \quad \ell_{n+1} \leq \lambda \ell_n - \ell_{n-1}.$$

By Claim 9.5 above, this is sufficient to guarantee the result. Consider X_n with boundary of length ℓ_n . Let $W_n = V_{n+1} \setminus V_{n-1}$. The set of all cyl_v for $v \in W_n$ is the set of all horizontal or vertical cylinders which cross ∂X_n . (These cylinders are all horizontal if n is odd and all vertical if n is even.) As every such cylinder has inverse modulus λ , we know that

$$\sum_{v \in W_n} \text{length}(\partial cyl_v) \leq \lambda \ell_n.$$




FIGURE 4. This figure illustrates the action of ϕ from the proof of Claim 9.5 on $\mathbb{R}^2$ in the case of $\lambda = \frac{5}{2}$. The left two pictures illustrate a triangle T of points satisfying $\frac{y}{x} < \omega^{-1}$ being (eventually) ejected from the positive quadrant. The right two pictures indicate how $\phi(P) \subset \omega P$.

(Each such cylinder cyl_v for $v \in W_n$ crosses ∂X_n at least twice. A cylinder which crosses segments of length l , must have circumference λl .) Moreover,

$$X_{n+1} = X_{n-1} \cup \bigcup_{v \in W_n} cyl_v.$$

Explicitly, X_{n+1} is formed from X_{n-1} by attaching horizontal (or vertical) cylinders to the horizontal (or vertical) boundaries of X_{n-1} . Thus, we have that

$$\partial X_{n-1} \cup \partial X_{n+1} \subset \bigcup_{b \in W_n} \partial cyl_v.$$

The sets ∂X_{n-1} and ∂X_{n+1} are disjoint other than they may contain common singular points. Hence equation 9.3 holds. $\square$

Remark 9.6. These inequalities of proposition 9.4 are sharp, as can be seen for the case $\mathcal{G} = \mathbb{N}$. Moreover, whenever $\mathcal{G}$ is a tree, the inequality of equation 9.3 is actually equality. Therefore, by the dynamics of linear maps we know that there is a dichotomy when $\mathcal{G}$ is a tree and $\lambda > 2$. Either

- (1) $\lim_{n \rightarrow \infty} \ell_{n+1}/\ell_n = \frac{\lambda + \sqrt{\lambda^2 - 4}}{2}$, or
- (2) $\ell_{n+1}/\ell_n = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2} < 1$ for all $n \geq 0$.

By a similar argument, we have bounds on the lengths of boundary edges of rectangles in the decomposition of each X_n into rectangles formed by intersecting horizontal and vertical cylinders.

Proposition 9.7 (Rectangle decomposition growth). *Set $\mathcal{W}_n$ equal to the maximal Euclidean length of an edge of a rectangle in the decomposition, R_n , of X_n into rectangles given in equation 9.1. If $\lambda > 2$, there is a constant $c > 0$ for which*

$$\mathcal{W}_n \leq c \left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2} \right)^n.$$

If $\lambda = 2$, there is a constant $c > 0$ for which $\mathcal{W}_n \leq cn$.

Proof. Set $\mathcal{W}'_n = \sum_{v \in V_n \setminus V_{n-2}} \mathbf{w}(v)$. This is the sum of the widths of the cylinders in $X_n \setminus X_{n-2}$. For this to make sense when $n = 0, 1$, we take $V_{-2} = V_{-1} = \emptyset$ and $X_{-2} = X_{-1} = \emptyset$. We will prove the proposition with $\mathcal{W}_n$ replaced by $\mathcal{W}'_n$. The proposition as stated follows from the fact that $\mathcal{W}_N \leq \max_{n \leq N} \mathcal{W}'_n$. Now note that for $n \geq 3$, the vertices in $V_n \setminus V_{n-2}$ are adjacent only to those in $V_{n-1} \setminus V_{n-3}$ and $V_{n+1} \setminus V_{n-1}$. Because $\mathbf{w}$ is an eigenvector with eigenvalue λ , we have the inequality

$$\mathcal{W}'_{n+1} \leq \lambda \mathcal{W}'_n - \mathcal{W}'_{n-1}.$$

This implies the bound given in the statement of the proposition by Claim 9.5. $\square$

Now as $U_n = \Phi^{\gamma_n^{-1}}(X_n)$, we can compute the Lebesgue transverse measure of the boundary of U_n in terms of the unit vector $\boldsymbol{\theta}$ and ℓ_n . Recall that the derivative $D(\Phi^g) = \rho_\lambda^g$. From this observation, we have

$$(9.4) \quad \mu(\partial U_n) = \begin{cases} |\rho_\lambda^{\gamma_n^{-1}}(\ell_n, 0) \wedge \boldsymbol{\theta}| & \text{if } n \text{ is even} \\ |\rho_\lambda^{\gamma_n^{-1}}(0, \ell_n) \wedge \boldsymbol{\theta}| & \text{if } n \text{ is odd} \end{cases} = \begin{cases} |(\ell_n, 0) \wedge \rho_\lambda^{\gamma_n}(\boldsymbol{\theta})| & \text{if } n \text{ is even} \\ |(0, \ell_n) \wedge \rho_\lambda^{\gamma_n}(\boldsymbol{\theta})| & \text{if } n \text{ is odd}, \end{cases}$$

where $(a, b) \wedge (c, d) = ad - bc$ is the usual wedge product in the plane. In any case, we have

$$(9.5) \quad \mu(\partial U_n) \leq \ell_n \|\rho_\lambda^{\gamma_n}(\boldsymbol{\theta})\|.$$

The proof of conservativity proceeds by observing that $\|\rho_\lambda^{\gamma_n}(\boldsymbol{\theta})\|$ decays faster than ℓ_n grows along a subsequence, and therefore Lemma 9.1 applies.

The surfaces U_n inherit a decomposition $\mathcal{Q}_n$ into polygons as described in equation 9.2. Recall that $\mathcal{W}_n$ denoted the maximal width of a rectangle in the decomposition R_n of X_n into polygons. Set $\mathcal{Y}_n$ equal to the maximal Lebesgue transverse measure of an edge of a polygon in $\mathcal{Q}_n$. By the same argument used in the previous paragraph, we have

$$(9.6) \quad \mathcal{Y}_n \leq \mathcal{W}_n \|\rho_\lambda^{\gamma_n}(\boldsymbol{\theta})\|.$$

Again, if $\|\rho_\lambda^{\gamma_n}(\boldsymbol{\theta})\|$ decays faster than $\mathcal{W}_n$ grows along a subsequence, then Lemma 9.2 applies. We will now work to specify the subsequence which makes these results hold.

We define a collection of troublesome elements of the group G . Let

$$\mathcal{T} = \{(hv^{-1})^k\} \cup \{v^{-1}(hv^{-1})^k\} \cup \{(h^{-1}v)^k\} \cup \{v(h^{-1}v)^k\} \cup \\ \{(vh^{-1})^k\} \cup \{h^{-1}(vh^{-1})^k\} \cup \{(v^{-1}h)^k\} \cup \{h(v^{-1}h)^k\},$$

where k is allowed to range over the set $\{0, 1, 2, 3, \dots\}$. We define a subsequence $\langle \gamma_{n_j} \rangle$ by $n_0 = 0$ and

$$n_j = \min\{n_j > n_{j-1} : \gamma_{n_j} \gamma_{n_{j-1}}^{-1} \notin \mathcal{T}\}.$$

Note that n_j is well defined for all j so long as $\boldsymbol{\theta}$ is a λ -renormalizable direction. See statement (2) of Theorem 3.6. (If n_j is not well defined for some j , then $\langle g_n \rangle$ is tail equivalent to one of the sequences in statement (2).)

Given the above arguments, the following two lemmas imply the conservativity of the straight-line flow in direction of the unit vector $\boldsymbol{\theta}$ pointed in a λ -renormalizable direction. The first lemma handles the case of $\lambda = 2$, and the second handles the case of $\lambda > 2$ which is made more technical because we have to work with the varying values of λ . The proofs of both these lemmas use the same ideas.

Lemma 9.8. *There is a positive $\epsilon < 1$ depending such that for all j*

$$\eta(n_1)\eta(n_2 - n_1) \dots \eta(n_j - n_{j-1}) \|\rho_2^{\gamma_{n_j}}(\boldsymbol{\theta})\| \leq \epsilon^j \quad \text{for all } j,$$

where $\eta(1) = \frac{3}{2}$ and $\eta(n) = n$ for integers $n \geq 2$.

The statement of the lemma was chosen because it admits a recursive proof. Given the lemma, we get the type of decay we really want:

Corollary 9.9. *For all j , there is a positive $\epsilon < 1$ such that*

$$n_j \|\rho_2^{\gamma_{n_j}}(\boldsymbol{\theta})\| \leq \epsilon^j \quad \text{for all } j.$$

Proof. This follows from the fact that for any finite collection of positive integers,

$$n_1 + n_2 + \dots + n_k \leq \eta(n_1)\eta(n_2) \dots \eta(n_k).$$

We first prove this for $k = 2$. We break into cases:

- If $n_1 = n_2 = 1$ then $n_1 + n_2 = 2$ and $\eta(n_1)\eta(n_2) = \frac{9}{4} > 2$.
- If $n_1 = 1$ and $n_2 \geq 2$ then $\frac{\eta(n_1)\eta(n_2)}{n_1+n_2} = \frac{\frac{3}{2}n_2}{n_2+1} = \frac{3}{2} - \frac{3}{2(n_2+1)} \geq 1$, with equality when $n_2 = 2$.
- If $n_1 \geq 2$ and $n_2 \geq 2$ then $\eta(n_1)\eta(n_2) \geq 2 \max(n_1, n_2) \geq n_1 + n_2$.

Now suppose the statement is true for $k \geq 2$. Then,

$$\begin{aligned} n_1 + n_2 + \dots + n_k + n_{k+1} &\leq \eta(n_1 + \dots + n_k)\eta(n_{k+1}) = (n_1 + \dots + n_k)\eta(n_{k+1}) \\ &\leq \eta(n_1)\eta(n_2) \dots \eta(n_k)\eta(n_{k+1}). \end{aligned}$$

□

Proof of Lemma 9.8. To simplify notation, write $\delta_j = \gamma_{n_j}$. It is sufficient to prove the statement for the case of $j = 1$. To see this, suppose the statement is true for $j = 1$ and all λ -renormalizable directions $\boldsymbol{\theta}$. Then

$$\begin{aligned} \eta(n_1) \dots \eta(n_j - n_{j-1} + 1) \|\rho_2^{\delta_j}(\boldsymbol{\theta})\| &= \eta(n_1) \dots \eta(n_j - n_{j-1} + 1) \|\rho_2^{\delta_j \delta_1^{-1}} \circ \rho_2^{\delta_1}(\boldsymbol{\theta})\| \\ &= \left(\eta(n_1) \rho_2^{\delta_1}(\boldsymbol{\theta}) \right) \left(\eta(n'_1) \dots \eta(n'_{j-1} - n'_{j-2}) \|\rho_2^{\delta'_{j-1}}(\boldsymbol{\theta}')\| \right) \\ &\leq \epsilon \eta(n'_1) \dots \eta(n'_{j-1} - n'_{j-2}) \|\rho_2^{\delta'_{j-1}}(\boldsymbol{\theta}')\|. \end{aligned}$$

where $\boldsymbol{\theta}' = \rho_2^{\delta_1}(\boldsymbol{\theta}) / \|\rho_2^{\delta_1}(\boldsymbol{\theta})\|$, $n'_j = n_{j+1} - n_1$, and $\delta'_j = \delta_{j+1} \delta_1^{-1}$. Note that δ'_j arises from the shrinking sequence for $\boldsymbol{\theta}'$ in the same way in which δ_j arises from the shrinking sequence of $\boldsymbol{\theta}$. The inequality arises from the statement of the lemma in the case $j = 1$. We repeat this argument $j - 1$ more times to obtain the statement of the lemma.

We will now concentrate on proving the lemma in the case of $j = 1$. We note that the possible words δ_1 are of one of the following eight forms:

$$(9.7) \quad \begin{array}{cccc} v^{-a}(hv^{-1})^k & v^a(h^{-1}v)^k & h^{-a}(vh^{-1})^k & h^a(v^{-1}h)^k \\ h^av^{-1}(hv^{-1})^k & h^{-a}v(h^{-1}v)^k & v^ah^{-1}(vh^{-1})^k & v^{-a}h(v^{-1}h)^k \end{array}$$

Here $k \geq 0$ and $a \in \mathbb{Z} \setminus \{0, 1\}$. In addition, $(a, k) \neq (-1, 0)$ for words in the first row. By the previous paragraph, it is enough to prove that if δ_1 is one of these words then

$$(9.8) \quad \eta(n_1) \|\rho_2^{\delta_1}(\boldsymbol{\theta})\| \leq \epsilon < 1,$$

where $n_1 = 2k + 1$ if δ_1 is chosen from the first line of equation 9.7, or $n_1 = 2k + 2$ if δ_1 is chosen from the second. We simplify our job more by noting that many of these words are equivalent under the dihedral group. See Remark 2.6. Thus we really only need to cover one case from the first line and one case from the second.

We will prove the statement for the two cases $\delta_1 = h^{-a}(vh^{-1})^k$ and $\delta_1 = h^a v^{-1}(hv^{-1})^k$. In both cases either $v\delta_1$ or $v^{-1}\delta_1$ is an element of the shrinking sequence for $\boldsymbol{\theta}$. (Otherwise δ_1 would be reducible or a longer word.) Then by Corollary 7.10, we know that in either case

$$\boldsymbol{\theta} \in \text{Shr}_{v\delta_1} \cup \text{Shr}_{v^{-1}\delta_1} = \pi_{\mathbb{S}^1} \circ \rho_2^{\delta_1^{-1}}(\text{Shr}_v \cup \text{Shr}_{v^{-1}}).$$

For $\lambda = 2$, $\text{Shr}_v \cup \text{Shr}_{v^{-1}} \subset \{(x, y) \in \mathbb{S}^1 : -1 < \frac{x}{y} < 1\}$. Therefore, $\boldsymbol{\theta} \in \pi_{\mathbb{S}^1} \circ \rho_2^{\delta_1^{-1}}(x, \pm 1)$ for some choice of $-1 < x < 1$ and some choice of ± 1 . In particular it follows that

$$\|\rho_2^{\delta_1}(\boldsymbol{\theta})\| = \frac{\|(x, \pm 1)\|}{\|\rho_2^{\delta_1^{-1}}(x, \pm 1)\|}.$$

Therefore, it is sufficient to prove the statement that

$$(9.9) \quad \inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} \geq \frac{1}{\epsilon} > 1.$$

(We can drop the ± 1 by symmetry.)

We will now work out the case of $\delta_1 = h^{-a}(vh^{-1})^k$ where $k \geq 0$, $a \in \mathbb{Z} \setminus \{0, 1\}$ and $(a, k) \neq (-1, 0)$. We need to estimate the length of

$$\rho_2^{\delta_1^{-1}}(x, 1) = \rho_2^{(hv^{-1})^k} \circ \rho_2^{h^a}(x, 1).$$

Note that $\rho_2^{h^a}(x, 1) = (x + 2a, 1)$. We break into cases depending on a . Recall $-1 < x < 1$.

- (1) $x + 2a > 3$ when $a \geq 2$, or
- (2) $x + 2a < -1$ when $k \geq 1$ and $a \leq -1$.
- (3) $x + 2a < -3$ when $k = 0$ and $a \leq -2$.

In each of these cases, we have shown that $\rho_2^{h^a}(x, 1)$ is a subset of a ray contained in the line $L = \{(x', 1) : x' \in \mathbb{R}\}$. Now consider the matrix

$$\rho_2^{hv^{-1}} = \begin{bmatrix} -3 & 2 \\ -2 & 1 \end{bmatrix}.$$

This matrix is a parabolic with eigenvector $\mathbf{v} = (1, 1)$ satisfying $\rho_2^{hv^{-1}}(\mathbf{v}_1) = -\mathbf{v}_1$. Let $\mathbf{u} = (1, 0)$. Then $\rho_2^{hv^{-1}}(\mathbf{u}) = -\mathbf{u} - 2\mathbf{v}$. We see that $L = \{\mathbf{v} + t\mathbf{u} : t \in \mathbb{R}\}$. We compute that

$$\rho_2^{(hv^{-1})^k}(\mathbf{v} + t\mathbf{u}) = (-1)^k((2kt + 1)\mathbf{v} + t\mathbf{u}).$$

It can be checked that the point of $\rho_2^{(hv^{-1})^k}(L)$ which lies closest to the origin is $\rho_2^{(hv^{-1})^k}(\mathbf{v} + t\mathbf{u})$ for $-1 \leq t < 0$. This is the image under $\rho_2^{(hv^{-1})^k}$ of $\mathbf{v} + t\mathbf{u} = (x', 1)$ for some $0 \leq x' < 1$. In particular, this point is not in any of the rays. It follows that the infimum we need to find

to apply equation 9.9 is obtained at the endpoint of the ray. We must check this equation in each of the three cases. In case (1), the endpoint is $(3, 1) = \mathbf{v} + 2\mathbf{u}$. Therefore,

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|\rho_2^{(hv^{-1})^k}(\mathbf{v} + 2\mathbf{u})\|}{\eta(2k+1)\sqrt{2}} = \frac{\|(1+4k)\mathbf{v} + 2\mathbf{u}\|}{\eta(2k+1)\sqrt{2}}.$$

When $k = 0$, we have

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|(3, 1)\|}{\frac{3\sqrt{2}}{2}} = \frac{\sqrt{10}}{\frac{3\sqrt{2}}{2}} = \frac{\sqrt{20}}{3} > 1.$$

When $k \geq 1$, we know $\eta(2k+1) = 2k+1$. We apply the triangle inequality

$$\begin{aligned} \inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} &> \frac{(1+4k)\|\mathbf{v}\| - 2\|\mathbf{u}\|}{(2k+1)\sqrt{2}} = \frac{(1+4k)\sqrt{2}-2}{2k+1} \\ &= 2 - \frac{1+\sqrt{2}}{2k+1} \geq 2 - \frac{1+\sqrt{2}}{3} > 1. \end{aligned}$$

In case (2) the endpoint is $(-1, 1) = \mathbf{v} - 2\mathbf{u}$. We have $k \geq 1$. We check equation 9.9.

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|\rho_2^{(hv^{-1})^k}(\mathbf{v} - 2\mathbf{u})\|}{(2k+1)\sqrt{2}} = \frac{\|(1-4k)\mathbf{v} - 2\mathbf{u}\|}{(2k+1)\sqrt{2}}.$$

In case $k = 1$, we have $\|(1-4k)\mathbf{v} - 2\mathbf{u}\| = \|(-5, -3)\| = \sqrt{34}$. Therefore,

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\sqrt{34}}{4\sqrt{2}} = \frac{\sqrt{17}}{4} > 1.$$

In case $k \geq 2$, we can apply the triangle inequality.

$$\begin{aligned} \inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} &> \frac{\|(1-4k)\mathbf{v} - 2\mathbf{u}\|}{(2k+1)\sqrt{2}} > \frac{(4k-1)\sqrt{2}-2}{(2k+1)\sqrt{2}} = \frac{4k-1-\sqrt{2}}{2k+1} \\ &= 2 - \frac{3+\sqrt{2}}{2k+1} \geq 2 - \frac{3+\sqrt{2}}{5} > 1. \end{aligned}$$

Finally, we consider case (3). Here $k = 0$ and the ray endpoint is $(-3, 1)$ so,

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|(-3, 1)\|}{\frac{3}{2}\sqrt{2}} = \frac{\sqrt{20}}{3} > 1.$$

Now we consider the case, $\delta_1 = h^a v^{-1}(h v^{-1})^k$. We must estimate the length of

$$\rho_2^{\delta_1^{-1}}(x, 1) = \rho_2^{(vh^{-1})^k} \circ \rho_2^v \circ \rho_2^{h^{-a}}(x, 1).$$

Noting that $\rho_\lambda^{h^{-a}}(x, 1) = (x - 2a, 1)$, we break into two cases depending on the choice of $k \geq 0$ and $a \notin \{0, 1\}$.

- (1') $x - 2a < -3$ when $a \geq 2$, or
- (2') $x - a > 1$ when $a \leq -1$.

Let $\mathbf{v}$ and $\mathbf{u}$ be as in the previous paragraph. We have $\rho_2^{vh^{-1}}(\mathbf{v}) = -\mathbf{v}$ and $\rho_2^{vh^{-1}}(\mathbf{u}) = -(\mathbf{u} - 2\mathbf{v})$. Let $L = \{(x', 1)\}$. Set $L' = \rho_2^v(L) = \{(t, 1+2t) : t \in \mathbb{R}\}$. The closest point to the origin on L' is $(\frac{-2}{5}, \frac{1}{5})$. Now consider the closest point on $\rho_2^{(vh^{-1})^k}(L')$. Using the action of this parabolic, it can be checked that this closest point is of the form $\rho_2^{(vh^{-1})^k}(t, 1+2t)$

where $-1 < t \leq \frac{-2}{5}$. It follows that the closest point on $\rho_2^{(vh^{-1})^k v}(L)$ is also of this form. The preimage of this point lies in

$$\{\rho_2^{v^{-1}}(t, 1+2t) : -1 < t \leq \frac{-2}{5}\} = \{(t, 1) : -1 < t \leq \frac{-2}{5}\}.$$

Again we observe, that the rays in cases (1') and (2') do not contain such points. Therefore, the minimum length must occur at the endpoints of the rays. We now check equation 9.9 in case (1'). Here the endpoint is $(-3, 1)$ and $\rho_2^v(-3, 1) = (-3, -5) = -5\mathbf{v} + 2\mathbf{u}$. We simplify equation 9.9 using the triangle inequality.

$$\begin{aligned} \inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} &> \frac{\|\rho_2^{(vh^{-1})^k}(-5\mathbf{v}+2\mathbf{u})\|}{(2k+2)\sqrt{2}} = \frac{\|(-5-4k)\mathbf{v}+2\mathbf{u}\|}{(2k+2)\sqrt{2}} \\ &\geq \frac{(5+4k)\sqrt{2}-2}{(2k+2)\sqrt{2}} = 2 - \frac{\sqrt{2}-1}{2k+2} \geq 2 - \frac{\sqrt{2}-1}{2} > 1. \end{aligned}$$

In case (2'), the endpoint is $(1, 1)$ and $\rho_2^v(1, 1) = (1, 3) = 3\mathbf{v} - 2\mathbf{u}$. Therefore,

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|\rho_2^{(vh^{-1})^k}(3\mathbf{v} - 2\mathbf{u})\|}{(2k+2)\sqrt{2}} = \frac{\|(3+4k)\mathbf{v} - 2\mathbf{u}\|}{(2k+2)\sqrt{2}}.$$

When $k = 0$, we have $(3+4k)\mathbf{v} - 2\mathbf{u} = (1, 3)$. Therefore, in this case,

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{\|(3, 1)\|}{2\sqrt{2}} = \frac{\sqrt{5}}{2} > 1.$$

When $k \geq 1$, we apply the triangle inequality

$$\inf_{-1 < x < 1} \frac{\|\rho_2^{\delta_1^{-1}}(x, 1)\|}{\eta(n_1)\sqrt{2}} > \frac{(3+4k)\sqrt{2}-2}{(2k+2)\sqrt{2}} = \frac{3+4k-\sqrt{2}}{2k+2} = 2 - \frac{1+\sqrt{2}}{2k+2} \geq 2 - \frac{1+\sqrt{2}}{4} > 1.$$

This concludes the argument. The constant ϵ can be taken so that $\frac{1}{\epsilon}$ is the minimum of the finite number of constants used the cases above. $\square$

Lemma 9.10. *If $\lambda > 2$, there is a positive $\epsilon < 1$ depending only on λ such that*

$$\left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}\right)^{n_j} \|\rho_\lambda^{\gamma_{n_j}}(\boldsymbol{\theta})\| \leq \epsilon^j \quad \text{for all } j.$$

Proof. To simplify notation, write $\delta_j = \gamma_{n_j}$ and $\omega = (\lambda + \sqrt{\lambda^2 - 4})/2$. As in the previous case, it is in fact sufficient to prove the statement for $j = 1$. Suppose the statement for all λ -renormalizable directions $\boldsymbol{\theta}$ in the case of $j = 1$. Then we have

$$\omega^{n_{j+1}} \|\rho_\lambda^{\delta_{j+1}}(\boldsymbol{\theta})\| = \omega^{n_{j+1}} \|\rho_\lambda^{\delta_{j+1}\delta_1^{-1}} \circ \rho_\lambda^{\delta_1}(\boldsymbol{\theta})\| = \omega^{n_{j+1}} \|\rho_\lambda^{\delta_{j+1}\delta_1^{-1}}(\boldsymbol{\theta}')\| \|\rho_\lambda^{\delta_1}(\boldsymbol{\theta})\| < \epsilon \omega^{n'_j} \|\rho_\lambda^{\delta'_j}(\boldsymbol{\theta}')\|$$

where $\boldsymbol{\theta}' = \rho_\lambda^{\delta_1}(\boldsymbol{\theta})/\|\rho_\lambda^{\delta_1}(\boldsymbol{\theta})\|$, $n'_j = n_{j+1} - n_1$, and $\delta'_j = \delta_{j+1}\delta_1^{-1}$. Again, we can apply induction to obtain the statement of the lemma.

We now prove the lemma in the case of $j = 1$. As in the previous proof, it is sufficient to prove the lemma in the cases of $\delta_1 = h^{-a}(vh^{-1})^k$ and $\delta_1 = h^a v^{-1}(hv^{-1})^k$. Here, $k \geq 0$ and $a \in \mathbb{Z} \setminus \{0, 1\}$. For $\delta_1 = h^{-a}(vh^{-1})^k$ we can also assume $(a, k) \neq (-1, 0)$. It is enough to prove that if δ_1 is one of these words then

$$(9.10) \quad \omega^{n_1} \|\rho_\lambda^{\delta_1}(\boldsymbol{\theta})\| \leq \epsilon < 1.$$

Here $n_1 = 2k + 1$ if $\delta_1 = h^{-a}(vh^{-1})^k$ and $n_1 = 2k + 2$ if $\delta_1 = h^a v^{-1}(hv^{-1})^k$.

In both cases either $v\delta_1$ or $v^{-1}\delta_1$ is an element of the shrinking sequence for θ . Then by Corollary 7.10, we know that in either case

$$\theta \in \text{Shr}_{v\delta_1} \cup \text{Shr}_{v^{-1}\delta_1} = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{\delta_1^{-1}} (\text{Shr}_v \cup \text{Shr}_{v^{-1}}),$$

where $\pi_{\mathbb{S}^1}$ denotes projection onto the unit circle. From Proposition 7.13, we know

$$(9.11) \quad \mathcal{R}_\lambda \cap (\text{Shr}_v \cup \text{Shr}_{v^{-1}}) \subset \pi_{\mathbb{S}^1}(\{(x, \pm 1) : -\omega^{-1} < x < \omega^{-1}\}).$$

In particular, $\pi_{\mathbb{S}^1} \circ \rho_\lambda^{\delta_1^{-1}}(\theta)$ lies in this set. Without loss of generality, we may assume $\theta = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{\delta_1^{-1}}(x, 1)$ for some x with $-\omega^{-1} < x < \omega^{-1}$. Thus,

$$(9.12) \quad \begin{aligned} \|\rho_\lambda^{\delta_1}(\theta)\| &\leq \sup_{-\omega^{-1} < x < \omega^{-1}} \frac{\|(x, 1)\|}{\|\rho_\lambda^{\delta_1^{-1}}(x, 1)\|} \\ &\leq \sup_{-\omega^{-1} < x < \omega^{-1}} \frac{\sqrt{\lambda\omega^{-1}}}{\|\rho_\lambda^{\delta_1^{-1}}(x, 1)\|} = \frac{\sqrt{\lambda\omega^{-1}}}{\inf_{-\omega^{-1} < x < \omega^{-1}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\|}. \end{aligned}$$

The last inequality follows from the fact that $\|(x, 1)\| \leq \|(\omega^{-1}, 1)\| = \sqrt{\lambda\omega^{-1}}$. In particular from equation 9.10, it is sufficient to prove that

$$(9.13) \quad \inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| \geq \frac{1}{\epsilon} > 1.$$

for some $\epsilon < 1$.

Now we concentrate on the case $\delta_1 = h^{-a}(vh^{-1})^k$. We need to compute the infimum of $\rho_\lambda^{\delta_1^{-1}}(x, 1) = \rho_\lambda^{(hv^{-1})^k h^a}(x, 1)$ over those x satisfying $-\omega^{-1} < x < \omega^{-1}$. We have that $\rho_\lambda^{h^a}(x, 1) = (x + a\lambda, 1)$. Now, we break into cases depending on a and k . As $a \notin \{0, 1\}$, $k \geq 0$ and $(a, k) \neq (-1, 0)$, either

- (1) $x + a\lambda \geq 2\lambda - \omega^{-1} = \lambda + \omega$ when $a \geq 2$, or
- (2) $x + a\lambda \leq \omega^{-1} - \lambda = -\omega$ when $k \geq 1$ and $a \leq -1$.
- (3) $x + a\lambda \leq \omega^{-1} - 2\lambda = -\lambda - \omega$ when $k = 0$ and $a \leq -2$.

(For the equalities above, we use the identity $\omega + \omega^{-1} = \lambda$.) We will compute $\inf_{x'} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\|$, where x' ranges over the possible values of $x + a\lambda$ allowed by the inequalities in cases (1)-(3).

First consider the infimum over the line $\rho_\lambda^{(hv^{-1})^k}(\mathbb{R} \times \{1\})$. The two eigenvectors of $\rho_\lambda^{hv^{-1}}$ are

$$(9.14) \quad \mathbf{v}_1 = (\omega, 1) \quad \text{and} \quad \mathbf{v}_2 = (\omega^{-1}, 1),$$

with $\rho_\lambda^{hv^{-1}}(\mathbf{v}_1) = -\omega^2 \mathbf{v}_1$ and $\rho_\lambda^{hv^{-1}}(\mathbf{v}_2) = -\omega^{-2} \mathbf{v}_2$. Therefore,

$$\inf_{x' \in \mathbb{R}} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|\rho_\lambda^{(hv^{-1})^k}(x^\circ, 1)\|,$$

where $0 \leq x^\circ < \omega^{-1}$. As the inequalities above exclude the possibility of $x' = x^\circ$, we conclude that the infimum in cases (1)-(3) must be realized at the endpoints of the rays.

In case (1), we have

$$\inf_{x' \geq \lambda + \omega} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|\rho_\lambda^{(hv^{-1})^k}(\lambda + \omega, 1)\|.$$

We break into two subcases depending if $k = 0$ or $k \geq 1$. If $k = 0$, then

$$\inf_{x' \geq \lambda + \omega} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|(\lambda + \omega, 1)\| > \lambda + \omega$$

Applying this to equation 9.13 in this case yields

$$\inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| \geq \frac{\omega^{-1}}{\sqrt{\lambda\omega^{-1}}}(\lambda + \omega) = \sqrt{\frac{(\lambda + \omega)^2}{\lambda\omega}} > \sqrt{2}$$

Now consider $k \geq 1$. We have that $(\lambda + \omega, 1) = \frac{1}{\omega - \omega^{-1}}(2\omega\mathbf{v}_1 - \lambda\mathbf{v}_2)$, with $\mathbf{v}_i$ as in equation 9.14. Thus,

$$\begin{aligned} \|\rho_\lambda^{(hv^{-1})^k}(\lambda + \omega, 1)\| &= \frac{\omega^{2k+1}}{\omega - \omega^{-1}} \|2\mathbf{v}_1 - \lambda\omega^{-4k}\mathbf{v}_2\| \geq \frac{\omega^{2k+1}}{\omega - \omega^{-1}} (2\|v_1\| - \lambda\omega^{-4}\|v_2\|) \\ &= \frac{\omega^{2k+1}}{\omega - \omega^{-1}} (2\sqrt{\lambda\omega} - \lambda\omega^{-4}\sqrt{\lambda\omega^{-1}}) = \frac{\omega^{2k+1}\sqrt{\lambda\omega^{-1}}}{\omega - \omega^{-1}} (2\omega - \lambda\omega^{-4}). \end{aligned}$$

Applying this to equation 9.13 yields

$$\begin{aligned} \inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| &\geq \frac{\omega^{-2k-1}}{\sqrt{\lambda\omega^{-1}}} \left(\frac{\omega^{2k+1}\sqrt{\lambda\omega^{-1}}}{\omega - \omega^{-1}} (2\omega - \lambda\omega^{-4}) \right) \\ &= \frac{2\omega - \lambda\omega^{-4}}{\omega - \omega^{-1}} = \frac{\omega - \omega^{-1} + \lambda(1 - \omega^{-4})}{\omega - \omega^{-1}} > 1 \end{aligned}$$

In case (2), we have $k \geq 1$ and

$$\inf_{x' \leq -\omega} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|\rho_\lambda^{(hv^{-1})^k}(-\omega, 1)\|.$$

We may write $(-\omega, 1) = \frac{1}{\omega - \omega^{-1}}(-\lambda\mathbf{v}_1 + 2\omega\mathbf{v}_2)$. Therefore,

$$\begin{aligned} \|\rho_\lambda^{(hv^{-1})^k}(-\omega, 1)\| &= \frac{1}{\omega - \omega^{-1}} \|2\omega^{1-2k}\mathbf{v}_2 - \omega^{2k}\lambda\mathbf{v}_1\| \geq \frac{1}{\omega - \omega^{-1}} (\omega^{2k}\lambda\|\mathbf{v}_1\| - 2\omega^{1-2k}\|\mathbf{v}_2\|) \\ &= \frac{\omega^{2k+1}\sqrt{\lambda\omega^{-1}}}{\omega - \omega^{-1}} (\lambda - 2\omega^{-4k}) > \frac{\omega^{2k+1}\sqrt{\lambda\omega^{-1}}}{\omega - \omega^{-1}} (\lambda - 2\omega^{-4}). \end{aligned}$$

Applying this to equation 9.13 yields

$$\begin{aligned} \inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| &\geq \frac{\omega^{-2k-1}}{\sqrt{\lambda\omega^{-1}}} \left(\frac{\omega^{2k+1}\sqrt{\lambda\omega^{-1}}}{\omega - \omega^{-1}} (\lambda - 2\omega^{-4}) \right) \\ &= \frac{\lambda - 2\omega^{-4}}{\omega - \omega^{-1}} = \frac{\omega - \omega^{-1} + 2(\omega^{-1} - \omega^{-4})}{\omega - \omega^{-1}} > 1. \end{aligned}$$

Case (3) is simpler because $k = 0$. For δ_1 as in this case, we have that

$$\inf_{x' \leq -\lambda - \omega} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|(-\lambda - \omega, 1)\| > \lambda + \omega$$

and we can proceed as in the first subcase of case (1).

Now we consider the case, $\delta_1 = h^a v^{-1}(hv^{-1})^k$. Noting that $\rho_\lambda^{h^{-a}}(x, 1) = (x - a\lambda, 1)$, we break into two cases depending on the choice of $k \geq 0$ and $a \notin \{0, 1\}$.

(1') $x - a\lambda \leq \omega^{-1} - 2\lambda = -\lambda - \omega$ when $a \geq 2$, or

(2') $x - a\lambda \geq \lambda - \omega^{-1} = \omega$ when $a \leq -1$.

We will analyze $\inf_{x'} \|\rho_\lambda^{(vh^{-1})^k v}(x', 1)\|$, where x' ranges over the possible values of $x - a\lambda$ allowed by the inequalities in cases (1') and (2'). This time, we have that

$$\inf_{x' \in \mathbb{R}} \|\rho_\lambda^{(vh^{-1})^k v}(x', 1)\| = \|\rho_\lambda^{(vh^{-1})^k v}(x^\circ, 1)\|,$$

where $-\frac{\lambda}{1+\lambda^2} \leq x^\circ < \omega^{-1}$. This interval is in the complement of the set of values allowed for $x - a\lambda$ by cases (1) and (2), therefore the value of $\inf_{x'=x-a\lambda} \|\rho_\lambda^{(vh^{-1})^k v}(x', 1)\|$ is realized at the endpoints of the rays of these cases.

In case (1') we have,

$$\inf_{x' \leq -\lambda - \omega} \|\rho_\lambda^{(hv^{-1})^k}(x', 1)\| = \|\rho_\lambda^{(vh^{-1})^k v}(-\lambda - \omega, 1)\| = \|\rho_\lambda^{(vh^{-1})^k}(-\lambda - \omega, -\lambda^2 - \omega^2)\|.$$

As before $\mathbf{v}_1 = (\omega, 1)$ and $\mathbf{v}_2 = (\omega^{-1}, 1)$ are eigenvectors of $\rho_\lambda^{vh^{-1}}$, but here $\rho_\lambda^{vh^{-1}}(\mathbf{v}_1) = \omega^{-2}\mathbf{v}_1$ and $\rho_\lambda^{vh^{-1}}(\mathbf{v}_2) = \omega^2\mathbf{v}_2$. We write our vector in terms of these eigenvectors as

$$(-\lambda - \omega, -\lambda^2 - \omega^2) = \frac{\lambda}{\omega^3 - \omega}\mathbf{v}_1 + \frac{-2\omega^4}{\omega^2 - 1}\mathbf{v}_2.$$

Thus, we see

$$\rho_\lambda^{(vh^{-1})^k v}(-\lambda - \omega, 1) = \frac{\lambda}{\omega^3 - \omega}\omega^{-2k}\mathbf{v}_1 + \frac{-2\omega^4}{\omega^2 - 1}\omega^{2k}\mathbf{v}_2.$$

By the triangle inequality, we have

$$\|\rho_\lambda^{(vh^{-1})^k v}(-\lambda - \omega, 1)\| \geq \frac{2\omega^{2k+4}\|\mathbf{v}_2\|}{\omega^2 - 1} - \frac{\lambda\omega^{-2k-1}\|\mathbf{v}_1\|}{\omega^2 - 1} = \frac{\sqrt{\lambda\omega^{-1}}}{\omega^2 - 1}(2\omega^{2k+4} - \lambda\omega^{-2k}).$$

In this case, we apply equation 9.13 and see

$$\begin{aligned} \inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| &\geq \frac{\omega^{-2k-2}}{\sqrt{\lambda\omega^{-1}}} \left(\frac{\sqrt{\lambda\omega^{-1}}}{\omega^2 - 1} (2\omega^{2k+4} - \lambda\omega^{-2k}) \right) \\ &= \frac{2\omega^2 - \lambda\omega^{-4k-2}}{\omega^2 - 1} = \frac{2\omega^2 - \omega^{-4k-1} - \omega^{-4k-3}}{\omega^2 - 1} > \frac{2(\omega^2 - 1)}{\omega^2 - 1} = 2. \end{aligned}$$

In case (2'), we have the following calculations.

$$\inf_{x' \geq \omega} \|\rho_\lambda^{(vh^{-1})^k v}(x', 1)\| = |\rho_\lambda^{(vh^{-1})^k v}(\omega, 1)| = |\rho_\lambda^{(vh^{-1})^k}(\omega, \lambda\omega + 1)|.$$

We may write $(\omega, \lambda\omega + 1) = \frac{1}{\omega^2 - 1}(-2\mathbf{v}_1 + \omega^3\lambda\mathbf{v}_2)$. Therefore,

$$\rho_\lambda^{(vh^{-1})^k}(\omega, \lambda\omega + 1) = \frac{1}{\omega^2 - 1}(-2\omega^{-2k}\mathbf{v}_1 + \omega^{3+2k}\lambda\mathbf{v}_2).$$

By the triangle inequality, we have

$$|\rho_\lambda^{(vh^{-1})^k}(\omega, \lambda\omega + 1)| \leq \frac{\sqrt{\lambda\omega^{-1}}}{\omega^2 - 1}(\omega^{3+2k}\lambda - 2\omega^{1-2k}).$$

We apply equation 9.13 and see

$$\begin{aligned} \inf_{-\omega^{-1} < x < \omega^{-1}} \frac{\omega^{-n_1}}{\sqrt{\lambda\omega^{-1}}} \|\rho_\lambda^{\delta_1^{-1}}(x, 1)\| &\geq \frac{\omega^{-2k-2}}{\sqrt{\lambda\omega^{-1}}} \left(\frac{\sqrt{\lambda\omega^{-1}}}{\omega^2 - 1} (\omega^{3+2k}\lambda - 2\omega^{1-2k}) \right) \\ &= \frac{\omega\lambda - 2\omega^{-4k-1}}{\omega^2 - 1} = \frac{\omega^2 + 1 - 2\omega^{-4k-1}}{\omega^2 - 1} \\ &\geq \frac{\omega^2 + 1 - 2\omega^{-1}}{\omega^2 - 1}, \end{aligned}$$

and the fraction $\frac{\omega^2 + 1 - 2\omega^{-1}}{\omega^2 - 1}$ is strictly greater than one because $1 - 2\omega^{-1} > -1$. This completes the proof of equation 9.13 in all cases. $\square$

10. QUADRANTS AND SHRINKING SEQUENCES


10.1. Quadrants and expansion. Up to this point the λ -shrinking sequence $\langle g_i \rangle$ of a vector $\boldsymbol{\theta}$ has been characterized by the property that

$$\|\boldsymbol{\theta}\| > \|\rho_\lambda^{g_1}(\boldsymbol{\theta})\| > \|\rho_\lambda^{g_2}(\boldsymbol{\theta})\| > \dots$$

In this section, we will see that this condition is essentially equivalent to saying that each $\rho_\lambda^{g_i}(\boldsymbol{\theta})$ lies in some specific quadrant depending on the shrinking sequence. This section culminates with a proof of theorem 6.9.

Recall definitions 6.14 and 6.15 of sign pairs and quadrants in $\mathbb{R}^2$.

Definition 10.1 (Expanding sign action). The group action $\Sigma : G \times SP \rightarrow SP$ on the set of sign pairs is the action determined by the following diagram.



Now recall Definition 7.1 of $Exp_\lambda(g)$. Motivation for the definition of the expanding sign action comes from the following.

Proposition 10.2 (Quadrants and expansion). *Let $\langle g_i \rangle$ be a geodesic ray or segment and $s \in SP$. Suppose that $\theta \in cl(Exp_\lambda(g_1) \cap \mathcal{Q}_s)$. Define $s_i = \Sigma^{g_i}(s)$. Then $\rho_\lambda^{g_i}(\theta) \in cl(\mathcal{Q}_{s_i})$ for all i .*

Remark 10.3. The technical condition $\theta \in cl(Exp_\lambda(g_1) \cap \mathcal{Q}_s)$ allows the proposition to handle horizontal and vertical θ . If θ is not horizontal or vertical, the condition that $\theta \in cl(Exp_\lambda(g_1) \cap \mathcal{Q}_s)$ is equivalent to saying that $\theta \in \mathcal{Q}_s \cap \mathbb{S}^1$ and $\|\rho_\lambda^{g_1}(\theta)\| \geq \|\theta\|$.

Proof. First we will prove that the statement is true for $i = 1$. Any such statement must be invariant under the action of the dihedral group on $\langle g_i \rangle$ and θ . See Remark 2.6. Thus we may assume that $g_1 = h$ and $\theta \in cl(\mathcal{Q}_{++} \cup \mathcal{Q}_{--})$. We have $\theta \in Exp_\lambda(h) \cap (cl(\mathcal{Q}_{++} \cup \mathcal{Q}_{--}))$. Thus,

$$\rho_\lambda^h(\theta) \in \rho_\lambda^h(Exp_\lambda(h) \cap (cl(\mathcal{Q}_{++} \cup \mathcal{Q}_{--}))) \subset \mathcal{Q}_{++}.$$

This agrees with the fact that $\Sigma^h(-+) = \Sigma^h(++) = ++$. So, the proposition is true for $i = 1$, by dihedral group invariance.

To see the statement is true for $i > 1$, we apply induction. Proposition 7.8 implies that the sequence $\|\rho_\lambda^{g_i}(\theta)\|$ is non-strictly increasing. Assume $\rho_\lambda^{g_i}(\theta) \in cl(\mathcal{Q}_{s_i})$. We have $\|\rho_\lambda^{g_{i+1}}(\theta)\| \geq \|\rho_\lambda^{g_i}(\theta)\|$. Then by the first paragraph, $\rho_\lambda^{g_{i+1}}(\theta) \in cl(\mathcal{Q}_{s'})$, where $s' = \Sigma^{g_{i+1}g_i^{-1}}(s_i) = s_{i+1}$. $\square$

Proposition 10.4. *Let $g \in \{h, v, h^{-1}, v^{-1}\}$ and $s \in SP$. If $\rho_\lambda^g(\mathcal{Q}_s) \subset \mathcal{Q}_s$ then for any $\mathbf{v} \in \mathcal{Q}_s$ we have $\|\rho_\lambda^g(\mathbf{v})\| > \|\mathbf{v}\|$.*

Proof. By the dihedral group action, we may assume $g = h$. We have $\rho_\lambda^h(\mathcal{Q}_{++}) \subset \mathcal{Q}_{++}$ and $\rho_\lambda^h(\mathcal{Q}_{--}) \subset \mathcal{Q}_{--}$, but $\rho_\lambda^h(\mathcal{Q}_{+-}) \not\subset \mathcal{Q}_{+-}$ and $\rho_\lambda^h(\mathcal{Q}_{-+}) \not\subset \mathcal{Q}_{-+}$. By Proposition 7.3 we have $\mathcal{Q}_{++} \cup \mathcal{Q}_{--} \subset Exp_\lambda(h)$. $\square$

By combining these two propositions, we have the following.

Corollary 10.5. *Suppose $\langle g_i \rangle$ is a geodesic ray and $\rho_\lambda^{g_1}(\mathcal{Q}_s) \subset \mathcal{Q}_s$, or equivalently*


$$(g_1, s) \in \{(h, ++), (h, --), (v, ++), (v, --), (h^{-1}, +-), (h^{-1}, -+), (v^{-1}, +-), (v^{-1}, -+)\}.$$

Then for all i , $\rho_\lambda^{g_i}(cl(\mathcal{Q}_s)) \subset cl(\mathcal{Q}_{s_i})$ where $s_i = \Sigma^{g_i}(s)$.

Recall that given a λ -shrinkable direction $\theta \in \mathbb{S}^1$ with shrinking sequence $\langle g_0, g_1, g_2, \dots \rangle$, we can associate a sign sequence $\langle s_0, s_1, s_2, \dots \rangle$ with $s_i \in SP$ so that $\rho_\lambda^{g_i}(\theta) \in \mathcal{Q}_{s_i}$ for all i . See Definition 6.17. These two sequences are related by the following.

Proposition 10.6 (Shrinking and sign sequences). *Let $\langle g_i \rangle$ and $\langle s_i \rangle$ be shrinking sequences and sign pairs for a λ -shrinkable direction $\theta \in \mathbb{S}^1$. Then, there is a unique (infinite)*

path in the following diagram which begins at the node labeled $\mathcal{Q}_{s_0}$ and follows edges labeled $g_1, g_2g_1^{-1}, g_3g_2^{-2}$, et cetera. Moreover, an edge labeled $g_i g_{i-1}^{-1}$ joins the node $\mathcal{Q}_{s_{i-1}}$ to $\mathcal{Q}_{s_i}$.



Proof. We will prove that we can determine s_{i+1} from s_i , $g_{i+1}g_i^{-1}$ and $g_{i+2}g_{i+1}^{-1}$. (This is similar to the diagram; a node in a path is uniquely determined from the previous node and the labels on the adjacent arrows.) Consider the following observations.

- (1) We have $\rho_\lambda^{g_{i+1}}(\theta) \in \text{Shr}_\lambda(g_{i+2}g_{i+1}^{-1}) \cap \mathcal{Q}_{s_{i+1}}$. So the intersection $\text{Shr}_\lambda(g_{i+2}g_{i+1}^{-1}) \cap \mathcal{Q}_{s_{i+1}}$ is non-empty. Therefore, $g_{i+2}g_{i+1}^{-1} \in \{h, v\}$ implies $s_{i+1} \in \{+-, --\}$ and $g_{i+2}g_{i+1}^{-1} \in \{h^{-1}, v^{-1}\}$ implies $s_{i+1} \in \{++, --\}$.
- (2) Similarly, we have $\rho_\lambda^{g_{i+1}}(\theta) \in \rho_\lambda^{g_{i+1}g_i^{-1}}(\mathcal{Q}_{s_i}) \cap \mathcal{Q}_{s_{i+1}}$, so this intersection is non-empty. Therefore, we have the following.

$$\begin{aligned}
s_i = +- \text{ and } g_{i+1}g_i^{-1} = h &\text{ implies } s_{i+1} \in \{+-, --\}. \\
s_i = -+ \text{ and } g_{i+1}g_i^{-1} = h &\text{ implies } s_{i+1} \in \{++, +- \}. \\
s_i = +- \text{ and } g_{i+1}g_i^{-1} = v &\text{ implies } s_{i+1} \in \{++, +- \}. \\
s_i = -+ \text{ and } g_{i+1}g_i^{-1} = v &\text{ implies } s_{i+1} \in \{-+, -- \}. \\
s_i = ++ \text{ and } g_{i+1}g_i^{-1} = h^{-1} &\text{ implies } s_{i+1} \in \{++, +- \}. \\
s_i = -- \text{ and } g_{i+1}g_i^{-1} = h^{-1} &\text{ implies } s_{i+1} \in \{+-, -- \}. \\
s_i = ++ \text{ and } g_{i+1}g_i^{-1} = v^{-1} &\text{ implies } s_{i+1} \in \{++, +- \}. \\
s_i = -- \text{ and } g_{i+1}g_i^{-1} = v^{-1} &\text{ implies } s_{i+1} \in \{-+, -- \}.
\end{aligned}$$

These two observations combine to uniquely determine s_{i+1} as in the diagram. $\square$

10.2. Critical times for the shrinking sequence. Let $\langle s_n \rangle$ be the sign sequence of θ . We have the following important definition.

Definition 10.7. An integer $n \geq 1$ for which $s_{n-1} = s_n$ is a *critical time*.

There are various relationships between the sign sequence and the shrinking sequence $\langle g_n \rangle$ that appear at critical times.

Proposition 10.8. *The following statements are equivalent.*

- (1) *The integer $n > 0$ is a critical time.*
- (2) $\rho_\lambda^{g_{n-1}g_n^{-1}}(\mathcal{Q}_{s_n}) \subset \mathcal{Q}_{s_n}$.
- (3) *The pair $(g_{n-1}g_n^{-1}, s_n)$ lies in the set*

$$\{(h, ++), (h, --), (v, ++), (v, --), (h^{-1}, +-), (h^{-1}, -+), (v^{-1}, +-), (v^{-1}, -+)\}.$$
- (4) *There is a $\mathbf{v} \in \mathcal{Q}_{s_n}$ so that $\rho_\lambda^{g_n g_{n-1}^{-1}}(\mathbf{v}) \in \mathcal{Q}_{s_n}$.*
- (5) $\mathcal{Q}_{s_n} \subset \text{Exp}_\lambda(g_{n-1}g_n^{-1})$.
- (6) $g_{n+1}g_n^{-1} \in \{h^2, hv, vh, v^2, h^{-2}, h^{-1}v^{-1}, v^{-1}h^{-1}, v^{-2}\}$.

Proof. Let $\theta_n = \rho_\lambda^{g_n}(\theta)$. (1) implies (4) by taking $\mathbf{v} = \theta_n \in \mathcal{Q}_{s_n}$, because $\rho_\lambda^{g_{n-1}g_n^{-1}}(\theta_n) = \theta_n \in \mathcal{Q}_{s_n}$. Statement (2) through (5) can be seen to be equivalent by considering Corollary 7.4 with $g = g_{n-1}g_n^{-1}$ and $s = s_n$. Statement (2) implies (1) because $\theta_n \in \mathcal{Q}_{s_n}$ and $\theta_{n-1} = \rho_\lambda^{g_{n-1}g_n^{-1}}(\theta_n) \in \rho_\lambda^{g_{n-1}g_n^{-1}}(\mathcal{Q}_{s_n}) \subset \mathcal{Q}_{s_n}$. Therefore, $s_n = s_{n-1}$. Finally (1) implies (6) because $\|\theta_{n+1}\| < \|\theta_n\| < \|\theta_{n-1}\|$ and $\theta_n, \theta_{n-1} \in \mathcal{Q}_n$. Thus both $g_{n+1}g_n^{-1}$ and $g_n g_{n-1}^{-1}$ are labels arrows pointing away from $\mathcal{Q}_{s_n} = \mathcal{Q}_{s_{n-1}}$ in Proposition 10.6. It follows that their product is as stated in (6). Similarly, (6) implies (3), because $\mathcal{Q}_{s_n}$ has edges labeled $g_n g_{n-1}^{-1}$ and $g_{n+1}g_n^{-1}$ entering and leaving, respectively. Hence, s_n is as described in (3). $\square$

Corollary 10.9 (Critical times occur). *For $\theta \in \mathcal{R}_\lambda$, there are infinitely many critical times.*

Proof. We apply statement (6) of Proposition 10.8. Suppose the corollary is false for θ . Then for all but finitely many n , we would have $g_n g_{n-2}^{-1} \in \{h v^{-1}, h^{-1} v, v h^{-1}, v^{-1} h\}$. But, then $\langle g_i \rangle$ is tail equivalent to one of the sequences in statement (2) of Theorem 3.6. This contradicts the definition of a λ -renormalizable direction. $\square$

10.3. Interaction with the dot product. Recall the group automorphism $\gamma : G \rightarrow G$ defined as in equation 6.13. This automorphism has special significance for the representation ρ_λ^G .

Proposition 10.10. *For all λ and all $g \in G$, we have $\rho_\lambda^{\gamma(g)} = {}^t \rho_\lambda^{g^{-1}}$, the inverse transpose of ρ_λ^g .*

The proof follows by checking that it is true on the generators. We have a consequence analogous to equation 6.14. For all $\mathbf{v}, \mathbf{w} \in \mathbb{R}^2$ and all $g \in G$,

$$(10.1) \quad \mathbf{v} \cdot \mathbf{w} = \rho_\lambda^g(\mathbf{v}) \cdot \rho_\lambda^{\gamma(g)}(\mathbf{w}),$$

where $\cdot$ denotes the usual dot product $\mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$. We establish some simple corollaries of this observation.

Corollary 10.11 (Sign pairs at critical times). *Let $\langle g_i \rangle$ and $\langle s_i \rangle$ be shrinking sequences and sign pairs for a λ -shrinkable direction $\theta \in \mathbb{S}^1$. Suppose $t > 0$ is a critical time, then $s_t = \Sigma^{\gamma(g_t)}(s_0)$.*

Proof. Consider that $1 = \theta \cdot \theta = \rho_\lambda^{g_i}(\theta) \cdot \rho_\lambda^{\gamma(g_i)}(\theta)$ for all $i > 0$. Since $\|\rho_\lambda^{g_1}(\theta)\| < 1$ we know $\|\rho_\lambda^{\gamma(g_1)}(\theta)\| < 1$. Therefore, by proposition 10.2, we know $\rho_\lambda^{\gamma(g_i)}(\theta) \in \mathcal{Q}_{s'_i}$ where $s'_i = \Sigma^{\gamma(g_i)}(s_0)$ for all i . Since t is a critical time, we know $s_t = s_{t-1}$. By the dihedral group action, without loss of generality, we can assume that $s_{t-1} = s_t = ++$ and $g_t \circ g_t^{-1} = h^{-1}$. See Remark 2.6. Note that

$$s'_t \in \Sigma^{\gamma(h^{-1})}(SP) = \Sigma^v(SP) = \{++, --\}.$$

(See the diagram in Definition 10.1.) Since $\rho_\lambda^{\gamma(g_t)}(\theta) \in \mathcal{Q}_{s'_t}$, we know $s'_t = ++$ because $\rho_\lambda^{g_t}(\theta) \in \mathcal{Q}_{s_t} = \mathcal{Q}_{++}$ and $\rho_\lambda^{g_t}(\theta) \cdot \rho_\lambda^{\gamma(g_t)}(\theta) = 1$. $\square$

Corollary 10.12. *Suppose θ is λ -shrinkable, and $\mathbf{v} \in \mathbb{R}^2$ satisfies $\theta \cdot \mathbf{v} \neq 0$. Let $\langle g_n \rangle$ be the λ -shrinking sequence for θ . Then, $\|\rho_\lambda^{\gamma(g_n)}(\mathbf{v})\| \rightarrow \infty$ as $n \rightarrow \infty$.*

Proof. By equation 10.1, we have that $\theta \cdot \mathbf{v} = \rho_\lambda^{g_n}(\theta) \cdot \rho_\lambda^{\gamma(g_n)}(\mathbf{v})$. Therefore,

$$\|\rho_\lambda^{\gamma(g_n)}(\mathbf{v})\| \geq \frac{|\theta \cdot \mathbf{v}|}{\|\rho_\lambda^{g_n}(\theta)\|}.$$

The denominator of this expression tends to 0 as $n \rightarrow \infty$, so $\|\rho_\lambda^{\gamma(g_n)}(\mathbf{v})\| \rightarrow \infty$. $\square$

Of particular importance, we can conclude that if $\boldsymbol{\theta} \cdot \mathbf{v} \neq 0$ then eventually there is an i for which $\|\rho_\lambda^{\gamma(g_i)}(\mathbf{v})\| > \|\rho_\lambda^{\gamma(g_{i-1})}(\mathbf{v})\|$. Then for $n > i$, the quadrant containing $\rho_\lambda^{\gamma(g_n)}(\mathbf{v})$ is governed by the expanding sign action. See proposition 10.2. Then the following applies.

Proposition 10.13. *Let $\boldsymbol{\theta} \in \mathcal{R}_\lambda$. Let $\langle g_n \rangle$ be the λ -shrinking sequence of $\boldsymbol{\theta}$, and let $\langle s_n \rangle$ be the sign sequence. Suppose $\mathbf{v} \in \mathbb{R}^2$ satisfies $\boldsymbol{\theta} \cdot \mathbf{v} > 0$. By Corollary 10.12, there is an i for which $\|\rho_\lambda^{\gamma(g_i)}(\mathbf{v})\| > \|\rho_\lambda^{\gamma(g_{i-1})}(\mathbf{v})\|$. For any critical time $n \geq i$, we have $\rho_\lambda^{\gamma(g_n)}(\mathbf{v}) \in \mathcal{Q}_{s_n}$.*

Proof. Note that if $g \in \{h, v, h^{-1}, v^{-1}\}$ then the image $\Sigma^g(SP)$ consists of precisely two sign pairs. (For instance, $\Sigma^h(SP) = \{+-, -+\}$. See definition 10.1.) In particular, for $n \geq i$, we have $\rho_\lambda^{\gamma(g_n)}(\mathbf{v})$ lies in one of the two quadrants in $\Sigma^{g_n g_{n-1}^{-1}}(SP)$. Now suppose that n is a critical time. Statement (3) of Proposition 10.8 explicitly describes the possible pairs $(g_{n-1} g_n^{-1}, s_n)$. By inspection it can be observed that $\Sigma^{g_n g_{n-1}^{-1}}(SP) = \{s_n, -s_n\}$. Now recalling equation 10.1, we have that

$$\boldsymbol{\theta} \cdot \mathbf{v} = \rho_\lambda^{g_n}(\boldsymbol{\theta}) \cdot \rho_\lambda^{\gamma(g_n)}(\mathbf{v}).$$

Therefore $\boldsymbol{\theta} \cdot \mathbf{v} > 0$ and $\rho_\lambda^{g_n}(\boldsymbol{\theta}) \in \mathcal{Q}_{s_n}$ implies that $\rho_\lambda^{\gamma(g_n)}(\mathbf{v}) \in \mathcal{Q}_{s_n}$. $\square$

Now suppose that $\boldsymbol{\theta} \cdot \mathbf{v} = 0$. Let R denote the linear map $R : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ which rotates by $\frac{\pi}{2}$. As a matrix,

$$(10.2) \quad R = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Conjugation by this map induces an automorphism of ρ_λ^G . Inspection reveals that

$$(10.3) \quad R \circ \rho_\lambda^g = \rho_\lambda^{\gamma(g)} \circ R$$

for all $g \in G$. Note that this map R permutes the quadrants of $\mathbb{R}^2$ and therefore induces permutation $r : SP \rightarrow SP$ so that

$$(10.4) \quad R(\mathcal{Q}_s) = \mathcal{Q}_{r(s)}.$$

The punchline of this section is that we can detect the sign of $\boldsymbol{\theta} \cdot \mathbf{v}$ using the shrinking and sign sequences of $\boldsymbol{\theta}$.

Theorem 10.14. *Let $\boldsymbol{\theta} \in \mathcal{R}_\lambda$, and let $\langle g_n \rangle$ and $\langle s_n \rangle$ denote the λ -shrinking and sign sequences of $\boldsymbol{\theta}$, respectively. Choose any nonzero vector $\mathbf{v} \in \mathbb{R}^2$. Then,*

- (a) *If $\boldsymbol{\theta} \cdot \mathbf{v} > 0$, then there is an $i \geq 0$ such that for any critical time $n > i$ we have $\rho_\lambda^{\gamma(g_n)}(\mathbf{v}) \in \mathcal{Q}_{s_n}$.*
- (b) *If $\boldsymbol{\theta} \cdot \mathbf{v} < 0$, then there is an $i \geq 0$ such that for any critical time $n > i$ we have $-\rho_\lambda^{\gamma(g_n)}(\mathbf{v}) \in \mathcal{Q}_{s_n}$.*
- (c) *If $\boldsymbol{\theta} \cdot \mathbf{v} = 0$, then for all $i \geq 0$ we have $\rho_\lambda^{\gamma(g_n)}(\mathbf{v}) \in \mathcal{Q}_{r^{\pm 1}(s_n)}$ for some fixed choice of sign.*

Proof. Statement (a) follows directly from Proposition 10.13. Statement (b) follows by applying this proposition to $-\mathbf{v}$. Statement (c) follows from equation 10.3 by noting that

$$\rho_\lambda^{\gamma(g_n)} \circ R(\boldsymbol{\theta}) = R \circ \rho_\lambda^{g_n}(\boldsymbol{\theta}) \in \mathcal{Q}_{r(s_n)}.$$

$\square$

10.4. Proof of Theorem 6.9. The proof of theorem 6.9 follows almost directly from Theorem 10.14. We need to convert between the dot and wedge product. We recall that

$$(10.5) \quad \mathbf{v} \cdot \mathbf{w} = \mathbf{v} \wedge R(\mathbf{w}) = R^{-1}(\mathbf{v}) \wedge \mathbf{w}$$

with R defined as in equation 10.2. Recall that the Quadrant Sequence Proposition (Prop. 6.18) said that the only $\mathbf{v} \in \mathbb{S}^1$ for which $\rho_\lambda^{g_i}(\mathbf{v}) \in \overline{\mathcal{Q}_{s_i}}$ for all i is $\mathbf{v} = \boldsymbol{\theta}$. This statement is really a corollary of Theorem 10.14.

Proof of Proposition 6.18. Suppose such a $\mathbf{v}$ exists. Then $\mathbf{v} \wedge \boldsymbol{\theta} \neq 0$. Let $\mathbf{w} = R(\mathbf{v})$. Then,

$$\mathbf{v} \wedge \boldsymbol{\theta} = \boldsymbol{\theta} \cdot \mathbf{w} = \rho_\lambda^{g_n}(\boldsymbol{\theta}) \cdot \rho_\lambda^{\gamma(g_n)}(\mathbf{w}).$$

Since $\boldsymbol{\theta} \cdot \mathbf{w}$ is non-zero, Theorem 10.14 guarantees that there is a n for which $\pm \rho_\lambda^{\gamma(g_n)}(\mathbf{w}) \in \mathcal{Q}_{s_n}$, with the sign equal to the sign of $\boldsymbol{\theta} \cdot \mathbf{w}$. By equation 10.3, we have $\pm \rho_\lambda^{\gamma(g_n)}(\mathbf{w}) = \pm R \circ \rho_\lambda^{g_n}(\mathbf{v})$, so we may conclude that $\rho_\lambda^{g_n}(\mathbf{v}) \in \mathcal{Q}_{r(s_n)} \cup \mathcal{Q}_{r^{-1}(s_n)}$. This is a contradiction, since we assumed that $\rho_\lambda^{g_n}(\mathbf{v}) \in \overline{\mathcal{Q}_{s_n}}$. $\square$

Recall definition 6.8 of a $(\boldsymbol{\theta}, n)$ -survivor $m \in H^1(S, V, \mathbb{R})$. Theorem 6.9 stated that elements $m \in H^1(S, V, \mathbb{R})$ which arise from applying the Ψ function to locally finite invariant measures in direction $\boldsymbol{\theta}$ are $(\boldsymbol{\theta}, n)$ -survivors for all $n \geq 0$.

Proof of Theorem 6.9. We will apply Lemma 6.4. Let σ be a saddle connection in S , and let $\mathbf{v} = \text{hol } \sigma$. We will check that if $m(\llbracket \sigma \rrbracket) \neq 0$ then $\text{sgn}(m(\llbracket \sigma \rrbracket)) = \text{sgn}(\mathbf{v} \wedge \boldsymbol{\theta})$. We may assume that $\mathbf{v}$ is not parallel to $\boldsymbol{\theta}$, otherwise $m(\llbracket \sigma \rrbracket) = 0$ as saddle connections in direction $\boldsymbol{\theta}$ can not support invariant measures. Let $\langle g_n \rangle$ denote the shrinking sequence for $\boldsymbol{\theta}$, and $\langle s_n \rangle$ denote the sign sequence. By Proposition 6.18, there is a smallest n for which $\rho_\lambda^{g_n}(\mathbf{v}) \notin \mathcal{Q}_{s_n} \cup \mathcal{Q}_{-s_n}$. Let s denote the pair of signs for which $\rho_\lambda^{g_n}(\mathbf{v}) \in \mathcal{Q}_s$. Another way to write this condition is that $\mathbf{v} \notin \rho_\lambda^{g_n^{-1}}(\mathcal{Q}_{s_n} \cup \mathcal{Q}_{-s_n})$, which contains both $\pm \boldsymbol{\theta}$. The set of saddle connections whose homology classes are represented by $\Phi^{g_n^{-1}}(E)$ decompose S into parallelograms whose edges are parallel to $\rho_\lambda^{g_n^{-1}}(\mathbf{e}_1)$ and $\rho_\lambda^{g_n^{-1}}(\mathbf{e}_2)$. These classes form a basis for homology, and we can write

$$\llbracket \sigma \rrbracket = \sum_{i=1}^k \llbracket \sigma_i \rrbracket$$

where each $\llbracket \sigma_i \rrbracket \in \Phi^{g_n^{-1}}(E \cup -E)$ has holonomy which lies in the boundary $\rho_\lambda^{g_n^{-1}}(\partial \mathcal{Q}_s)$. As both $\boldsymbol{\theta}$ and $-\boldsymbol{\theta}$ do not lie in the closure of $\rho_\lambda^{g_n^{-1}}(\mathcal{Q}_s)$, we know that $\text{sgn}(\mathbf{w} \wedge \boldsymbol{\theta})$ is the same for all $\mathbf{w} \in \overline{\rho_\lambda^{g_n^{-1}}(\mathcal{Q}_s)}$. In particular, $\text{sgn}(\text{hol}(\sigma_i) \wedge \boldsymbol{\theta}) = \text{sgn}(\mathbf{v} \wedge \boldsymbol{\theta})$ for all i in the sum above. As m is a $(\boldsymbol{\theta}, n)$ -survivor, by definition we have either $m(\llbracket \sigma_i \rrbracket) = 0$ or

$$\text{sgn}(m(\llbracket \sigma_i \rrbracket)) = \text{sgn}(\text{hol}(\sigma_i) \wedge \boldsymbol{\theta}) = \text{sgn}(\mathbf{v} \wedge \boldsymbol{\theta}).$$

As each term in the sum has this same sign, assuming we have $m(\llbracket \sigma \rrbracket) \neq 0$ we conclude that

$$\text{sgn}(m(\llbracket \sigma \rrbracket)) = \text{sgn}\left(\sum_{i=1}^k m(\llbracket \sigma_i \rrbracket)\right) = \text{sgn}(\mathbf{v} \wedge \boldsymbol{\theta}),$$

so we may apply Lemma 6.4 as desired. $\square$

11. SURVIVORS AND OPERATORS

The purpose of this section is to prove Theorem 6.28, which says that all θ -survivors in $H^1(S, V, \mathbb{Z})$ arise as $\Xi(\mathbf{f})$ where $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ is a θ -survivor. In order to prove this statement, we will transform several results proved in section 10 about the interplay between the ρ_λ^G action on $\mathbb{R}^2$ and quadrants in $\mathbb{R}^2$ into statements about the Φ_*^G action on $H^1(S, V, \mathbb{R})$ and the Υ^G action on $\mathbb{R}^{\mathcal{V}}$.

11.1. Homology, Cohomology and $\mathbb{R}^{\mathcal{V}}$. We begin by proving Proposition 6.12, which says that $\Phi_*^g \circ \Xi = \Xi \circ \Upsilon^g$.

Proof of Proposition 6.12. Let $\llbracket x \rrbracket \in H_1(S, V, \mathbb{Z})$. By definition of Φ_*^g and of Ξ ,

$$(\Phi_*^g \circ \Xi(\mathbf{f}))(\llbracket x \rrbracket) = \Xi(f)(\Phi^{g^{-1}}(\llbracket x \rrbracket)) = \sum_{v \in \mathcal{V}} i(\Phi^{g^{-1}}(\llbracket x \rrbracket), \llbracket cyl_v \rrbracket) \mathbf{f}(v).$$

By acting by Φ^g on each side of each expression for intersection number, we have

$$(\Phi_*^g \circ \Xi(\mathbf{f}))(\llbracket x \rrbracket) = \sum_{v \in \mathcal{V}} i(\llbracket x \rrbracket, \Phi^g(\llbracket cyl_v \rrbracket)) \mathbf{f}(v).$$

Consider the case of $g = h^k$ with $k \in \mathbb{Z}$. By Proposition 6.7, we have

$$\Phi^{h^k}(\llbracket cyl_v \rrbracket) = \begin{cases} \llbracket cyl_v \rrbracket + k \sum_{w \sim v} \llbracket cyl_w \rrbracket & \text{if } v \in \mathcal{A} \\ \llbracket cyl_v \rrbracket & \text{if } v \in \mathcal{B}. \end{cases}$$

Thus, we may write

$$(\Phi_*^{h^k} \circ \Xi(\mathbf{f}))(\llbracket x \rrbracket) = \sum_{a \in \mathcal{A}} i(\llbracket x \rrbracket, \llbracket cyl_a \rrbracket) \mathbf{f}(a) + \sum_{b \in \mathcal{B}} i(\llbracket x \rrbracket, \llbracket cyl_b \rrbracket + k \sum_{a \sim b} \llbracket cyl_a \rrbracket) \mathbf{f}(b).$$

By regrouping terms, we see

$$(\Phi_*^{h^k} \circ \Xi(\mathbf{f}))(\llbracket x \rrbracket) = \sum_{a \in \mathcal{A}} i(\llbracket x \rrbracket, \llbracket cyl_a \rrbracket) (\mathbf{f}(a) + k \sum_{b \sim a} \mathbf{f}(b)) + \sum_{b \in \mathcal{B}} i(\llbracket x \rrbracket, \llbracket cyl_b \rrbracket) \mathbf{f}(b)$$

This last expression is equal to $\sum_{v \in \mathcal{V}} i(\llbracket x \rrbracket, \llbracket cyl_v \rrbracket) (\mathbf{H}^k(\mathbf{f}))(\mathbf{v}) = (\Xi \circ \mathbf{H}^k(\mathbf{f}))(\llbracket x \rrbracket)$. Thus, $\Xi \circ \mathbf{H}^k = \Phi_*^{h^k} \circ \Xi$ as desired. The argument is similar for the case of $g = v^k$. $\square$

Recall that $\mathbb{R}_c^{\mathcal{V}}$ represents the set of compactly supported functions from $\mathcal{V} \rightarrow \mathbb{R}$. We introduce a canonical linear map $\mathcal{Z} : H_1(S, V, \mathbb{R}) \rightarrow \mathbb{R}_c^{\mathcal{V}}$. Define

$$(11.1) \quad \mathcal{Z}(\llbracket x \rrbracket)(v) = i(\llbracket x \rrbracket, \llbracket cyl_v \rrbracket),$$

where $\llbracket cyl_v \rrbracket \in H_1(S \setminus V, \mathbb{Z})$ represents the homology class of the core curve of cylinder cyl_v for $v \in \mathcal{V}$. Recalling the bilinear pairing $\langle, \rangle : \mathbb{R}^{\mathcal{V}} \times \mathbb{R}_c^{\mathcal{V}} \rightarrow \mathbb{R}$ given in equation 6.11. By definition of Ξ (see equation 6.5), we have

$$(11.2) \quad \langle \mathbf{f}, \mathcal{Z}(\llbracket x \rrbracket) \rangle = \Xi(\mathbf{f})(\llbracket x \rrbracket).$$

Note that $\mathcal{Z}$ is not injective. A useful consequence of the construction is that

$$\mathcal{Z}(\llbracket x \rrbracket) = \mathcal{Z}(\llbracket y \rrbracket) \quad \text{implies} \quad \Xi(\mathbf{f})(\llbracket x \rrbracket) = \Xi(\mathbf{f})(\llbracket y \rrbracket).$$

We collect the following corollary to Proposition 6.12.

Corollary 11.1. *For all $\llbracket x \rrbracket \in H_1(S, V, \mathbb{R})$, we have $\Upsilon^g \circ \mathcal{Z}(\llbracket x \rrbracket) = \mathcal{Z} \circ \Phi^{\gamma(g)}(\llbracket x \rrbracket)$, where $\gamma : G \rightarrow G$ is the involutive homomorphism defined above equation 6.14.*

Proof. Note that two elements $\mathbf{x}, \mathbf{y} \in \mathbb{R}_c^\vee$ are equal if and only if $\langle \mathbf{f}, \mathbf{x} \rangle = \langle \mathbf{f}, \mathbf{y} \rangle$ for all $\mathbf{f} \in \mathbb{R}^\vee$. Let $\mathbf{f} \in \mathbb{R}^\vee$ be arbitrary. By equation 6.14,

$$\langle \mathbf{f}, \Upsilon^g \circ \mathcal{Z}(\llbracket x \rrbracket) \rangle = \langle \Upsilon^{\gamma(g)^{-1}}(\mathbf{f}), \mathcal{Z}(\llbracket x \rrbracket) \rangle.$$

Then by equation 11.2 and Proposition 6.12,

$$\langle \mathbf{f}, \Upsilon^g \circ \mathcal{Z}(\llbracket x \rrbracket) \rangle = \Xi(\Upsilon^{\gamma(g)^{-1}}(\mathbf{f}))(\llbracket x \rrbracket) = (\Phi_*^{\gamma(g)^{-1}} \circ \Xi(\mathbf{f}))(\llbracket x \rrbracket).$$

By the definition of $\Phi_*^{\gamma(g)^{-1}}$ given in equation 6.1, we continue

$$\langle \mathbf{f}, \Upsilon^g \circ \mathcal{Z}(\llbracket x \rrbracket) \rangle = \Xi(\mathbf{f})(\Phi^{\gamma(g)}(\llbracket x \rrbracket)) = \langle \mathbf{f}, \mathcal{Z} \circ \Phi^{\gamma(g)}(\llbracket x \rrbracket) \rangle.$$

Thus, $\Upsilon^g \circ \mathcal{Z} = \mathcal{Z} \circ \Phi^{\gamma(g)}$ as desired. $\square$

Recall that R is the linear map $\mathbb{R}^2 \rightarrow \mathbb{R}^2$ which rotates by $\frac{\pi}{2}$. Also, r is the induced permutation on the signs of quadrants in $\mathbb{R}^2$. See equations 10.2 and 10.4.

Proposition 11.2. *Let σ be a saddle connection and $s \in SP$. Then, $\text{hol } \sigma \in \text{cl}(\mathcal{Q}_s)$ if and only if $\mathcal{Z}(\llbracket \sigma \rrbracket) \in \hat{\mathcal{Q}}_{r(s)}$.*

Proof. Choose any s for which $\text{hol } \sigma \in \text{cl}(\mathcal{Q}_s)$. Write $s = (s_x, s_y)$ with $s_x, s_y \in \{1, -1\}$. Then $r(s) = (-s_y, s_x)$. Because $\text{hol } \sigma \in \text{cl}(\mathcal{Q}_s)$, we have $\text{sgn}(\pi_x(\text{hol } \sigma)) \in \{s_x, 0\}$ and $\text{sgn}(\pi_y(\text{hol } \sigma)) \in \{s_y, 0\}$. Let $\mathbf{a} \in \mathcal{A}$. Then

$$\text{sgn } \mathcal{Z}(\llbracket \sigma \rrbracket)(\mathbf{a}) = \text{sgn } i(\llbracket \sigma \rrbracket, \llbracket \text{cyl}_{\mathbf{a}} \rrbracket) = \text{sgn } (\text{hol } \sigma) \wedge (1, 0) = \text{sgn}(-\pi_y(\text{hol } \sigma)) \in \{-s_y, 0\}.$$

Similarly if $\mathbf{b} \in \mathcal{B}$, then

$$\text{sgn } \mathcal{Z}(\llbracket \sigma \rrbracket)(\mathbf{b}) = \text{sgn } i(\llbracket \sigma \rrbracket, \llbracket \text{cyl}_{\mathbf{b}} \rrbracket) = \text{sgn } (\text{hol } \sigma) \wedge (0, 1) = \text{sgn}(\pi_x(\text{hol } \sigma)) \in \{s_x, 0\}.$$

So, $\mathcal{Z}(\llbracket \sigma \rrbracket) \in \hat{\mathcal{Q}}_{r(s)}$. The converse follows by reversing this argument. $\square$

11.2. Quadrant tracking via expansion. Recall that conjugation by R preserves the subgroup $\rho_\lambda^G \subset SL(2, \mathbb{R})$. Equation 10.3 explains exactly how conjugation by R acts on G . This action on G also relates to the expanding sign action Σ^G . Namely,

$$(11.3) \quad \Sigma^g \circ r = r \circ \Sigma^{\gamma(g)}.$$

We view the following as a corollary of Proposition 10.2.

Corollary 11.3. *Let $\langle g_i \rangle$ be a geodesic ray and $s \in SP$. Assume that σ is a saddle connection for which $\text{hol } \sigma \in \text{cl}(\text{Exp}_\lambda(\gamma(g_1)) \cap \mathcal{Q}_s)$. Then, $\Upsilon^{g_i} \circ \mathcal{Z}(\llbracket \sigma \rrbracket) \in \hat{\mathcal{Q}}_{r(s_i)}$ where $s_i = \Sigma^{g_i}(s)$.*

Proof. By Proposition 10.2, we know $\rho_\lambda^{\gamma(g_i)}(\text{hol } \sigma) \in \text{cl}(\mathcal{Q}_{s_i})$ for all i . We also know that $\text{hol} \circ \Phi^{\gamma(g_i)}(\llbracket \sigma \rrbracket) = \rho_\lambda^{\gamma(g_i)}(\text{hol } \sigma)$ is the holonomy of a saddle connection. By Proposition 11.2, $\mathcal{Z} \circ \Phi^{\gamma(g_i)}(\llbracket \sigma \rrbracket) \in \hat{\mathcal{Q}}_{r(s_i)}$. By Corollary 11.1, we see $\Upsilon^{g_i} \circ \mathcal{Z}(\llbracket \sigma \rrbracket) = \mathcal{Z} \circ \Phi^{\gamma(g_i)}(\llbracket \sigma \rrbracket) \in \hat{\mathcal{Q}}_{r(s_i)}$. $\square$

Proposition 11.4 (Quadrant tracking). *Suppose $\langle g_i \rangle$ is a geodesic ray and $\rho_\lambda^{g_1}(\mathcal{Q}_s) \subset \mathcal{Q}_s$. Then $\Upsilon^{g_i}(\hat{\mathcal{Q}}_s) \subset \hat{\mathcal{Q}}_{s_i}$ where $s_i = \Sigma^{g_i}(s)$.*

Proof. Let $\mathbf{f} \in \hat{\mathcal{Q}}_s$. Given $\mathbf{v} \in \mathcal{V}$, let $\mathbf{e}_\mathbf{v} \in \mathbb{R}^\vee$ denote the function

$$(11.4) \quad \mathbf{e}_\mathbf{v}(\mathbf{w}) = \begin{cases} 1 & \text{if } \mathbf{v} = \mathbf{w} \\ 0 & \text{if } \mathbf{v} \neq \mathbf{w}. \end{cases}$$

Formally, we may write

$$\mathbf{f} = \sum_{v \in \mathcal{V}, \mathbf{f}(v) \neq 0} \mathbf{f}(v) \mathbf{e}_v.$$

This sum makes sense, because for any $v \in \mathcal{V}$, there are only finitely many terms whose support includes v . For each $v \in \mathcal{V}$ with $\mathbf{f}(v) \neq 0$ we can choose a horizontal or vertical saddle connection σ_v such that for any $w \in \mathcal{V}$ we have

$$i(\llbracket \sigma_v \rrbracket, \llbracket cyl_w \rrbracket) = \begin{cases} \text{sgn } \mathbf{f}(v) & \text{if } v = w \\ 0 & \text{otherwise.} \end{cases}$$

Then by definition of $\mathcal{Z}$ we have

$$(11.5) \quad \mathbf{f} = \sum_{v \in \mathcal{V}, \mathbf{f}(v) \neq 0} |\mathbf{f}(v)| \mathcal{Z}(\llbracket \sigma_v \rrbracket).$$

Note that each $\mathcal{Z}(\llbracket \sigma_v \rrbracket) \in \hat{\mathcal{Q}}_s$ and therefore by Proposition 11.2 we know $hol \sigma_v \in cl(\mathcal{Q}_{r^{-1}(s)})$. We now apply Υ^{g_i} to equation 11.5, yielding

$$(11.6) \quad \Upsilon^{g_i}(\mathbf{f}) = \sum_{v \in \mathcal{V}, \mathbf{f}(v) \neq 0} |\mathbf{f}(v)| \Upsilon^{g_i} \circ \mathcal{Z}(\llbracket \sigma_v \rrbracket) = \sum_{v \in \mathcal{V}, \mathbf{f}(v) \neq 0} |\mathbf{f}(v)| \mathcal{Z} \circ \Phi^{\gamma(g_i)}(\llbracket \sigma_v \rrbracket),$$

with the last equality following from Corollary 11.1. Now consider that if $\rho_\lambda^{g_1}(\mathcal{Q}_s) \subset \mathcal{Q}_s$, then $\rho_\lambda^{\gamma(g_1)}(cl(\mathcal{Q}_{r^{-1}(s)})) \subset cl(\mathcal{Q}_{r^{-1}(s)})$. (This follows by inspecting the pairs (g_1, s) allowed by Corollary 10.5.) By Corollary 10.5, we may conclude that

$$hol \Phi^{\gamma(g_i)}(\llbracket \sigma_v \rrbracket) \in cl(\mathcal{Q}_{s'_i}),$$

where $s'_i = \Sigma^{\gamma(g_i)} \circ r^{-1}(s)$. By equation 11.3, we have $s'_i = r^{-1} \circ \Sigma^{g_i}(s)$. Then it follows from Proposition 11.2 that

$$\mathcal{Z} \circ \Phi^{\gamma(g_i)}(\llbracket \sigma_v \rrbracket) \in \hat{\mathcal{Q}}_{r(s'_i)} = \hat{\mathcal{Q}}_{s_i},$$

where $s_i = r(s'_i) = \Sigma^{g_i}(s)$ as in the statement of the proposition. As each of the terms in the sum for $\Upsilon^{g_i}(\mathbf{f})$ in equation 11.6 lie in $\mathcal{Q}_{s_i}$, we know $\Upsilon^{g_i}(\mathbf{f}) \in \mathcal{Q}_{s_i}$. $\square$

Let $\langle g_i \rangle$ be the shrinking sequence for a λ -renormalizable direction $\boldsymbol{\theta} \in S^1$, and $\langle s_i \rangle$ denote the sign sequence. Recall definition 10.7 that n is a critical time if $s_{n-1} = s_n$. We have the following corollary to Proposition 11.4.

Corollary 11.5 (Critical times). *Suppose n is a critical time. Then if $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a $(\boldsymbol{\theta}, n)$ -survivor, then it is a $(\boldsymbol{\theta}, k)$ -survivor for all $k < n$.*

Proof. Let $\mathbf{f}_k = \Upsilon^{g_k}(\mathbf{f})$ and $\boldsymbol{\theta}_k = \rho_\lambda^{g_k}(\boldsymbol{\theta})$. We know that $\boldsymbol{\theta}_n \in \mathcal{Q}_{s_n}$ and $\mathbf{f}_n \in \hat{\mathcal{Q}}_{s_n}$. By proposition 10.8, we know $\rho_\lambda^{g_{n-1}g_n^{-1}}(\mathcal{Q}_{s_n}) \subset \mathcal{Q}_{s_n} = \mathcal{Q}_{s_{n-1}}$. Consider the geodesic segment $\langle h_i = g_{n-i}g_n^{-1} \rangle$. By definition of the sign sequence $\langle s_i \rangle$, we have $\boldsymbol{\theta}_{n-i} \in \mathcal{Q}_{s_{n-i}}$. By proposition 10.2, we know $s_{n-i} = \Sigma^{h_i}(s_n)$. Finally, by proposition 11.4, we know $\mathbf{f}_{n-i} = \Upsilon^{h_i}(\hat{\mathcal{Q}}_s) \subset \hat{\mathcal{Q}}_{s_{n-i}}$ too. Therefore, by definition, $\mathbf{f}_{n-i}$ is a $(\boldsymbol{\theta}, n-i)$ -survivor. $\square$

Corollary 11.6. *Suppose $\boldsymbol{\theta}$ is a λ -renormalizable direction. If for some $N \geq 0$ we know that $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ is a $(\boldsymbol{\theta}, k)$ -survivor for all $k \geq N$, then $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor.*

Proof. Corollary 10.9 guarantees that we have infinitely many critical times n . Choose such an $n > N$. Corollary 11.5 implies that $\mathbf{f}$ is a $(\boldsymbol{\theta}, k)$ -survivor for all $k < n$. $\square$

Corollary 11.7 (Group invariance). *Assume that $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor, for $\boldsymbol{\theta}$ a λ -renormalizable direction. Then for all $g \in G$, $\Upsilon^g(\mathbf{f})$ is a $\rho_\lambda^g(\boldsymbol{\theta})$ -survivor.*

Proof. Let $\mathbf{f}' = \Upsilon^g(\mathbf{f})$. Let $\langle g_i \rangle$ be the λ -shrinking sequence for $\boldsymbol{\theta}$, and $\langle g'_i \rangle$ be the λ -shrinking sequence for $\boldsymbol{\theta}' = \pi_{\mathbb{S}^1} \circ \rho_\lambda^g(\boldsymbol{\theta})$. Then the λ -shrinking sequences of $\boldsymbol{\theta}$ and $\boldsymbol{\theta}'$ are tail equivalent, as defined in section 3. Moreover, regardless of the choice of $g \in G$ there are positive integers m and n such that

$$\Upsilon^{g^m}(\mathbf{f}) = \Upsilon^{g'_n}(\mathbf{f}') \quad \text{and} \quad \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g^m}(\boldsymbol{\theta}) = \pi_{\mathbb{S}^1} \circ \rho_\lambda^{g'_n}(\boldsymbol{\theta}').$$

Then this is also true for m replaced by $m + k$ and n replaced by $n + k$ for all $k \geq 0$. Thus, corollary 11.6 implies that $\mathbf{f}'$ is a $\boldsymbol{\theta}'$ -survivor. $\square$

11.3. Survivors in $H^1(S, V, \mathbb{R})$ and $\mathbb{R}^\mathcal{V}$. In this section we will provide a proof of Theorem 6.28. The main idea of the proof is to understand the function $\mathbf{F} : H^1(S, V, \mathbb{R}) \rightarrow \mathbb{R}^\mathcal{V}$ given by

$$\mathbf{F}(m)(v) = m(\llbracket cyl_v \rrbracket),$$

where $\llbracket cyl_v \rrbracket$ represents the homology class of the cylinder associated to v oriented rightward or upward. Recall that $r : SP \rightarrow SP$ is the action on signs of quadrants induced by a rotation by $\frac{\pi}{2}$ radians. See equation 10.4. We have the following.

Proposition 11.8. $\mathbf{F} \circ \Phi_*^g = \Upsilon^{\gamma(g)} \circ \mathbf{F}$.

Proof. It is enough to handle the cases of $g \in \{h^k, v^k : k \in \mathbb{Z}\}$. Let $g = h^k$. Suppose $a \in \mathcal{A}$. Then by Proposition 6.7,

$$(\mathbf{F} \circ \Phi_*^{h^k}(m))(a) = m(\Phi^{h^{-k}}(\llbracket cyl_a \rrbracket)) = m(\llbracket cyl_a \rrbracket).$$

By definition of γ , we have $\gamma(h^k) = v^{-k}$. By equation 6.3,

$$(\Upsilon^{v^{-k}} \circ \mathbf{F}(m))(a) = \mathbf{F}(m)(a) = m(\llbracket cyl_a \rrbracket).$$

Now let $b \in \mathcal{B}$. We have

$$(\mathbf{F} \circ \Phi_*^{h^k}(m))(b) = m(\Phi^{h^{-k}}(\llbracket cyl_b \rrbracket)) = m(\llbracket cyl_b \rrbracket - k \sum_{a \sim b} \llbracket cyl_a \rrbracket).$$

$$(\Upsilon^{v^{-k}} \circ \mathbf{F}(m))(b) = \mathbf{F}(m)(b) - k \sum_{a \sim b} \mathbf{F}(m)(a) = m(\llbracket cyl_b \rrbracket) - k \sum_{a \sim b} m(\llbracket cyl_a \rrbracket).$$

These two expressions are equal by linearity of m . The proof is similar for the case of $g = v^k$. $\square$

Recall that R acts on $\mathbb{R}^2$ by rotation by $\frac{\pi}{2}$. See equation 10.2.

Proposition 11.9. *If $m \in H^1(S, V, \mathbb{R})$ is a $(\boldsymbol{\theta}, n)$ -survivor then $\mathbf{F}(m) \in \mathbb{R}^\mathcal{V}$ is a $(R^{-1}(\boldsymbol{\theta}), n)$ -survivor.*

Proof. To ease notation define $\boldsymbol{\theta}' = R^{-1}(\boldsymbol{\theta})$. Let $\langle g_n \rangle$ denote the shrinking sequence for $\boldsymbol{\theta}$. Let $g'_n = \gamma(g_n)$. By equation 10.3, $\langle g'_n \rangle$ is the shrinking sequence for $\boldsymbol{\theta}'$. Therefore, if $\langle s_n \rangle$ is the sign sequence of $\boldsymbol{\theta}$, then the sign sequence for $\boldsymbol{\theta}'$ is $\langle s'_n = r^{-1}(s_n) \rangle$. By proposition 6.20, we must show that $\Upsilon^{g'_n} \circ \mathbf{F}(m) \in \hat{\mathcal{Q}}_{s'_n}$. By proposition 11.8, $\Upsilon^{g'_n} \circ \mathbf{F}(m) = \mathbf{F}(\Phi_*^{g_n}(m))$. For $v \in \mathcal{V}$, we have

$$\mathbf{F}(\Phi_*^{g_n}(m))(v) = (\Phi_*^{g_n}(m))(\llbracket cyl_v \rrbracket).$$

Define $x, y \in \mathbb{R}$ so that $(x, y) = \boldsymbol{\theta}$. We may apply Proposition 6.10 because m is a $(\boldsymbol{\theta}, n)$ -survivor and because $\llbracket cyl_v \rrbracket$ can be written as a sum of homology classes of saddle connections parallel to $hol(\llbracket cyl_v \rrbracket)$. Therefore, for all $a \in \mathcal{A}$, unless $\mathbf{F}(\Phi_*^{g_n}(m))(a) = 0$,

$$\text{sgn } \mathbf{F}(\Phi_*^{g_n}(m))(a) = \text{sgn } hol(\llbracket cyl_a \rrbracket) \wedge \rho_\lambda^{g_n}(\boldsymbol{\theta}) = \text{sgn } (1, 0) \wedge \rho_\lambda^{g_n}(\boldsymbol{\theta}) = \text{sgn } y.$$

Similarly, for $b \in \mathcal{B}$, unless $\mathbf{F}(\Phi_*^{g_n}(m))(b) = 0$, we have

$$\text{sgn } \mathbf{F}(\Phi_*^{g_n}(m))(b) = \text{sgn } hol(\llbracket cyl_b \rrbracket) \wedge \rho_\lambda^{g_n}(\boldsymbol{\theta}) = \text{sgn } (0, 1) \wedge \rho_\lambda^{g_n}(\boldsymbol{\theta}) = -\text{sgn } x.$$

Noting that $(x, y) \in \mathcal{Q}_{s_n}$, we see $(y, -x) \in \mathcal{Q}_{r^{-1}(s_n)}$. Thus, $\mathbf{F}(\Phi_*^{g_n}(m))(v) \in \hat{\mathcal{Q}}_{s'_n}$ as desired. $\square$

Proof of Theorem 6.28. Suppose that $m \in H^1(S, V, \mathbb{R})$ is a $\boldsymbol{\theta}$ -survivor, and suppose that there is no $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ for which $m = \Xi(\mathbf{f})$. Then there are horizontal or vertical saddle connections $\sigma, \sigma' \in E$ both of which cross cyl_v for some $v \in V$ and for which $m(\llbracket \sigma \rrbracket) \neq m(\llbracket \sigma' \rrbracket)$. Note that for such a pair of saddle connections, we have

$$\Phi^g(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket) = \llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket$$

for all $g \in G$, because the difference $\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket$ has zero algebraic intersection number with each horizontal and vertical cylinder. Consider the shrinking sequence $\langle g_n \rangle$ of $\boldsymbol{\theta}$. We have

$$(11.7) \quad (\Phi_*^{g_n}(m))(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket) = m(\Phi^{g_n^{-1}}(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket)) = m(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket).$$

So, this quantity remains constant.

On the other hand, let $\mathbf{g} = \mathbf{F}(m)$. By Proposition 11.9, $\mathbf{g}$ is a $\boldsymbol{\theta}' = R(\boldsymbol{\theta})$ -survivor. The λ -shrinking sequence of $\boldsymbol{\theta}'$ is $\langle g'_n = \gamma(g_n) \rangle$ by equation 10.3. Now we utilize the subsequence decay property. By this property, there is a subsequence $\langle g'_{n_i} \rangle$ such that for all $v \in \mathcal{V}$ we have

$$\lim_{i \rightarrow \infty} \Upsilon^{g'_{n_i}}(\mathbf{g})(v) = 0.$$

The quantity $\Upsilon(g'_{n_i})(\mathbf{g})(v)$ has meaning to us because

$$\Upsilon^{g'_{n_i}}(\mathbf{g})(v) = (\Upsilon^{g'_{n_i}} \circ \mathbf{F}(m))(v) = (\mathbf{F} \circ \Phi_*^{g'_{n_i}}(m))(v) = \Phi_*^{g'_{n_i}}(m)(\llbracket cyl_v \rrbracket).$$

Because m is a $(\boldsymbol{\theta}, n)$ -survivor for all n , we know that for any saddle connection $\nu \in E$ satisfies either $\Phi_*^{g_n}(m)(\llbracket \nu \rrbracket) = 0$ or

$$(11.8) \quad \text{sgn } \Phi_*^{g_n}(m)(\llbracket \nu \rrbracket) = \text{sgn } (hol(\llbracket \nu \rrbracket) \wedge \rho_\lambda^{g_n}(\boldsymbol{\theta})).$$

In particular, choose and orient a boundary component of cyl_v for some $v \in \mathcal{V}$. This component is made up of some number of oriented horizontal or vertical saddle connections $\nu_1, \dots, \nu_k$. Equation 11.8 implies that all nonzero $\Phi_*^{g_n}(m)(\llbracket \nu_j \rrbracket)$ have the same sign regardless of the choice of $j = 1, \dots, k$. Since $\llbracket \nu_1 \rrbracket + \dots + \llbracket \nu_k \rrbracket = \llbracket cyl_v \rrbracket$ in $H_1(S, V, \mathbb{Z})$, it follows that

$$\lim_{i \rightarrow \infty} \Phi_*^{g'_{n_i}}(m)(\llbracket \nu_j \rrbracket)(\mathbf{g})(v) = 0 \quad \text{for all } j = 1, \dots, k.$$

Now note that every horizontal and every vertical saddle connection lies in the boundary of some cylinder cyl_v for $v \in \mathcal{V}$. In particular, this holds for σ and σ' . We conclude that

$$\lim_{i \rightarrow \infty} \Phi_*^{g'_{n_i}}(m)(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket) = 0.$$

This contradicts our assumption that $m(\llbracket \sigma \rrbracket - \llbracket \sigma' \rrbracket) \neq 0$ via equation 11.7. $\square$

11.4. Survivors and the pairing of $\mathbb{R}^\nu$ with $\mathbb{R}_c^\nu$. We begin this section by summarizing results from the previous two sections. Then we will state some consequences of our previous work.

Let θ be a λ -renormalizable direction. Let $\langle g_i \rangle$ and $\langle s_i \rangle$ be the shrinking and sign sequences of θ . Recall $\mathcal{S}_\theta$ denotes the set of θ -survivors in $\mathbb{R}^\nu$. By definition 6.13 this is

$$\mathcal{S}_\theta = \{\mathbf{f} \in \mathbb{R}^\nu : \Upsilon^{g_n}(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_n} \text{ for all } n \geq 0\}.$$

Alternately, by Theorems 6.9 and 6.28, $\Xi(\mathcal{S}_\theta)$ is the set of all cohomology classes arising from transverse invariant measures to the foliation in the direction θ . Then Lemma 6.4 implies that

$$(11.9) \quad \mathcal{S}_\theta = \{\mathbf{f} \in \mathbb{R}^\nu : \begin{array}{l} \text{sgn}\langle \mathbf{f}, \mathcal{Z}(\llbracket \sigma \rrbracket) \rangle = \text{sgn}(\text{hol } \sigma \wedge \theta) \text{ for all saddle} \\ \text{connections } \sigma \text{ with } \langle \mathbf{f}, \mathcal{Z}(\llbracket \sigma \rrbracket) \rangle \neq 0 \end{array} \}.$$

Here, the quantity $\langle \mathbf{f}, \mathcal{Z}(\llbracket \sigma \rrbracket) \rangle$ represents the value assigned to $\llbracket \sigma \rrbracket$ by the associated cohomology class (which is potentially induced by an invariant measure). See equation 11.2.

The later sections of this paper are primarily interested in questions of the following sort. Given $\mathbf{x} \in \mathbb{R}_c^\nu$ with some properties and any $\mathbf{f} \in \mathcal{S}_\theta$, is it true that $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$? We pursue this sort of question from the following point of view. One way to show that $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$ is to find saddle connections σ_j so that $\mathbf{x} = \sum_j \mathcal{Z}(\llbracket \sigma_j \rrbracket)$ and $\text{hol } \sigma_j \wedge \theta > 0$. If we can find this, then $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$ follows from equation 11.9.

There is a second, more operator theoretic point of view as well.

Proposition 11.10 (Positivity checks). *Let θ be a λ -renormalizable direction. Let $\langle g_i \rangle$ and $\langle s_i \rangle$ be the shrinking and sign sequences of θ . Let $\mathbf{x} \in \mathbb{R}_c^\nu$. Either of the following two statements imply that for all $\mathbf{f} \in \mathcal{S}_\theta$, we have $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$.*

- (1) *There exist a finite collection oriented saddle connections $\{\sigma_j\}$ so that $\text{hol } \sigma_j \wedge \theta > 0$ for each j and $\mathbf{x} = \sum_j \mathcal{Z}(\llbracket \sigma_j \rrbracket)$.*
- (2) *There is a critical time t so that $\Upsilon^{\gamma(g_t)}(\mathbf{x}) \in \hat{\mathcal{Q}}_{s_t}$.*

Remark 11.11. In fact, it can be observed that these two statements are equivalent. That (1) implies (2) follows essentially from Theorem 10.14. Conversely, if we have (2), then $\Upsilon^{\gamma(g_t)}(\mathbf{x})$ can be expressed as $\mathcal{Z}(\llbracket \tau_1 \rrbracket + \dots + \llbracket \tau_N \rrbracket)$ where each τ_j is an appropriately oriented horizontal or vertical saddle connection, so that $\text{hol } \tau_j \in \text{cl}(\mathcal{Q}_{r^{-1}(s_t)})$. Then

$$\mathbf{x} = \Upsilon^{\gamma(g_t)^{-1}} \circ \mathcal{Z}(\llbracket \tau_1 \rrbracket + \dots + \llbracket \tau_N \rrbracket) = \mathcal{Z} \circ \Phi^{g_t^{-1}}(\llbracket \tau_1 \rrbracket + \dots + \llbracket \tau_N \rrbracket)$$

and we may take $\sigma_j = \Phi^{g_t^{-1}}(\tau_j)$.

Proof of Proposition 11.10. For statement (1), this follows from the paragraph preceding the proposition. In the case of statement (2), we have $\langle \mathbf{f}, \mathbf{x} \rangle = \langle \Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{x}) \rangle$. Since both $\Upsilon^{g_t}(\mathbf{f})$ and $\Upsilon^{\gamma(g_t)}(\mathbf{x})$ lie in $\hat{\mathcal{Q}}_{s_t}$, their product is non-negative. $\square$

The following strengthens the utility of statement (2) of Proposition 11.10.

Proposition 11.12. *Let $\theta \in \mathcal{R}_\lambda$ and let $\mathbf{f}$ be a θ -survivor. Let $\langle g_i \rangle$ and $\langle s_i \rangle$ be the shrinking and sign sequences of θ . If $\mathbf{x} \in \mathbb{R}_c^\nu \cap \hat{\mathcal{Q}}_{s_0}$, then for any critical time t , $\Upsilon^{\gamma(g_t)}(\mathbf{x}) \in \hat{\mathcal{Q}}_{s_t}$.*

Proof. We can write $\mathbf{x} = \sum_{j=1}^N \mathcal{Z}(\llbracket \sigma_j \rrbracket)$ where each σ_j is a horizontal or vertical saddle connection oriented so that $\text{hol } \sigma_j \in \text{cl}(\mathcal{Q}_{r^{-1}(s_0)})$. See Proposition 11.2. Then,

$$(11.10) \quad \Upsilon^{\gamma(g_t)}(\mathbf{x}) = \sum_{j=1}^N \mathcal{Z} \circ \Phi^{g_t}(\llbracket \sigma_j \rrbracket),$$

by Corollary 11.1. Looking at the action on holonomy, we have

$$\text{hol} \circ \Phi^{g_t}(\llbracket \sigma_j \rrbracket) = \rho_\lambda^{g_t}(\text{hol } \sigma_j).$$

By Proposition 7.3, since $\rho_\lambda^{g_1}$ shrinks the vector $\theta \in \mathcal{Q}_{s_0}$, we know $\rho_\lambda^{g_1}$ expands every vector in $\mathcal{Q}_{r^{-1}(s_0)}$. Therefore, the quadrant containing $\rho_\lambda^{g_t}(\text{hol } \sigma_j)$ is governed by the expanding sign action. Namely by Proposition 10.2, we must have $\rho_\lambda^{g_i}(\text{hol } \sigma_j) \in \text{cl}(\mathcal{Q}_{s'_i})$ where $s'_i = \Sigma^{g_i} \circ r^{-1}(s_0)$ for all i . Therefore, by Proposition 11.2 we know each term from equation 11.10 satisfies

$$\mathcal{Z} \circ \Phi^{g_t}(\llbracket \sigma_j \rrbracket) \in \hat{\mathcal{Q}}_{r(s'_t)}.$$

By equation 11.3, we know $s'_i = r^{-1} \circ \Sigma^{\gamma(g_i)}(s_0)$, and so $r(s'_t) = \Sigma^{\gamma(g_t)}(s_0)$. Therefore $r(s'_t) = s_t$ by Corollary 10.11. It follows from equation 11.10 that $\Upsilon^{\gamma(g_t)}(\mathbf{x}) \in \hat{\mathcal{Q}}_{s_t}$ as desired. $\square$

12. SUBGRAPHS AND COVERS

In this section, we will consider how the group of operators Υ^G is affected by restricting to subgraphs and lifting to finite covers. This will be relevant to proofs in section 14, where we prove the adjacency sign property and the decay properties hold when $\mathcal{G}$ has no vertices of valance one.

12.1. Subgraphs. Throughout this subsection, $\mathcal{H} \subset \mathcal{G}$ will be a connected infinite subgraph. We will use subscripts to distinguish quantities related to the two graphs. For instance, we write $\mathcal{V}_{\mathcal{H}} \subset \mathcal{V}_{\mathcal{G}}$ to indicate that the vertex set of $\mathcal{H}$ is a subset of the vertex set of $\mathcal{G}$. The subgraph $\mathcal{H}$ inherits a bipartite structure from $\mathcal{G}$. Namely, $\mathcal{A}_{\mathcal{H}} = \mathcal{A}_{\mathcal{G}} \cap \mathcal{V}_{\mathcal{H}}$ and $\mathcal{B}_{\mathcal{H}} = \mathcal{B}_{\mathcal{G}} \cap \mathcal{V}_{\mathcal{H}}$. We use $\Upsilon_{\mathcal{G}}^G : \mathbb{R}^{\mathcal{V}_{\mathcal{G}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ and $\Upsilon_{\mathcal{H}}^G : \mathbb{R}^{\mathcal{V}_{\mathcal{H}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$ to denote the appropriate groups of operators on the two spaces. We abuse notation by identifying $\mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$ with the subset of $\mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ whose support is contained in $\mathcal{V}_{\mathcal{H}}$.

Proposition 12.1. *Let $\mathbf{f} \in \mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$. Suppose that $\mathbf{f} \in \hat{\mathcal{Q}}_s$ for some $s \in SP$. Write $s = (a, b)$ with $a, b \in \{\pm 1\}$. Then for all integers $k \neq 0$, the following statements hold.*

- (1) $\mathbf{H}_{\mathcal{G}}^k(\mathbf{f}) - \mathbf{H}_{\mathcal{H}}^k(\mathbf{f})$ is supported on a subset of $\mathcal{A}_{\mathcal{G}}$ and the sign of all non-zero values of this function is $b \cdot \text{sgn}(k)$.
- (2) $\mathbf{V}_{\mathcal{G}}^k(\mathbf{f}) - \mathbf{V}_{\mathcal{H}}^k(\mathbf{f})$ is supported on a subset of $\mathcal{B}_{\mathcal{G}}$ and the sign of all non-zero values of this function is $a \cdot \text{sgn}(k)$.

Proof. We will prove statement (1). Statement (2) has a similar proof. By considering the definition of $\mathbf{H}$ given in equation 6.2, for all $\mathbf{v} \in \mathcal{V}_{\mathcal{G}}$ we have

$$(\mathbf{H}_{\mathcal{G}}^k(\mathbf{f}) - \mathbf{H}_{\mathcal{H}}^k(\mathbf{f}))(\mathbf{v}) = \begin{cases} \sum_{b \sim_{\mathcal{G}} \mathbf{v}, b \not\sim_{\mathcal{H}} \mathbf{v}} k\mathbf{f}(b) & \text{if } \mathbf{v} \in \mathcal{A}_{\mathcal{G}} \\ 0 & \text{if } \mathbf{v} \in \mathcal{B}_{\mathcal{G}}, \end{cases}$$

where the sum should be interpreted as over all edges leaving $\mathbf{v}$ that appear in $\mathcal{G}$ but not in $\mathcal{H}$. The conclusion follows. $\square$

Corollary 12.2. *Let $\langle g_i \rangle$ be a geodesic ray in G . Suppose $\mathbf{x} \in \mathbb{R}^{\mathcal{V}_{\mathcal{H}}} \cap \hat{\mathcal{Q}}_{s_0}$ for some $s_0 \in SP$, and define $s_i = \Sigma^{g_i}(s_0)$. Then for all $i \geq 1$,*

$$\Upsilon_{\mathcal{G}}^{g_i g_1^{-1}}(\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x})) \in \hat{\mathcal{Q}}_{s_i}.$$

Proof. By Proposition 12.1 and comparison to Definition 10.1 of the expanding sign action, we see that $\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x}) \in \hat{\mathcal{Q}}_{s_1}$. Moreover, because of the description of the support as a subset of either $\mathcal{A}$ or $\mathcal{B}$, we have

$$\Upsilon_{\mathcal{G}}^{g_1}(\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x})) = \Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x}).$$

Since $s_1 = \Sigma^{g_1}(s_0)$, we know $s_1 = \Sigma^{g_1}(s_1)$. Therefore we have $\rho_{\lambda}^{g_1}(\mathcal{Q}_{s_1}) \subset \mathcal{Q}_{s_1}$ and thus

$$\Upsilon_{\mathcal{G}}^{g_i g_1^{-1}}(\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x})) = \Upsilon_{\mathcal{G}}^{g_i}(\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x})) \in \hat{\mathcal{Q}}_{s'_i}$$

where $s'_i = \Sigma^{g_i}(s_1)$ by Proposition 11.4. Finally, note that

$$s_i = \Sigma^{g_i}(s_0) = \Sigma^{g_i g_1^{-1}}(s_1) = \Sigma^{g_i}(s_1) = s'_i$$

since $\Sigma^{g_1}(s_1) = s_1$. Thus, $\Upsilon_{\mathcal{G}}^{g_i g_1^{-1}}(\Upsilon_{\mathcal{G}}^{g_1}(\mathbf{x}) - \Upsilon_{\mathcal{H}}^{g_1}(\mathbf{x})) \in \hat{\mathcal{Q}}_{s_i}$ as desired. $\square$

For $\mathbf{v} \in \mathcal{V}_{\mathcal{H}}$, we use $\mathbf{e}_{\mathbf{v}}$ to denote the element of either $\mathbb{R}_{\mathcal{C}}^{\mathcal{V}_{\mathcal{G}}}$ or $\mathbb{R}_{\mathcal{C}}^{\mathcal{V}_{\mathcal{H}}}$ (depending on context) which assigns one to $\mathbf{v}$ and zero to everything else.

Theorem 12.3. *Suppose $\mathcal{H} \subset \mathcal{G}$ is an infinite connected subgraph. Let $\langle g_i \rangle$ be any geodesic ray in G . Then, for any vertex $\mathbf{v} \in \mathcal{V}_{\mathcal{H}}$,*

$$|\Upsilon_{\mathcal{H}}^{g_i}(\mathbf{e}_{\mathbf{v}})(\mathbf{w})| \leq |\Upsilon_{\mathcal{G}}^{g_i}(\mathbf{e}_{\mathbf{v}})(\mathbf{w})|$$

for all $\mathbf{w} \in \mathcal{V}_{\mathcal{H}}$ and all $i \geq 0$.

Proof. Give $\mathcal{H}$ an arbitrary ribbon graph structure, and find an arbitrary eigenvector of the adjacency operator. Then Corollary 11.3 gives a sequence $s_i \in SP$ such that $\Upsilon_{\mathcal{H}}^{g_i}(\mathbf{e}_{\mathbf{v}}) \in \hat{\mathcal{Q}}_{s_i}$ and $\Upsilon_{\mathcal{G}}^{g_i}(\mathbf{e}_{\mathbf{v}}) \in \hat{\mathcal{Q}}_{s_i}$. In particular, $s_i = \Sigma^{g_i}(s_0)$ for the appropriate choice of $s_0 \in SP$.

Now we will inductively define elements $\mathbf{y}_i \in \mathbb{R}_{\mathcal{C}}^{\mathcal{V}_{\mathcal{G}}}$ for each integer $i \geq 1$. Assuming $\mathbf{y}_1, \dots, \mathbf{y}_{i-1}$ are defined, we define $\mathbf{y}_i$ to be the unique element of so that the following equation holds

$$\Upsilon_{\mathcal{G}}^{g_i}(\mathbf{e}_{\mathbf{v}}) - \Upsilon_{\mathcal{H}}^{g_i}(\mathbf{e}_{\mathbf{v}}) = \Upsilon_{\mathcal{G}}^{g_i g_1^{-1}}(\mathbf{y}_1) + \Upsilon_{\mathcal{G}}^{g_i g_2^{-1}}(\mathbf{y}_2) + \dots + \mathbf{y}_i.$$

We will show that

$$(12.1) \quad \Upsilon_{\mathcal{G}}^{g_i g_j^{-1}}(\mathbf{y}_j) \in \hat{\mathcal{Q}}_{s_i} \quad \text{for all } i \geq j.$$

In particular, for all $\mathbf{w} \in \mathcal{V}_{\mathcal{G}}$ we have

$$\Upsilon_{\mathcal{G}}^{g_i}(\mathbf{e}_{\mathbf{v}})(\mathbf{w}) = \Upsilon_{\mathcal{H}}^{g_i}(\mathbf{e}_{\mathbf{v}})(\mathbf{w}) + \Upsilon_{\mathcal{G}}^{g_i g_1^{-1}}(\mathbf{y}_1)(\mathbf{w}) + \Upsilon_{\mathcal{G}}^{g_i g_2^{-1}}(\mathbf{y}_2)(\mathbf{w}) + \dots + \mathbf{y}_i(\mathbf{w}),$$

and all terms in this sum are either zero or have the same sign. Therefore, equation 12.1 implies the theorem.

To shorten our formulas, let $\mathbf{x}_j = \Upsilon_{\mathcal{H}}^{g_j}(\mathbf{e}_{\mathbf{v}})$. By induction, we observe that

$$\mathbf{y}_j = \Upsilon_{\mathcal{G}}^{g_j g_{j-1}^{-1}}(\mathbf{x}_{j-1}) - \Upsilon_{\mathcal{H}}^{g_j g_{j-1}^{-1}}(\mathbf{x}_{j-1}).$$

Let $i \geq j$. Recall from the first paragraph that $\mathbf{x}_j \in \hat{\mathcal{Q}}_{s_j}$. Applying Corollary 12.2 to the geodesic ray $\langle g_i g_j^{-1} \rangle_{i \geq j}$ yields that $\Upsilon^{g_i g_j^{-1}}(\mathbf{y}_j) \in \hat{\mathcal{Q}}_{s'_{i,j}}$ where $s'_{i,j} = \Sigma^{g_i g_j^{-1}}(s_j)$. The conclusion follows from the observation that $s_j = \Sigma^{g_j}(s_0)$ and therefore

$$s'_{i,j} = \Sigma^{g_i g_j^{-1}}(s_j) = \Sigma^{g_i g_j^{-1}} \circ \Sigma^{g_j}(s_0) = s_i.$$

This proves equation 12.1 as desired. $\square$

12.2. Finite covers. Now let $\tilde{\mathcal{G}}$ be a finite cover of $\mathcal{G}$. We will let $\pi : \mathcal{V}_{\tilde{\mathcal{G}}} \rightarrow \mathcal{V}_{\mathcal{G}}$ denote the restriction of the covering map to the vertices. This map induces a map $\pi_* : \mathbb{R}^{\mathcal{V}_{\tilde{\mathcal{G}}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ given by

$$(12.2) \quad \pi_*(\tilde{\mathbf{f}})(\mathbf{v}) = \sum_{\tilde{\mathbf{v}} \in \pi^{-1}(\mathbf{v})} \tilde{\mathbf{f}}(\tilde{\mathbf{v}}).$$

Since π arose from a covering map, we have the following.

Proposition 12.4. *For all $g \in G$, $\Upsilon^g \circ \pi_* = \pi_* \circ \Upsilon^g_{\tilde{\mathcal{G}}}$.*

This may be proved by checking that the equation holds for powers of the generators of G .

Finally, to use the previous propositions in our setting, we need to be able to find nice subgraphs of finite covers of $\mathcal{G}$. We use $\mathcal{G}_{\mathbb{Z}}$ to denote the graph with vertex set consisting of the integers which is formed by drawing edges between subsequent integers.

Lemma 12.5. *Suppose $\mathcal{G}$ is an infinite connected graph so that every vertex has finite valance. Further suppose that $\mathcal{G}$ has no vertices of valance one. Let $\mathbf{v}$ be any vertex of $\mathcal{G}$. Then either*

- (1) *there is an embedding $\phi : \mathcal{G}_{\mathbb{Z}} \rightarrow \mathcal{G}$ such that $\phi(0) = \mathbf{v}$, or*
- (2) *there is a double cover $\tilde{\mathcal{G}}$ of $\mathcal{G}$ (depending on $\mathbf{v}$) and an embedding $\tilde{\phi} : \mathcal{G}_{\mathbb{Z}} \rightarrow \tilde{\mathcal{G}}$ such that $\tilde{\phi}(0)$ is a lift of $\mathbf{v}$.*

Proof. For $S \subset \mathbb{Z}$ let $\mathcal{G}_S$ denote the graph with vertex set S which is formed by adding edges between all pairs of integers whose difference is 1. Because $\mathcal{G}$ is infinite and has bounded valance, there are vertices of arbitrary large distance from $\mathbf{v}$. Thus, we can define a metric embedding $\psi_0 : \mathcal{G}_{\{0, -1, -2, \dots\}} \rightarrow \mathcal{V}_{\mathcal{G}}$ such that $\psi_0(0) = \mathbf{v}$. (For each $n \geq 0$, choose $\psi_0(-n)$ so that its distance from $\mathbf{v}$ is n .) Now for $n > 0$ inductively define $\psi_n : \mathcal{G}_{\{n, n-1, \dots\}} \rightarrow \mathcal{V}_{\mathcal{G}}$ so that

- ψ_n restricted to $\mathcal{G}_{\{n-1, n-2, \dots\}}$ is ψ_{n-1} , and
- ψ_n applied to the edge $(n-1, n)$ is distinct from the edge $(n-2, n-1)$. (This can be done because $\mathcal{G}$ has no vertices of valance one.)

Now assume ψ_n is always injective. Then the limit $\lim_{n \rightarrow \infty} \psi_n$ is an embedding of $\mathcal{G}_{\mathbb{Z}} \rightarrow \mathcal{G}$. So assume ψ_n is not injective for some n . Let N be the smallest integer for which ψ_n is not injective. Then, there is an $M < N$ for which $\psi_N(N) = \psi_N(M)$. Let $\mathcal{H}$ be the circular subgraph of $\mathcal{G}$ consisting of the image $\psi_N([M, N])$. Let $[\mathcal{H}] \in H_1(\mathcal{G}; \mathbb{Z})$ denote the corresponding homology class with $\mathbb{Z}$ coefficients. Note that $[\mathcal{H}]$ must be primitive. Let $p : H_1(\mathcal{G}; \mathbb{Z}) \rightarrow \mathbb{Z}_2$ be any group homomorphism so that $p([\mathcal{H}]) = 1$. Let $p' : \pi_1(\mathcal{G}) \rightarrow \mathbb{Z}_2$ be the homomorphism constructed by taking homology class of an element of $\pi_1(\mathcal{G})$ and then applying p . Let $\tilde{\mathcal{G}}$ denote the double cover of $\mathcal{G}$ which corresponds to the kernel of p . Then $\psi_N([M, N])$ lifts to an embedding $\tilde{\psi}_N(M)$ and $\tilde{\psi}_N(N)$ are both lifts of the point

$\psi_N(N) = \psi_N(M)$. Also consider the two disjoint lifts of ψ_M . These two lifts are rays which end at $\tilde{\psi}_N(M)$ and $\tilde{\psi}_N(N)$. The two lifts of the ray ψ_M and one of the lifts of the segment $\psi_N|_{[M,N]}$ can be stitched together to form our desired embedding $\tilde{\phi} : \mathcal{G}_{\mathbb{Z}} \rightarrow \tilde{\mathcal{G}}$. $\square$

13. GENERALIZED FARKAS' THEOREM

We will use a generalization of Farkas' theorem given by Craven and Koliha in theorem 2 of [CK77]. We will introduce some of their terminology and then give their result. Then we will apply this to prove Lemma 6.35. We also prove Lemma 6.41 with techniques developed in this section.

If X is a real vector space, its algebraic dual $X^\sharp$ is the collection of linear functionals $X \rightarrow \mathbb{R}$. $X^\sharp$ is also a real vector space and we have a bilinear pairing $\langle \cdot, \cdot \rangle : X \times X^\sharp \rightarrow \mathbb{R}$ given by $\langle \mathbf{x}, \mathbf{f} \rangle = \mathbf{f}(\mathbf{x})$. A subset $X^+ \subset X^\sharp$ is said to *separate points in X* if for any distinct $\mathbf{x}_1, \mathbf{x}_2 \in X$ there is a $\mathbf{f} \in X^+$ such that $\langle \mathbf{x}_1, \mathbf{f} \rangle \neq \langle \mathbf{x}_2, \mathbf{f} \rangle$. The pair $\langle X, X^+ \rangle$ is a *dual pair* if X^+ is a subspace of $X^\sharp$ which separates points in X . The *weak topology* on X with respect to the dual pair $\langle X, X^+ \rangle$ is the coarsest topology which makes each functional in X^+ continuous.

A convex cone in X is a subset $S \subset X$ such that $S + S \subset S$ and $\alpha S \subset S$ for all $\alpha \geq 0$. If $\langle X, X^+ \rangle$ is a dual pair, then the *anticone* of S is

$$S^+ = \{\mathbf{f} \in X^+ : \langle \mathbf{x}, \mathbf{f} \rangle \geq 0 \text{ for all } \mathbf{x} \in S\}.$$

The *algebraic adjoint* of a linear map $M : X \rightarrow Y$ is the map $M^\sharp : Y^\sharp \rightarrow X^\sharp$ defined by

$$\langle \mathbf{x}, M^\sharp(\mathbf{g}) \rangle = \langle M(\mathbf{x}), \mathbf{g} \rangle.$$

If $\langle X, X^+ \rangle$ and $\langle Y, Y^+ \rangle$ are dual pairs then the linear map $M : X \rightarrow Y$ is weakly-continuous if and only if $M^\sharp(Y^+) \subset X^+$. In this case we define the adjoint $M^+ : Y^+ \rightarrow X^+$ to be the restriction of $M^\sharp$ to Y^+ .

Theorem 13.1 (Generalized Farkas' theorem [CK77]). *Let $\langle X, X^+ \rangle$ and $\langle Y, Y^+ \rangle$ be dual pairs, let $S \subset X$ be a convex cone, and let $M : X \rightarrow Y$ be a weakly-continuous linear map. If $M(S)$ is weakly-closed then the following are equivalent conditions on $\mathbf{b} \in Y$:*

- (1) *The equation $M\mathbf{x} = \mathbf{b}$ has a solution $\mathbf{x} \in S$.*
- (2) *If $\mathbf{y}^+ \in Y^+$ satisfies $M^+(\mathbf{y}^+) \in S^+$ then $\langle \mathbf{b}, \mathbf{y}^+ \rangle \geq 0$.*

Lemma 6.35 is a special case of this theorem. To see this, consider the dual pair $\langle \mathbb{R}^\mathcal{V}, \mathbb{R}_c^\mathcal{V} \rangle$. Here the weak topologies are simply the topologies of pointwise convergence. We observe that the adjoint of the adjacency operator $\mathbf{A} : \mathbb{R}^\mathcal{V} \rightarrow \mathbb{R}^\mathcal{V}$ is just the restriction of $\mathbf{A}$ to $\mathbb{R}_c^\mathcal{V}$. (As in the rest of the paper, we abuse notation by using $\mathbf{A}$ to refer to either of these maps.) In particular, $\mathbf{A}$ is weakly-continuous. We will prove that the convex cone $\mathbf{A}(\hat{\mathcal{Q}}_{++})$ is weakly-closed below. The anticone of $\hat{\mathcal{Q}}_{++} \subset \mathbb{R}^\mathcal{V}$ is precisely $\hat{\mathcal{Q}}_{++} \cap \mathbb{R}_c^\mathcal{V}$. By the theorem above, given any $\mathbf{f} \in \mathbb{R}^\mathcal{V}$, the following statements are equivalent.

- (1) The equation $\mathbf{A}(\mathbf{g}) = \mathbf{f}$ has a solution $\mathbf{g} \in \hat{\mathcal{Q}}_{++}$.
- (2) If $\mathbf{x} \in \mathbb{R}_c^\mathcal{V}$ satisfies $\mathbf{A}(\mathbf{x}) \in \hat{\mathcal{Q}}_{++}$, then $\langle \mathbf{f}, \mathbf{x} \rangle \geq 0$.

Thus, we have proved Lemma 6.35 given the following.

Proposition 13.2. *The convex cone $\mathbf{A}(\hat{\mathcal{Q}}_{++}) \subset \mathbb{R}^\mathcal{V}$ is weakly closed.*

Proof. Suppose $\langle \mathbf{g}_i \in \mathbf{A}(\hat{\mathcal{Q}}_{++}) \rangle$ is a sequence weakly-converging to $\mathbf{g}_\infty \in \mathbb{R}^\mathcal{V}$. We must show that $\mathbf{g}_\infty \in \mathbf{A}(\hat{\mathcal{Q}}_{++})$. We may choose $\mathbf{f}_i \in \hat{\mathcal{Q}}_{++}$ such that $\mathbf{A}(\mathbf{f}_i) = \mathbf{g}_i$ for all i . The idea of the

proof is to use a Cantor diagonalization argument to produce a subsequence of $\langle \mathbf{f}_i \rangle$ which converges to some $\mathbf{f}_\infty$, which necessarily lies in $\hat{\mathcal{Q}}_{++}$. Then, we have $\mathbf{A}(\mathbf{f}_\infty) = \mathbf{g}_\infty$ by the weak-continuity of $\mathbf{A}$. And therefore $\mathbf{g}_\infty \in \mathbf{A}(\hat{\mathcal{Q}}_{++})$.

We enumerate $\mathcal{V} = \{v_1, v_2, \dots\}$. We will first find a subsequence $\langle \mathbf{f}_{i(1,j)} \rangle_j$ of $\langle \mathbf{f}_i \rangle$ such that $\lim_{j \rightarrow \infty} \mathbf{f}_{i(1,j)}$ exists. Let w be a vertex adjacent to v_1 . Then by the formula for $\mathbf{A}(\mathbf{f}_i)$, we know $0 \leq \mathbf{f}_i(v_1) \leq \mathbf{A}(\mathbf{f}_i)(w) = \mathbf{g}_i(w)$. Moreover, $\lim_{i \rightarrow \infty} \mathbf{g}_i(w) \rightarrow \mathbf{g}_\infty(w)$. Thus, for all but finitely many i we have

$$0 \leq \mathbf{f}_i(v_1) \leq \mathbf{g}_\infty(w) + 1.$$

Thus by compactness of the interval $[0, \mathbf{g}_\infty(w) + 1]$, we can find a subsequence $\langle \mathbf{f}_{i(1,j)} \rangle_j$ of $\langle \mathbf{f}_i \rangle$ such that $\lim_{j \rightarrow \infty} \mathbf{f}_{i(1,j)}$ exists. We repeat this argument inductively. For each $n \geq 1$, we can find a subsequence $\langle \mathbf{f}_{i(n+1,j)} \rangle_j$ of $\langle \mathbf{f}_{i(n,j)} \rangle_j$ such that $\lim_{j \rightarrow \infty} \mathbf{f}_{i(n+1,j)}(v_{n+1})$ exists. Then the diagonal sequence $\langle \mathbf{f}_{i(n,n)} \rangle_n$ satisfies $\lim_{n \rightarrow \infty} \mathbf{f}_{i(n,n)}(v_j)$ exists for all j . We set $\mathbf{f}_\infty = \lim_{n \rightarrow \infty} \mathbf{f}_{i(n,n)}$ and proceed as in the previous paragraph. $\square$

The generalized Farkas' theorem given above is also a key point in the proof of Lemma 6.41, which proves the surjectivity of the map $A : \mathcal{S}_\theta \rightarrow \mathcal{S}_{\bar{\theta}}$. This lemma assumes the critical decay property and the adjacency sign property, which allows us to use Proposition 6.40. This proposition gives us a $(\theta, 0)$ -survivor which maps onto any element of $\mathcal{S}_{\bar{\theta}}$.

Proof of Lemma 6.41. Let $\langle g_i \rangle$ and $\langle s_i \rangle$ denote the shrinking and sign sequences of θ . Then $\langle \bar{g}_i \rangle$ and $\langle \bar{s}_i \rangle$ are the corresponding sequences of $\bar{\theta}$. See the beginning of section 6.6. To simplify notation define $\mathbf{g}_i = \Upsilon^{g_i}(\mathbf{g})$.

For each critical time t define the set $H_t = \{\mathbf{h} \in \hat{\mathcal{Q}}_{s_t} : \mathbf{A}(\mathbf{h}) = \mathbf{g}_t\}$. Applying Proposition 6.40 with $\mathbf{g}$ replaced by $\mathbf{g}_t$ and θ replaced by $\pi_{\mathbb{S}^1} \circ \rho_\lambda^{g_t}(\theta)$ implies that H_t is non-empty. For $\mathbf{h} \in H_t$, we have

$$\mathbf{A} \circ \Upsilon^{g_t^{-1}}(\mathbf{h}) = \Upsilon^{g_t^{-1}} \circ \mathbf{A}(\mathbf{h}) = \mathbf{g}.$$

Therefore any $\Upsilon^{g_t^{-1}}(\mathbf{h})$ represents a candidate $\mathbf{f}$. Set $F_t = \Upsilon^{g_t^{-1}}(H_t)$. By definition, we have

$$(13.1) \quad F_t = \{\mathbf{f}_0 \in \mathbb{R}^{\mathcal{V}} : \mathbf{A}(\mathbf{f}_0) = \mathbf{g} \text{ and } \mathbf{f}_0 \in \Upsilon^{g_t}(\hat{\mathcal{Q}}_{s_t})\}.$$

Note that lemma 6.36 implies that

$$F_t = \{\mathbf{f}_0 : \mathbf{A}(\mathbf{f}_0) = \mathbf{g} \text{ and } \mathbf{f}_0 \text{ is a } (\theta, i)\text{-survivor for all } i \leq t\}.$$

Thus, for critical times $t_1 \leq t_2$ we have $F_{t_2} \subset F_{t_1}$. Thus

$$\{\mathbf{f} \in \mathcal{S}_\theta : \mathbf{A}(\mathbf{f}) = \mathbf{g}\} = \bigcap_t F_t,$$

where the intersection is taken over the critical times. We will use the notation $F_\infty = \bigcap_t F_t$. We must show that F_∞ is non-empty.

Note that each F_t is non-empty, because each H_t is. Furthermore, we note that each F_t is weakly-closed. This can be most easily seen by looking at equation 13.1. It follows because both $\mathbf{A}$ and Υ^{g_t} are weakly-continuous and $\hat{\mathcal{Q}}_{s_t}$ is weakly-closed.

For each critical time t choose $\mathbf{f}_t \in F_t$. We will find a weak limit point, $\mathbf{f}_\infty$, of the a subsequence of $\langle \mathbf{f}_t \mid t \text{ a critical time} \rangle$. As each $\mathbf{f}_t$ is a $(\theta, 0)$ -survivor, we know that $\mathbf{f}_t \in \mathcal{Q}_{s_0}$. Without loss of generality, we may assume that $s_0 = ++$. Therefore, $\mathbf{f}_t, \mathbf{g} \in \hat{\mathcal{Q}}_{++}$. For each $v \in \mathcal{V}$, there is a $w \sim v$ and we have

$$0 \leq \mathbf{f}_t(v) \leq \mathbf{A}(\mathbf{f}_t)(w) = \mathbf{g}(w).$$

In particular the sequence of $\mathbf{f}_t(v)$ is bounded for each $v \in \mathcal{V}$. Therefore, we can apply a Cantor diagonal argument as in the proof of Proposition 13.2. This produces our desired limit $\mathbf{f}_\infty$. Since each F_t is closed, and the sequence is nested, we see that $\mathbf{f}_\infty \in F_\infty = \bigcap_t F_t$. $\square$

14. GRAPHS WITHOUT VERTICES OF VALANCE ONE

In this section, we will primarily consider graphs $\mathcal{G}$ with no vertices of valance one. We prove the following results about these graphs.

Theorem 14.1 (Decay). *Suppose $\mathcal{G}$ has no vertices of valance one. Let $\boldsymbol{\theta} \in \mathcal{R}_\lambda$ and let $\langle g_n \rangle$ be the λ -shrinking sequence of $\boldsymbol{\theta}$. Then for any $\mathbf{f} \in \mathcal{S}_{\boldsymbol{\theta}}$ and any $v \in \mathcal{V}$, the sequence*

$$|\Upsilon^{g_n}(\mathbf{f})(v)|$$

tends monotonically to zero as $n \rightarrow \infty$.

We break the proof of this theorem into two parts. We will prove that the sequence is monotonically decreasing in section 14.2. In sections 14.3 and 14.4, we will prove that this limit is zero. Clearly, this theorem implies that any subsequence also decays to zero. Thus, we have the following corollary.

Corollary 14.2 (Decay properties). *If $\mathcal{G}$ has no vertices of valance one, then $S(\mathcal{G}, \mathbf{w})$ has the subsequence decay property and the critical decay property.*

We will prove that graphs with no vertices of valance one have the adjacency sign property in sections 14.5 and 14.6. See Theorem 14.25 and Corollary 14.26

14.1. Facts about eigenfunctions. In this section, we discuss some facts about eigenfunctions of graphs which distinguish graphs with no vertices of valance one. Note that $\mathcal{G}_\mathbb{Z}$ is the only infinite connected graph with no vertices of valance one and no vertices of valance larger than three. Thus, we will pay particular attention to the case of vertices of valance larger than two.

To distinguish graphs with vertices of valance one, we make the following definition.

Definition 14.3. For an integer $k \geq 2$, a k -spoke is a subset $\{v_1, \dots, v_k\} \subset \mathcal{V}$ such that

- (1) $\text{val}(v_1) = 1$.
- (2) For each $i \in \mathbb{N}$ with $2 \leq i \leq k - 1$, $\text{val}(v_i) = 2$, $v_i \sim v_{i-1}$ and $v_i \sim v_{i+1}$.

Note that there is no condition on the valance of the vertex v_k of a k -spoke. See figure 5 for an example.




FIGURE 5. $\{v_1, \dots, v_6\}$ is a 6-spoke.

Proposition 14.4 (Eigenfunctions and spokes). *Let $\{v_1, \dots, v_k\}$ be a spoke. Assume $\mathbf{w} \in \mathbb{R}^V$ is a positive function satisfying $\mathbf{A}\mathbf{w} = \lambda\mathbf{w}$. If $\lambda = 2$, then for all j with $2 \leq j \leq k$, we have $\mathbf{w}(v_j) = j\mathbf{w}(v_1)$. If $\lambda > 2$ then*

$$\mathbf{w}(v_j) = \frac{\sinh(jz)}{\sinh(z)} \mathbf{w}(v_1) \quad \text{where } z = \ln \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} = \cosh^{-1}\left(\frac{\lambda}{2}\right).$$

Sketch of proof. Note that $\mathbf{w}(v_2) = \lambda\mathbf{w}(v_1)$ and $\mathbf{w}(v_j) = \lambda\mathbf{w}(v_{j-1}) - \mathbf{w}(v_{j-2})$. Thus, the values of $\mathbf{w}(v_j)$ are determined by the previous values, and thus the value of $\mathbf{w}(v_j)$ is uniquely determined by the value of $\mathbf{w}(v_1)$. Finally by inspection and trigonometry, it can be checked that the values stated in the proposition do give rise to a solution to these equations. $\square$

Note that every vertex of valance one belongs to a spoke. The following handles vertices with greater valance.

Proposition 14.5 (Detecting spokes). *Let $\mathbf{w}$ be a positive function satisfying $\mathbf{A}\mathbf{w} = \lambda\mathbf{w}$. Let x be a vertex with $\text{val}(x) \geq 2$. Then, x belongs to a spoke if and only if there is a $y \sim x$ such that*

$$\frac{\mathbf{w}(y)}{\mathbf{w}(x)} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}.$$

Proof. We will assume $\lambda > 2$. (There are only three infinite connected graphs with a positive eigenfunction with eigenvalue 2.)

First suppose that $x = v_j$ belongs to a spoke. Let $y = v_{j-1}$. Then by proposition 14.4 and angle addition formulas, we know that

$$\frac{\mathbf{w}(y)}{\mathbf{w}(x)} = \frac{\sinh((j-1)z)}{\sinh(jz)} = \frac{\sinh(jz) \cosh z - \cosh(jz) \sinh(z)}{\sinh(jz)}.$$

We have $\cosh(z) = \lambda/2$ and $\sinh(z) = \sqrt{\lambda^2 - 4}/2$. Thus,

$$\frac{\mathbf{w}(y)}{\mathbf{w}(x)} = \frac{\lambda}{2} - \frac{\cosh(jz)\sqrt{\lambda^2 - 4}}{2\sinh(jz)} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}.$$

Now we will approach the converse. First, we claim that if a and b are two adjacent vertices, with

$$(14.1) \quad \frac{\mathbf{w}(b)}{\mathbf{w}(a)} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2},$$

then $\text{val}(b) \leq 2$. Suppose $\text{val}(b) = k \geq 3$. Let a and $c_1, \dots, c_{k-1}$ be the vertices adjacent to b . We have that

$$\lambda = \frac{\mathbf{w}(a) + \sum_{i=1}^{k-1} \mathbf{w}(c_i)}{\mathbf{w}(b)} > \frac{\lambda + \sqrt{\lambda^2 - 4}}{2} + \frac{\sum_{i=1}^{k-1} \mathbf{w}(c_i)}{\mathbf{w}(b)}.$$

Thus, there is a j such that

$$\frac{\mathbf{w}(c_j)}{\mathbf{w}(b)} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2(k-1)} = \frac{2}{(k-1)(\lambda + \sqrt{\lambda^2 - 4})} \leq \frac{1}{\lambda + \sqrt{\lambda^2 - 4}} \leq \frac{1}{\lambda}.$$

In summary $\mathbf{w}(c_j) < \mathbf{w}(b)/\lambda$. Therefore,

$$(\mathbf{A}\mathbf{w})(c_j) = \sum_{d \sim c_j} \mathbf{w}(d) \geq \mathbf{w}(b) > \lambda\mathbf{w}(c_j),$$

which contradicts our assumption that $\mathbf{A}\mathbf{w} = \lambda\mathbf{w}$.

Now suppose $a \sim b$, $\text{val}(b) = 2$ and equation 14.1 holds. We will show a belongs to a spoke, completing the proof. Let c denote the other vertex adjacent to b . We have $\lambda\mathbf{w}(b) = \mathbf{w}(a) + \mathbf{w}(c)$. Thus,

$$\frac{\mathbf{w}(c)}{\mathbf{w}(b)} = \lambda - \frac{\mathbf{w}(a)}{\mathbf{w}(b)} > \lambda - \left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right)^{-1} = \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}.$$

Thus, equation 14.1 is satisfied with a replaced by b and b replaced by c . By the claim above, we know $\text{val}(c) \leq 2$. By induction, we see that either $\{a, b, c\}$ is a subset of a spoke, or there is an infinite sequence of vertices $\{x_0 = a, x_1 = b, x_2 = c, x_3, x_4, \dots\}$ with each x_j for $j \geq 1$ satisfying $\text{val}(x_j) = 2$, $x_j \sim x_{j-1}$, and $x_j \sim x_{j+1}$. We will show that $\{a, b, c\}$ must be a subset of a spoke, by proving this other possibility is false. Note that the values of $\mathbf{w}(x_j)$ is uniquely determined by $\mathbf{w}(x_{j-1})$ and $\mathbf{w}(x_{j-2})$ according to the rule $\mathbf{w}(x_j) + \mathbf{w}(x_{j-2}) = \lambda\mathbf{w}(x_{j-1})$. In particular, the value of each $\mathbf{w}(x_j)$ may be determined inductively from $\mathbf{w}(x_0)$ and $\mathbf{w}(x_1)$. Any such solution can be written as

$$\mathbf{w}(x_j) = r\left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}\right)^j + s\left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right)^j$$

for all j and some $r, s \in \mathbb{R}$. We will now solve for r and s . We have $\mathbf{w}(x_0) = r + s$. Then,

$$\begin{aligned} \frac{\mathbf{w}(b)}{\mathbf{w}(a)} &= \frac{\mathbf{w}(x_1)}{\mathbf{w}(x_0)} = \frac{r\left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}\right) + s\left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right)}{\mathbf{w}(x_0)} = \\ &= \frac{r\left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}\right) + (\mathbf{w}(x_0) - r)\left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right)}{\mathbf{w}(x_0)} = \left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right) + r\sqrt{\lambda^2 - 4}. \end{aligned}$$

Thus, by equation 14.1, we have $r < 0$. It follows that there is a $j \in \mathbb{N}$ such that $\mathbf{w}(x_j) < 0$. This contradicts our initial assumption that $\mathbf{w}$ is a positive eigenfunction. Thus, a is an element of a spoke. $\square$

Recall that the constant r is the spectral radius of the adjacency operator acting on $\ell_2(\mathcal{V})$. It is also the minimal eigenvalue of a positive eigenfunction.

Corollary 14.6. *Suppose $\mathcal{G}$ contains a vertex of valance n which is not a member of a spoke. Then the spectral radius of the adjacency operator satisfies $r \geq \frac{n}{\sqrt{n-1}}$.*

Proof. Let v be the vertex of valance n which is not a member of a spoke, and let $\mathbf{f}$ be a positive eigenfunction. Then by proposition 14.5, if $w \sim v$ we know $\frac{\mathbf{f}(w)}{\mathbf{f}(v)} \geq \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}$. Since $\sum_{w \sim v} \mathbf{f}(w) = \lambda\mathbf{f}(v)$, we know that

$$\lambda \geq n\left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right).$$

This is equivalent to the inequality given in the corollary. $\square$

14.2. Monotonic decay. In this subsection we prove the monotonicity part of Theorem 14.1.

Lemma 14.7 (Monotonic Decay). *Suppose $\mathcal{G}$ has no vertices of valance one. Let $\boldsymbol{\theta} \in \mathbb{S}^1$ be a λ -renormalizable direction with λ -shrinking sequence $\langle g_n \rangle$. For every $\boldsymbol{\theta}$ -survivor $\mathbf{f} \in \mathbb{R}^{\mathcal{V}}$ and every $v \in \mathcal{V}$, the sequence $|\Upsilon^{g_n}(\mathbf{f})(v)|$ decreases (non-strictly) monotonically in n .*

We will simplify the statement of this lemma. Applying Corollary 11.7, we know that $\Upsilon^{g_1}(\mathbf{f})$ is a $\rho_\lambda^{g_1}(\boldsymbol{\theta})$ -survivor. By induction, it is sufficient to prove that

$$(14.2) \quad |\Upsilon^{g_1}(\mathbf{f})(v)| \leq |\mathbf{f}(v)|,$$

for all λ -renormalizable $\boldsymbol{\theta}$, all $\boldsymbol{\theta}$ -survivors $\mathbf{f}$, and all $v \in \mathcal{V}$. Up to the dihedral group action, we may assume that $\boldsymbol{\theta} \in \mathcal{Q}_{++}$ and that $g_1 = h^{-1}$. See Remark 2.6. Since $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor, we know $\mathbf{f} \in \hat{\mathcal{Q}}_{++}$. By equation 6.2, we know $\mathbf{H}(\mathbf{f})(b) = b$ for all $b \in \mathcal{B}$. So it suffices to consider $v \in \mathcal{A}$. Suppose that $\rho_\lambda^{h^{-1}}(\boldsymbol{\theta}) \in \mathcal{Q}_{++}$. Recalling the formula in equation 6.2 for $\mathbf{H}(\mathbf{f})$, we have that for $a \in \mathcal{A}$,

$$\mathbf{H}^{-1}(\mathbf{f})(a) = \mathbf{f}(a) - \sum_{b \sim a} \mathbf{f}(b) \geq 0.$$

But, each $\mathbf{f}(b) \geq 0$ as $\mathbf{f} \in \hat{\mathcal{Q}}_{++}$, so $\mathbf{H}^{-1}(\mathbf{f})(a) \leq \mathbf{f}(a)$. Hence, equation 14.2 is trivially true when $\rho_\lambda^{h^{-1}}(\boldsymbol{\theta}) \in \mathcal{Q}_{++}$. By Proposition 10.6, the alternative is that $\rho_\lambda^{h^{-1}}(\boldsymbol{\theta}) \in \mathcal{Q}_{-+}$. In this case $\mathbf{H}^{-1}(\mathbf{f})(a) < 0$, so equation 14.2 is equivalent to showing

$$(14.3) \quad \mathbf{f}(a) \geq -\mathbf{H}^{-1}(\mathbf{f})(a).$$

In fact, we have that $\mathbf{H}^{-1}(\mathbf{f})(a) = \mathbf{f}(a) - \mathbf{A}(\mathbf{f})(a)$ by Proposition 6.33. Thus, equation 14.3 is equivalent to showing that $2\mathbf{f}(a) \geq \mathbf{A}(\mathbf{f})(a)$. Perhaps more usefully, we have

$$2\mathbf{f}(a) - \mathbf{A}(\mathbf{f})(a) = 2\langle \mathbf{f}, \mathbf{e}_a \rangle - \langle \mathbf{A}(\mathbf{f}), \mathbf{e}_a \rangle = \langle \mathbf{f}, 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a) \rangle.$$

In summary, the following lemma implies the Monotonic Decay lemma (14.7) above.

Lemma 14.8. *Suppose $\text{val}(\mathcal{G}) \geq 2$. Let $\boldsymbol{\theta} \in \mathbb{S}^1 \cap \mathcal{Q}_{++}$ be a λ -renormalizable direction with shrinking sequence $\langle g_n \rangle$. Assume that $g_1 = h^{-1}$. Then for every $\boldsymbol{\theta}$ -survivor $\mathbf{f}$ and every $a \in \mathcal{A}$, we have $\langle \mathbf{f}, 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a) \rangle \geq 0$.*

Remark 14.9. In the paragraph preceding the lemma, we indicated that the case of interest is when $\rho_\lambda^{h^{-1}}(\boldsymbol{\theta}) \in \mathcal{Q}_{-+}$. The conclusion of the lemma follows trivially when $\rho_\lambda^{h^{-1}}(\boldsymbol{\theta}) \in \mathcal{Q}_{++}$. This is because the conclusion of the lemma is equivalent to equation 14.3. When $\mathbf{H}^{-1}(\mathbf{f})(a) \geq 0$, equation 14.3 holds for trivial reasons.

Recall, the *valance* of a vertex $v \in \mathcal{V}$ is the number $\text{val}(v) = \#\{x \in \mathcal{V} : x \sim v\}$.

Proposition 14.10. *The conclusion of lemma 14.8 holds when a has valance larger than three and is not a member of a spoke.*

Proof. By assumption $g_1 = h^{-1}$. Thus, by Proposition 7.13, we know $\boldsymbol{\theta} = (x, y) \in \mathbb{S}^1$ satisfies

$$0 < \frac{y}{x} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}.$$

We will show that there exists two saddle connections σ_1 and σ_2 such that

- (1) $\mathcal{Z}(\llbracket \sigma_1 \rrbracket + \llbracket \sigma_2 \rrbracket) = 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a)$, and
- (2) $\text{hol}(\sigma_1) \wedge \boldsymbol{\theta} \geq 0$ and $\text{hol}(\sigma_2) \wedge \boldsymbol{\theta} \geq 0$.

We will show these two statements imply the proposition. Statement (1) implies that

$$\langle \mathbf{f}, 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a) \rangle = \langle \mathbf{f}, \mathcal{Z}(\llbracket \sigma_1 \rrbracket + \llbracket \sigma_2 \rrbracket) \rangle.$$

We wish to show that this quantity is non-negative. This follows from statement (2) of Proposition 11.10 with $\mathbf{x} = \mathcal{Z}(\llbracket \sigma_1 \rrbracket + \llbracket \sigma_2 \rrbracket)$. Thus, the existence of such σ_1 and σ_2 imply the proposition.

Consider our surface $S = S(\mathcal{G}, \mathbf{w})$, where $\mathbf{A}\mathbf{w} = \lambda\mathbf{w}$. We may assume that $\mathbf{w}(\mathbf{a}) = 1$ by scaling $\mathbf{w}$ if necessary. The cylinder $cyl_{\mathbf{a}}$ has a decomposition into rectangles of the form

$$cyl_{\mathbf{a}} = \bigcup_{\mathbf{b} \sim \mathbf{a}} (cyl_{\mathbf{a}} \cap cyl_{\mathbf{b}}).$$

Let $k = val(\mathbf{a}) \geq 3$. We may number these rectangles $R_1, \dots, R_k$ so that each R_i is adjacent to $R_{i+1 \pmod k}$. Similarly, number the relevant vertices $\mathbf{b}_1, \dots, \mathbf{b}_k \in \mathcal{B}$ so that $R_i = cyl_{\mathbf{a}} \cap cyl_{\mathbf{b}_i}$ for all i . Choose $j \in \{1, \dots, k\}$ so that

$$(14.4) \quad \mathbf{w}(\mathbf{b}_j) + \mathbf{w}(\mathbf{b}_{j+1 \pmod k}) = \min\{\mathbf{w}(\mathbf{b}_i) + \mathbf{w}(\mathbf{b}_{i+1 \pmod k}) : i = 1, \dots, k\}.$$

Choose σ_1 to be the diagonal of the rectangle $R_j \cup R_{j+1 \pmod k}$ which can be oriented downward and leftward. Choose σ_2 to be the diagonal of the complimentary rectangle, $cyl_{\mathbf{a}} \setminus (R_j \cup R_{j+1 \pmod k})$, oriented downward and leftward. See figure 6.




FIGURE 6. An example decomposition of $cyl_{\mathbf{a}}$ into rectangles. In this case, $val(\mathbf{a}) = 4$ and $j = 3$ is the index satisfying equation 14.4.

We now verify statements (1) and (2) hold for our choices of σ_1 and σ_2 . Note that $\sigma_1 \cup \sigma_2$ cross $cyl_{\mathbf{a}}$ twice with positive algebraic sign, and cross each of $cyl_{\mathbf{b}}$ for $\mathbf{b} \sim \mathbf{a}$ once with negative algebraic sign. Moreover, $\sigma_1 \cup \sigma_2$ cross no other cylinders. This verifies statement (1).

Now we verify statement (2). Write $hol(\sigma_1) = (-w_1, -1)$ and $hol(\sigma_2) = (-w_2, -1)$ with $w_1 > 0$ and $w_2 > 0$. Now recall that $\boldsymbol{\theta} = (x, y) \in S^1 \cap \mathcal{Q}_{++}$ is λ -renormalizable and that $g_1 = h^{-1}$. Thus, Proposition 7.13 implies that $\frac{y}{x} < \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}$. For $i = 1, 2$, we have

$$(14.5) \quad hol(\sigma_i) \wedge \boldsymbol{\theta} = (-w_i, -1) \wedge (x, y) = x - w_i y = x(1 - w_i \frac{y}{x}) > \frac{x}{2} (2 - w_i (\lambda - \sqrt{\lambda^2 - 4})).$$

(Note $x > 0$ since $\boldsymbol{\theta} \in \mathcal{Q}_{++}$.) To proceed we must find upper bounds for w_1 and w_2 . Because of equation 14.4, we know that

$$w_1 \leq \frac{2\lambda}{val(\mathbf{a})} \leq \frac{2\lambda}{3}.$$

(w_1 must be less than or equal to the average length of pairs of adjacent rectangles.) Therefore, by continuing equation 14.5 for $i = 1$ we have

$$\begin{aligned} hol(\sigma_1) \wedge \boldsymbol{\theta} &> \frac{x}{2} \left(2 - \left(\frac{2\lambda}{3} \right) (\lambda - \sqrt{\lambda^2 - 4}) \right) = \frac{x}{3} (3 + \lambda \sqrt{\lambda^2 - 4} - \lambda^2) \\ &= \frac{x}{3} \left(\frac{2\lambda^2 - 9}{\lambda \sqrt{\lambda^2 - 4} + \lambda^2 - 3} \right) \geq 0. \end{aligned}$$

Here, the ≥ 0 statement is somewhat subtle. The denominator $\lambda \sqrt{\lambda^2 - 4} + \lambda^2 - 3$ is positive because $\lambda \geq 2$, and the numerator $2\lambda^2 - 9 \geq 0$ because $\lambda \geq \frac{3\sqrt{2}}{2}$ by Corollary 14.6. To get

an upper bound for w_2 we find a lower bound for w_1 . We know a is not a member of a spoke and σ_1 crosses two rectangles, so by Proposition 14.5 we have

$$w_1 \geq \lambda - \sqrt{\lambda^2 - 4}.$$

Since the cylinder cyl_a has inverse modulus λ and width one, we know $w_1 + w_2 = \lambda$. Therefore $w_2 \leq \sqrt{\lambda^2 - 4}$. By continuing equation 14.5 for $i = 2$ we have

$$\begin{aligned} hol(\sigma_2) \wedge \boldsymbol{\theta} &> \frac{x}{2}(2 - (\sqrt{\lambda^2 - 4})(\lambda - \sqrt{\lambda^2 - 4})) = \frac{x}{2}(\lambda^2 - 2 - \lambda\sqrt{\lambda^2 - 4}) \\ &= \frac{x}{2}\left(\frac{4}{\lambda^2 - 2 - \lambda\sqrt{\lambda^2 - 4}}\right) > 0. \end{aligned}$$

This proves statement (2) and concludes the proof. $\square$

Proposition 14.11. *Lemma 14.8 holds when $val(a) = 2$ unless a belongs to a spoke.*

Proof. Let b_1 and b_2 denote the two vertices adjacent to a . For $i = 1, 2$, let σ_i denote the diagonal of rectangle $cyl_a \cap cyl_{b_i}$ which can be oriented downward and leftward. We observe that $\mathcal{Z}(\llbracket \sigma_1 \rrbracket + \llbracket \sigma_2 \rrbracket) = 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a)$. Write $\boldsymbol{\theta} = (x, y) \in \mathcal{Q}_{++}$. Because $g_1 = h^{-1}$, Proposition 7.13 implies that $y < x(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2})$. For $i = 1, 2$, we have $hol(\sigma_i) = (-\mathbf{w}(b_i), -\mathbf{w}(a))$. By Proposition 14.5, $\mathbf{w}(b_i) \geq \mathbf{w}(a)(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2})$. We have $\mathbf{w}(b_1) + \mathbf{w}(b_2) = \lambda\mathbf{w}(a)$. Thus,

$$\mathbf{w}(b_1) = \lambda\mathbf{w}(a) - \mathbf{w}(b_2) \leq \mathbf{w}(a)\left(\lambda - \frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right) = \mathbf{w}(a)\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}.$$

And similarly, $\mathbf{w}(b_2) \leq \mathbf{w}(a)(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2})$. Thus, we compute

$$hol(\sigma_i) \wedge \boldsymbol{\theta} = x\mathbf{w}(a) - y\mathbf{w}(b_i) > x\mathbf{w}(a) - \left(x\left(\frac{\lambda - \sqrt{\lambda^2 - 4}}{2}\right)\right)\left(\mathbf{w}(a)\left(\frac{\lambda + \sqrt{\lambda^2 - 4}}{2}\right)\right) = 0.$$

Thus, if $\mathbf{f}$ is a $\boldsymbol{\theta}$ -survivor, $\langle \mathbf{f}, \mathcal{Z}(\llbracket \sigma_i \rrbracket) \rangle > 0$. And therefore $\langle \mathbf{f}, 2\mathbf{e}_a - \mathbf{A}(\mathbf{e}_a) \rangle \geq 0$ as desired. $\square$

14.3. Effective decay for the integers. In this subsection, we will only consider the case when $\mathcal{G} = \mathcal{G}_{\mathbb{Z}}$, the graph whose vertex set is $\mathbb{Z}$ and edges join two integers if and only if they differ by one. Our decomposition $\mathcal{V} = \mathcal{A} \cup \mathcal{B}$ is a decomposition into even and odd integers.

Lemma 14.12 (Effective decay for the integers). *Let $\langle g_i \rangle$ be a shrinking sequence for a renormalizable direction $\boldsymbol{\theta}$. Then, there is a critical time $t > 0$ for which*

$$|\Upsilon^{g_t}(\mathbf{f})(v)| \leq \frac{1}{2}|\mathbf{f}(v)|$$

for any $\boldsymbol{\theta}$ -survivor $\mathbf{f}$ and any $v \in \mathbb{Z}$.

The idea of the proof is the following. We claim it is sufficient to find a critical time t for which

$$(14.6) \quad |\Upsilon^{\gamma(g_t)}(\mathbf{e}_v)(v)| \geq 2 \quad \text{for all } v \in \mathbb{Z}.$$

(The bulk of this section will be spent proving equation 14.6.) Let $\mathbf{e}_v \in \mathbb{R}_{\mathbb{C}}^{\mathbb{Z}}$ be as in equation 11.4. Set $\mathbf{e}'_v = \pm \mathbf{e}_v \in \hat{\mathcal{Q}}_s$ where the sign is chosen depending on the quadrant $\mathcal{Q}_s$ containing $\boldsymbol{\theta}$. By equation 6.14, we have

$$(14.7) \quad |\mathbf{f}(v)| = \langle \mathbf{f}, \mathbf{e}'_v \rangle = \langle \Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{e}'_v) \rangle = \sum_{w \in \mathcal{V}} (\Upsilon^{g_t}(\mathbf{f})(w)) (\Upsilon^{\gamma(g_t)}(\mathbf{e}'_v)(w)).$$

By Proposition 11.12, we have $\Upsilon^{\gamma(g_t)}(\mathbf{e}'_v) \in \hat{\mathcal{Q}}_{st}$ for all critical times t . Since both $\Upsilon^{g_t}(\mathbf{f})$ and $\Upsilon^{\gamma(g_t)}(\mathbf{e}'_v)$ lie in $\hat{\mathcal{Q}}_{st}$, all terms in the sum above are non-negative. Therefore,

$$|\mathbf{f}(v)| = \langle \Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{e}'_v) \rangle \geq (\Upsilon^{g_t}(\mathbf{f})(v)) (\Upsilon^{\gamma(g_t)}(\mathbf{e}'_v)(v)) \geq 2|\Upsilon^{g_t}(\mathbf{f})(v)|,$$

as claimed by the lemma.

Proposition 14.13. *Consider the geodesic ray defined*

$$g_n = \begin{cases} (v^{-1}h)^{\frac{n}{2}} & \text{if } n \text{ is even} \\ h(v^{-1}h)^{\frac{n-1}{2}} & \text{if } n \text{ is odd.} \end{cases}$$

Then for all $n \geq 1$ and all $a \in \mathcal{A}$ and $b \in \mathcal{B}$,

$$\begin{aligned} \Upsilon^{g_n}(\mathbf{e}_a) &= \begin{cases} \sum_{c \in [a-n+1, a+n-1]} \mathbf{e}_c & \text{if } n \equiv 1 \pmod{4}, \\ \sum_{c \in \mathcal{A} \cap [a-n+1, a+n-1]} \mathbf{e}_c - \sum_{c \in \mathcal{B} \cap [a-n+1, a+n-1]} \mathbf{e}_c & \text{if } n \equiv 2 \pmod{4}, \\ -\sum_{c \in [a-n+1, a+n-1]} \mathbf{e}_c & \text{if } n \equiv 3 \pmod{4}, \\ -\sum_{c \in \mathcal{A} \cap [a-n+1, a+n-1]} \mathbf{e}_c + \sum_{c \in \mathcal{B} \cap [a-n+1, a+n-1]} \mathbf{e}_c & \text{if } n \equiv 0 \pmod{4}. \end{cases} \\ \Upsilon^{g_n}(\mathbf{e}_b) &= \begin{cases} \sum_{c \in [b-n, b+n]} \mathbf{e}_c & \text{if } n \equiv 1 \pmod{4}, \\ \sum_{c \in \mathcal{A} \cap [b-n, b+n]} \mathbf{e}_c - \sum_{c \in \mathcal{B} \cap [b-n, b+n]} \mathbf{e}_c & \text{if } n \equiv 2 \pmod{4}, \\ -\sum_{c \in [b-n, b+n]} \mathbf{e}_c & \text{if } n \equiv 3 \pmod{4}, \\ -\sum_{c \in \mathcal{A} \cap [b-n, b+n]} \mathbf{e}_c + \sum_{c \in \mathcal{B} \cap [b-n, b+n]} \mathbf{e}_c & \text{if } n \equiv 0 \pmod{4}. \end{cases} \end{aligned}$$

It is straightforward to check that the formulas in the proposition above follow from the definition of the action Υ^G .

Lemma 14.14. *Let $\langle g_n \rangle$ be a geodesic ray for which $\rho_\lambda^{g_1}(\mathcal{Q}_s) \subset \mathcal{Q}_s$. Then for all $\mathbf{x} \in \mathbb{R}_c^\vee \cap \hat{\mathcal{Q}}_s$ and all $v \in \mathbb{R}_c^\vee$, the sequence $|\Upsilon^{g_n}(\mathbf{x})(v)|$ is non-decreasing.*

Proof. Consider the orbit under the sign action, $s_n = \Sigma^{g_n}(s)$. Note that the condition that $\rho_\lambda^{g_1}(\mathcal{Q}_s) \subset \mathcal{Q}_s$ guarantees that $\Upsilon^{g_n}(\hat{\mathcal{Q}}_s) \subset \hat{\mathcal{Q}}_{s_n}$ for all n , by Proposition 11.4.

Suppose the lemma is false. Then there is a geodesic ray $\langle g_n \rangle$ such that $|\Upsilon^{g_n}(\mathbf{x})(v)| < |\Upsilon^{g_{n-1}}(\mathbf{x})(v)|$. We may assume that n is minimal over all geodesic rays, all $\mathbf{x}$ and all possible choices of n . Since the statement is invariant under the dihedral group, we may assume that $g_1 = h$ and $s = ++$. We will show that

$$(14.8) \quad g_n = \begin{cases} (v^{-1}h)^{\frac{n}{2}} & \text{if } n \text{ is even} \\ h(v^{-1}h)^{\frac{n-1}{2}} & \text{if } n \text{ is odd.} \end{cases}$$

Otherwise, there is a $i \leq n-2$ for which $g_{i+2}g_i^{-1} \in \{vh, h^2, h^{-1}v^{-1}, v^{-2}\}$. (This is the first i for which g_{i+2} differs from the form above.) In this case, $s_{i+1} = s_{i+2}$, by the definition of the sign action. Moreover, $\rho_\lambda^{g_{i+2}g_i^{-1}}(\mathcal{Q}_{s_{i+1}}) \subset \mathcal{Q}_{s_{i+1}}$. So letting $\mathbf{y} = \Upsilon^{g_{i+1}}(\mathbf{x})$ and considering the geodesic ray $\langle g_{i+1+n}g_{i+1}^{-1} \rangle_n$ gives a shorter counter example.

Finally the case of g_n as in equation 14.8 follows from Proposition 14.13 above. Observe that

$$\Upsilon^{g_n}(\mathbf{x})(v) = \sum_{w \in \mathcal{V}} \mathbf{x}(w) \Upsilon^{g_n}(\mathbf{e}_w)(v).$$

Note that the sign of each non-zero term is only dependent on n , and that by Proposition 14.13 the sequence $|\Upsilon^{g_n}(\mathbf{e}_w)(v)|$ is non-decreasing regardless of the choice of w . $\square$

This proposition further lowers the bar for proving Lemma 14.12. It is sufficient to find any time n for which

$$(14.9) \quad |\Upsilon^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2 \quad \text{for all } v \in \mathbb{Z}.$$

We will then apply Lemma 14.14. We know that $\mathcal{Q}_{s_0}$ contains the vector $\boldsymbol{\theta}$, which is shrunk by $\rho_\lambda^{g_1}$. Therefore, Corollaries 7.4 and 7.5 imply that $\rho_\lambda^{\gamma(g_1)}(\mathcal{Q}_{s_0}) \subset \mathcal{Q}_{s_0}$. Thus, applies Then by Lemma 14.14, given any critical time $t \geq n$, we have $|\Upsilon^{\gamma(g_t)}(\mathbf{e}_v)(v)| > |\Upsilon^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2$. (We have such a critical time because of Corollary 10.9.) Thus, equation 14.9 implies equation 14.6 which implies Lemma 14.12.

Recall Definition 3.7 of a renormalizing sequence. (It is the possible shrinking sequences for a λ -renormalizable direction.)

Proposition 14.15. *Let $\langle g_n \rangle$ be any renormalizing sequence. Then there is an n for which g_n is of one of the following forms.*

- (1) $g_n = h^c v^b h^a$ for some $a \in \mathbb{Z}$ and non-zero $b, c \in \mathbb{Z}$ such that $(b, c) \notin \{(1, -1), (-1, 1)\}$.
- (2) $g_n = h^{-b} (h^{-b} v^b)^d h^a$ for some $a \in \mathbb{Z}$, $b \in \{\pm 1\}$, and $d > 0$.
- (3) $g_n = h^f v^e (h^{-b} v^b)^d h^a$ for some $a \in \mathbb{Z}$, $b \in \{\pm 1\}$, $d > 0$, and nonzero $e, f \in \mathbb{Z}$ such that $(e, f) \neq (b, -b)$.

Moreover, for any such g_n , we have $|\Upsilon^{g_n}(\mathbf{e}_v)(v)| \geq 2$ for all $v \in \mathcal{A}$.

Proof. There is an $a \in \mathbb{Z}$ (possibly zero) of maximal absolute value such that $h^a = g_{|a|}$. (A renormalizing sequence cannot be $g_n = h^{\pm n}$ for all n , by definition.) For all $v \in \mathcal{A}$ and all $a \in \mathbb{Z}$, we have $\mathbf{H}^a(\mathbf{e}_v) = \mathbf{e}_v$ by definition of $\mathbf{H}$. See equation 6.2. Then to be a geodesic ray, one of $vh^a, v^{-1}h^a \in \{g_n\}$. Thus, there is a non-zero b of maximal absolute value such that $v^b h^a = g_{|a|+|b|}$. By definition of $\mathbf{V}$, we have

$$\Upsilon^{g_{|a|+|b|}}(\mathbf{e}_v) = \mathbf{V}^b \circ \mathbf{H}^a(\mathbf{e}_v) = \mathbf{e}_v + b(\mathbf{e}_{v-1} + \mathbf{e}_{v+1}).$$

Then there is a non-zero c of maximal absolute value such that $h^c v^b h^a = g_{|a|+|b|+|c|}$. We have

$$\Upsilon^{g_{|a|+|b|+|c|}}(\mathbf{e}_v) = \mathbf{e}_v + b(\mathbf{e}_{v-1} + \mathbf{e}_{v+1}) + bc(\mathbf{e}_{v-2} + 2\mathbf{e}_v + \mathbf{e}_{v+2}).$$

Therefore $|\Upsilon^{g_{|a|+|b|+|c|}}(\mathbf{e}_v)(v)| = |1 + 2bc|$. This quantity is larger than one unless $(b, c) \in \{(1, -1), (-1, 1)\}$. This handles the case (1) of the proposition.

If case (1) does not apply, then $b = \pm 1$, and $c = -b$. There is a maximal integer $d \geq 1$ such that $(h^{-b} v^b)^d h^a = g_{|a|+2d}$. By conjugating by an element of the dihedral group and applying Proposition 14.13, we see that for some $\alpha, \beta \in \{\pm 1\}$ we have

$$\Upsilon^{g_{|a|+2d}}(\mathbf{e}_v)(v) = \Upsilon^{g_{|a|+2d}}(\mathbf{e}_v)(v \pm 2) = \alpha \quad \text{and} \quad \Upsilon^{g_{|a|+2d}}(\mathbf{e}_v)(v \pm 1) = \beta.$$

The choices of α and β are given by the following rules that

$$\alpha = (-1)^d \quad \text{and} \quad \beta = -b(-1)^d.$$

The element $g_{|a|+2d+1}$ must be given by either $h^{-b} g_{|a|+2d}$ or $v^{\pm 1} g_{|a|+2d}$. Assume that $g_{|a|+2d+1} = h^{-b} g_{|a|+2d+1}$. Then,

$$\Upsilon^{g_{|a|+2d+1}}(\mathbf{e}_v)(v) = \Upsilon^{g_{|a|+2d}}(\mathbf{e}_v)(v) - 2\Upsilon^{g_{|a|+2d+1}}(\mathbf{e}_v)(v \pm 1) = \alpha - 2b\beta = 3(-1)^d,$$

which is of absolute value larger than 2. This handles case (2).

The only remaining possibility is that $g_{|a|+2d+1} = v^{\pm 1} g_{|a|+2d}$. Then there is a nonzero e of maximal magnitude for which $g_{|a|+2d+|e|} = v^e g_{|a|+2d}$. We compute

$$\Upsilon^{g_{|a|+2d+|e|}}(\mathbf{e}_v)(v) = \alpha \quad \text{and} \quad \Upsilon^{g_{|a|+2d+|e|}}(\mathbf{e}_v)(v \pm 1) = \beta - 2e\alpha = \alpha(-b + 2e).$$

There is then an integer $f \neq 0$ such that $h^f g_{|a|+2d+|e|} = g_{|a|+2d+|e|+|f|}$. We compute that

$$\Upsilon^{g_{|a|+2d+|e|+|f|}}(\mathbf{e}_v)(v) = \alpha + 2f \Upsilon^{g_{|a|+2d+|e|}}(\mathbf{e}_v)(v \pm 1) = \alpha(1 + 2f(-b + 2e)).$$

This quantity has magnitude one only if $f(-b + 2e) = -1$. Therefore, we must have $e = b$, and $f = -b$. This handles case (3). Finally, if $e = b$ and $f = -b$, then

$$g_{|a|+2d+|e|+|f|} = h^{-b} v^b (h^{-b} v^b)^d h^a,$$

which contradicts the maximality of d . $\square$

Corollary 14.16. *Let $\langle g_n \rangle$ be any renormalizing sequence. Then there is an N such that for all $n \geq N$.*

$$|\Upsilon^{g_n}(\mathbf{e}_v)(v)| \geq 2$$

for all $v \in \mathcal{V}$.

Proof. Applying Proposition 14.15, there is an integer n_1 for which $|\Upsilon^{g_{n_1}}(\mathbf{e}_a)(a)| \geq 2$ for all $a \in \mathcal{A}$. By applying dihedral symmetry and Proposition 14.15 again, there is a integer n_2 for which $|\Upsilon^{\gamma(g_{n_2})}(\mathbf{e}_b)(b)| \geq 2$ for all $b \in \mathcal{B}$. Set $n = \max \{n_1, n_2\}$. By Lemma 14.14, for all $v \in \mathcal{V}$ we have $|\Upsilon^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2$. $\square$

Proof of Lemma 14.12 (Effective decay for the integers). From the discussion below the statement of the lemma, we see it is sufficient to prove the statement in equation 14.9. Since θ is renormalizable, its shrinking sequence $\langle g_n \rangle$ is renormalizable. The automorphism γ sends renormalizable sequences to renormalizable sequences. Therefore, the sequence $\langle \gamma(g_n) \rangle$ is also renormalizable. Thus, equation 14.9 follows directly from Corollary 14.16. $\square$

14.4. Effective decay for graphs without vertices of valance one. Essentially, we use covers and subgraphs to deduce effective decay for graphs with no vertices of valance one from effective decay for $\mathcal{G}_{\mathbb{Z}}$.

Lemma 14.17 (Effective decay). *Let $\mathcal{G}$ be any infinite connected bipartite graph with bounded valance and without vertices of valance one. Let $\langle g_i \rangle$ be a shrinking sequence for a renormalizable direction θ . Then, there is a critical time $t > 0$ for which*

$$|\Upsilon_{\mathcal{G}}^{g_t}(\mathbf{f})(v)| \leq \frac{1}{2} |\mathbf{f}(v)|$$

for any θ -survivor $\mathbf{f} \in \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ and any $v \in \mathcal{V}_{\mathcal{G}}$.

Proof. The sequence $\langle \gamma(g_n) \rangle$ is a renormalizable sequence. By Corollary 14.16, there is an N such that for all $n \geq N$, $|\Upsilon_{\mathbb{Z}}^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2$ for all $v \in \mathbb{Z}$, where $\Upsilon_{\mathbb{Z}}$ denotes the action associated to the graph $\mathcal{G}_{\mathbb{Z}}$. We will show that for $n \geq N$ we also have $|\Upsilon_{\mathcal{G}}^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2$ for all $v \in \mathcal{V}_{\mathcal{G}}$.

Let $\langle s_i \rangle$ denote the sign sequence of θ . Without loss of generality, we may assume $s_0 = ++$, $g_1 \in \{h^{-1}, v^{-1}\}$, and $\gamma(g_1) \in \{h, v\}$. In particular, $\rho_{\lambda}^{\gamma(g_1)}(\mathcal{Q}_{++}) \subset \mathcal{Q}_{++}$. Then by Corollary 11.3, for any infinite connected graph bipartite $\mathcal{H}$ we have $\Upsilon_{\mathcal{H}}(\hat{\mathcal{Q}}_{++}) \subset \mathcal{Q}_{s'_i}$ where $s'_i = \Sigma^{\gamma(g_n)}(++)$. Choose any $v \in \mathcal{V}$. By Lemma 12.5, there is a $\tilde{\mathcal{G}}$ which is either $\mathcal{G}$ or a double cover of $\mathcal{G}$, and an embedding $\tilde{\phi} : \mathcal{G}_{\mathbb{Z}} \rightarrow \tilde{\mathcal{G}}$ such that $\tilde{\phi}(0)$ is a lift of v . Let $\pi : \tilde{\mathcal{G}} \rightarrow \mathcal{G}$ denote this covering, and $\pi_* : \mathbb{R}^{\mathcal{V}_{\tilde{\mathcal{G}}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ be as in equation 12.2. Then by Proposition 12.4 for $n \geq N$, we have

$$|\Upsilon_{\mathcal{G}}^{\gamma(g_n)}(\mathbf{e}_v)(v)| = |\Upsilon_{\tilde{\mathcal{G}}}^{\gamma(g_n)}(\mathbf{e}_{\phi(0)})(\phi(0))|.$$

Then by Theorem 12.3, we know that

$$|\Upsilon_{\tilde{\mathcal{G}}}^{\gamma(g_n)}(\mathbf{e}_{\phi(0)})(\phi(0))| \geq |\Upsilon_{\mathbb{Z}}^{\gamma(g_n)}(\mathbf{e}_0)(0)| \geq 2,$$

so that $|\Upsilon_{\mathcal{G}}^{\gamma(g_n)}(\mathbf{e}_v)(v)| \geq 2$ as desired.

We complete the proof by following the logic applied to the case of $\mathcal{G} = \mathbb{Z}$. By Corollary 10.9, there is a critical time $t \geq N$. Then, for any θ -survivor $\mathbf{f} \in \mathbb{R}^{\mathcal{V}_G}$ and any $v \in \mathcal{V}_G$,

$$\mathbf{f}(v) = \langle \mathbf{f}, \mathbf{e}_v \rangle = \langle \Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{e}_v) \rangle \geq (\Upsilon^{g_t}(\mathbf{f})(v)) (\Upsilon^{\gamma(g_t)}(\mathbf{e}_v)(v)) \geq 2\Upsilon^{g_t}(\mathbf{f})(v).$$

Here the first inequality follows because both $\Upsilon^{g_t}(\mathbf{f}), \Upsilon^{\gamma(g_t)}(\mathbf{e}_v) \in \mathcal{Q}_{s_t}$ by Proposition 11.12. See equation 14.7 for another example of this. (In this case, we have no absolute values because $s_0 = ++$.) We deduce $\Upsilon^{g_t}(\mathbf{f})(v) \leq \frac{1}{2}\mathbf{f}(v)$ as claimed by the lemma. $\square$

14.5. A perturbed group action. Let $\mathbf{Ax} = \mathbf{y}$. For any $g \in G$, we can compute $\Upsilon^g(\mathbf{x}) - \mathbf{x}$ inductively in terms of $\mathbf{y}$. In this section we will explain how this is done.

Fixing any $\mathbf{y} \in \mathbb{R}_c^{\mathcal{V}}$, define the following “affine” actions on $\mathbb{R}_c^{\mathcal{V}}$.

$$\mathbf{H}_{\mathbf{y}}(\mathbf{z}) = \mathbf{H}(\mathbf{z}) + \pi_{\mathcal{A}}(\mathbf{y}) \quad \text{and} \quad \mathbf{V}_{\mathbf{y}}(\mathbf{z}) = \mathbf{V}(\mathbf{z}) + \pi_{\mathcal{B}}(\mathbf{y}).$$

Powers of these functions are given by

$$\mathbf{H}_{\mathbf{y}}^k(\mathbf{z}) = \mathbf{H}^k(\mathbf{z}) + k\pi_{\mathcal{A}}(\mathbf{y}) \quad \text{and} \quad \mathbf{V}_{\mathbf{y}}^k(\mathbf{z}) = \mathbf{V}^k(\mathbf{z}) + k\pi_{\mathcal{B}}(\mathbf{y}).$$

Thus, they generate a nonlinear action $\mathcal{X}_{\mathbf{y}} : G \times \mathbb{R}_c^{\mathcal{V}} \rightarrow \mathbb{R}_c^{\mathcal{V}}$, defined by $\mathcal{X}_{\mathbf{y}}^{h^k} = \mathbf{H}_{\mathbf{y}}^k$ and $\mathcal{X}_{\mathbf{y}}^{v^k} = \mathbf{V}_{\mathbf{y}}^k$.

Proposition 14.18. *For all $g \in G$, $c \in \mathbb{R}$ and $\mathbf{y}, \mathbf{y}_1, \mathbf{y}_2, \mathbf{z}_1, \mathbf{z}_2 \in \mathbb{R}_c^{\mathcal{V}}$, we have*

- (1) $\mathcal{X}_{\mathbf{y}}^g(\mathbf{z}_1 + \mathbf{z}_2) = \mathcal{X}_{\mathbf{y}}^g(\mathbf{z}_1) + \Upsilon^g(\mathbf{z}_2)$.
- (2) $\mathcal{X}_{\mathbf{y}_1 + \mathbf{y}_2}^g(\mathbf{0}) = \mathcal{X}_{\mathbf{y}_1}^g(\mathbf{0}) + \mathcal{X}_{\mathbf{y}_2}^g(\mathbf{0})$.
- (3) $\mathcal{X}_{c\mathbf{y}}^g(\mathbf{0}) = c\mathcal{X}_{\mathbf{y}}^g(\mathbf{0})$.

Proof. All statements follow from induction on the word length of g . The statements are clearly true when g is the identity. Assume statement (1) is true for g_0 . We will show it is true for $h^k g_0$.

$$\begin{aligned} \mathcal{X}_{\mathbf{y}}^{h^k g_0}(\mathbf{z}_1 + \mathbf{z}_2) &= \mathcal{X}_{\mathbf{y}}^{h^k}(\mathcal{X}_{\mathbf{y}}^{g_0}(\mathbf{z}_1) + \Upsilon^{g_0}(\mathbf{z}_2)) = \mathbf{H}^k(\mathcal{X}_{\mathbf{y}}^{g_0}(\mathbf{z}_1) + \Upsilon^{g_0}(\mathbf{z}_2)) + k\pi_{\mathcal{A}}(\mathbf{y}) \\ &= \mathbf{H}^k(\mathcal{X}_{\mathbf{y}}^{g_0}(\mathbf{z}_1)) + k\pi_{\mathcal{A}}(\mathbf{y}) + \Upsilon^{h^k g_0}(\mathbf{z}_2) = \mathcal{X}_{\mathbf{y}}^{h^k g_0}(\mathbf{z}_1) + \Upsilon^{h^k g_0}(\mathbf{z}_2). \end{aligned}$$

A similar equation holds for $v^k g_0$. Now assume (2) is true for g_0 . Then,

$$\begin{aligned} \mathcal{X}_{\mathbf{y}_1 + \mathbf{y}_2}^{h^k g_0}(\mathbf{0}) &= \mathbf{H}^k(\mathcal{X}_{\mathbf{y}_1 + \mathbf{y}_2}^{g_0}(\mathbf{0})) + k\pi_{\mathcal{A}}(\mathbf{y}_1 + \mathbf{y}_2) \\ &= \mathbf{H}^k(\mathcal{X}_{\mathbf{y}_1}^{g_0}(\mathbf{0}) + \mathcal{X}_{\mathbf{y}_2}^{g_0}(\mathbf{0})) + k\pi_{\mathcal{A}}(\mathbf{y}_1) + k\pi_{\mathcal{A}}(\mathbf{y}_2) = \mathcal{X}_{\mathbf{y}_1}^{h^k g_0}(\mathbf{0}) + \mathcal{X}_{\mathbf{y}_2}^{h^k g_0}(\mathbf{0}). \end{aligned}$$

Again, a similar formula holds for $v^k g_0$. Statement (3) holds for similar reasons. $\square$

The following proposition connects this group action to the adjacency sign property.

Proposition 14.19. *Let $\mathbf{x} \in \mathbb{R}_c^{\mathcal{V}}$ and set $\mathbf{y} = \mathbf{Ax}$. Then for all $g \in G$,*

$$\Upsilon^g(\mathbf{x}) - \mathbf{x} = \mathcal{X}_{\mathbf{y}}^g(\mathbf{0}).$$

Proof. We may prove this by induction on the word length of g . The statement is clearly true when g is the identity. Now suppose we know the statement for $g_0 \in G$. We must prove the statement holds for $h^k g_0$ and $v^k g_0$ for $k = \pm 1$. We write

$$\begin{aligned} \Upsilon^{h^k g_0}(\mathbf{x}) - \mathbf{x} &= \mathbf{H}^k \circ \Upsilon^{g_0}(\mathbf{x}) - \mathbf{x} = \mathbf{H}^k(\Upsilon^{g_0}(\mathbf{x}) - \mathbf{x}) + \mathbf{H}^k(\mathbf{x}) - \mathbf{x} \\ &= \mathbf{H}^k \circ \mathcal{X}_{\mathbf{y}}^g(\mathbf{0}) + k\pi_{\mathcal{A}} \circ \mathbf{A}(\mathbf{x}) = \mathbf{H}^k \circ \mathcal{X}_{\mathbf{y}}^g(\mathbf{0}) + k\pi_{\mathcal{A}}(\mathbf{y}) = \mathcal{X}_{\mathbf{y}}^{h^k g_0}(\mathbf{0}). \end{aligned}$$

A similar statement holds for the case of $v^k g_0$. $\square$

We will now connect this operation to the adjacency sign property. Recall that for $\mathbf{v} \in \mathcal{V}$ the function $\mathbf{e}_{\mathbf{v}} \in \mathbb{R}_{\mathcal{C}}^{\mathcal{V}}$ is the function that assigns one to $\mathbf{v}$ and assigns $\mathbf{0}$ to all other vertices.

Definition 14.20. Let $\lambda \geq 2$ and suppose $\boldsymbol{\theta} \in \mathcal{R}_{\lambda} \cap \mathcal{Q}_{++}$ has λ -shrinking sequence $\langle g_i \rangle$. Let $s'_i = \Sigma^{\gamma(g_i)}(++)$. For a vertex $\mathbf{v} \in \mathcal{V}$ we say the graph $\mathcal{G}$ has $\mathbf{v}$ -ASP if for all $i \geq 0$, we have $\mathcal{X}_{\mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}$.

Proposition 14.21. Suppose that $\mathcal{G}$ has $\mathbf{v}$ -ASP for all $\mathbf{v} \in \mathcal{V}$. Then $\mathcal{G}$ has the adjacency sign property.

Proof. Let $\mathbf{x} \in \mathbb{R}_{\mathcal{C}}^{\mathcal{V}}$ and assume that $\mathbf{y} = \mathbf{A}(\mathbf{x}) \in \hat{\mathcal{Q}}_{++}$. Let t be a critical time. Then Proposition 14.19 implies that $\Upsilon^{\gamma(g_t)}(\mathbf{x}) - \mathbf{x} = \mathcal{X}_{\mathbf{y}}^{\gamma(g_t)}(\mathbf{0})$. By Proposition 14.18, we can write

$$(14.10) \quad \mathcal{X}_{\mathbf{y}}^{\gamma(g_t)}(\mathbf{0}) = \sum_{\mathbf{v} \in \mathcal{V}} \mathcal{X}_{\mathbf{y}(\mathbf{v})\mathbf{e}_{\mathbf{v}}}^{\gamma(g_t)}(\mathbf{0}) = \sum_{\mathbf{v} \in \mathcal{V}} \mathbf{y}(\mathbf{v}) \mathcal{X}_{\mathbf{e}_{\mathbf{v}}}^{\gamma(g_t)}(\mathbf{0}).$$

Note that each $\mathbf{y}(\mathbf{v}) \geq 0$ by the assumption that $\mathbf{y} \in \hat{\mathcal{Q}}_{++}$. In addition, each $\mathcal{X}_{\mathbf{e}_{\mathbf{v}}}^{\gamma(g_t)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_t}$ because $\mathcal{G}$ has $\mathbf{v}$ -ASP. Now let $\langle s_i \rangle$ denote the sign sequence of $\boldsymbol{\theta}$, and let $\mathbf{f} \in \mathcal{S}_{\boldsymbol{\theta}}$ be a $\boldsymbol{\theta}$ -survivor. By Proposition 6.20, we know $\Upsilon^{g_t}(\mathbf{f}) \in \hat{\mathcal{Q}}_{s_t}$. By Proposition 11.12, for all critical times t we have $s'_t = s_t$. Therefore, each $\mathcal{X}_{\mathbf{e}_{\mathbf{v}}}^{\gamma(g_t)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s_t}$. Each term in equation 14.10 lies in $\hat{\mathcal{Q}}_{s_t}$. It follows that $\langle \Upsilon^{g_t}(\mathbf{f}), \mathcal{X}_{\mathbf{y}}^{\gamma(g_t)}(\mathbf{0}) \rangle \geq 0$ as desired. $\square$

Lemma 14.22 (Subgraphs and $\mathbf{v}$ -ASP). Let $\mathbf{v} \in \mathcal{V}$, and assume that there is an infinite connected subgraph $\mathcal{H} \subset \mathcal{G}$ containing the vertex $\mathbf{v}$ such that $\mathcal{H}$ has $\mathbf{v}$ -ASP. Then, $\mathcal{G}$ also has $\mathbf{v}$ -ASP.

Proof. We recall our notation from section 12. We distinguish the actions of $\Upsilon_{\mathcal{G}} : G \times \mathbb{R}^{\mathcal{V}_{\mathcal{G}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ and $\Upsilon_{\mathcal{H}} : G \times \mathbb{R}^{\mathcal{V}_{\mathcal{H}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$. To distinguish the two $\mathcal{X}_{\mathbf{y}}$ actions, we use $\mathcal{X}_{\mathcal{G}, \mathbf{y}} : G \times \mathbb{R}^{\mathcal{V}_{\mathcal{G}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ and $\Upsilon_{\mathcal{H}, \mathbf{y}} : G \times \mathbb{R}^{\mathcal{V}_{\mathcal{H}}} \rightarrow \mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$. We will abuse notation by identifying $\mathbb{R}^{\mathcal{V}_{\mathcal{H}}}$ with the subset of $\mathbb{R}^{\mathcal{V}_{\mathcal{G}}}$ which is supported on $\mathcal{V}_{\mathcal{H}}$.

We must show that $\mathcal{X}_{\mathcal{G}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}$ for all i . The statement of the proposition guarantees $\mathcal{X}_{\mathcal{H}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}$ for all i . We will show that for all i

$$(14.11) \quad \mathcal{X}_{\mathcal{G}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) - \mathcal{X}_{\mathcal{H}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}.$$

This implies the proposition.

Now we will inductively define elements $\mathbf{y}_i \in \mathbb{R}_{\mathcal{C}}^{\mathcal{V}_{\mathcal{G}}}$ for each integer $i \geq 1$. Assuming $\mathbf{y}_1, \dots, \mathbf{y}_{i-1}$ are defined, we define $\mathbf{y}_i$ to be the unique element of so that the following equation holds

$$(14.12) \quad \mathcal{X}_{\mathcal{G}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) - \mathcal{X}_{\mathcal{H}, \mathbf{e}_{\mathbf{v}}}^{\gamma(g_i)}(\mathbf{0}) = \Upsilon_{\mathcal{G}}^{\gamma(g_i g_1^{-1})}(\mathbf{y}_1) + \Upsilon_{\mathcal{G}}^{\gamma(g_i g_2^{-1})}(\mathbf{y}_2) + \dots + \mathbf{y}_i.$$

We will show that

$$(14.13) \quad \Upsilon_{\mathcal{G}}^{\gamma(g_i g_j^{-1})}(\mathbf{y}_j) \in \hat{\mathcal{Q}}_{s'_i}. \quad \text{for all } i \geq j.$$

This implies that equation 14.11 holds, because it holds for each term in the sum given in equation 14.12.

We observe by combining the cases $i = j$ and $i = j - 1$ of equation 14.12 that

$$\mathbf{y}_j = \mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_j)}(\mathbf{0}) - \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_j)}(\mathbf{0}) - \Upsilon_{\mathcal{G}}^{\gamma(g_j g_{j-1}^{-1})}(\mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}) - \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0})).$$

Define the following two quantities.

$$\mathbf{a}_j = \mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_j)}(\mathbf{0}) - \Upsilon_{\mathcal{G}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}).$$

$$\mathbf{b}_j = -\mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_j)}(\mathbf{0}) + \Upsilon_{\mathcal{H}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}).$$

Observe that

$$\mathbf{y}_j = \mathbf{a}_j + \mathbf{b}_j + \Upsilon_{\mathcal{G}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}) - \Upsilon_{\mathcal{H}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}).$$

By Proposition 14.18, we have

$$\mathbf{a}_j = \mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_j g_{j-1}^{-1})}(\mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0})) + \Upsilon_{\mathcal{G}}^{\gamma(g_j g_{j-1}^{-1})}(-\mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0})) = \mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_j g_{j-1}^{-1})}(\mathbf{0}).$$

Similarly,

$$\mathbf{b}_j = -\mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_j g_{j-1}^{-1})}(\mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0})) - \Upsilon_{\mathcal{H}}^{\gamma(g_j g_{j-1}^{-1})}(-\mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0})) = -\mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_j g_{j-1}^{-1})}(\mathbf{0}).$$

Thus $\mathbf{a}_j + \mathbf{b}_j = \mathbf{0}$, because $\mathcal{X}_{\mathcal{G}, \mathbf{e}_v}^{\gamma(g_j g_{j-1}^{-1})}(\mathbf{0})$ equals $\pm \pi_{\mathcal{A}}(\mathbf{e}_v)$ when $g_j g_{j-1}^{-1} = h^{\pm 1}$ and equals $\pm \pi_{\mathcal{B}}(\mathbf{e}_v)$ when $g_j g_{j-1}^{-1} = v^{\pm 1}$. Thus,

$$\mathbf{y}_j = \Upsilon_{\mathcal{G}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}) - \Upsilon_{\mathcal{H}}^{\gamma(g_j g_{j-1}^{-1})} \circ \mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}).$$

Now we will use the assumption that $\mathcal{H}$ has v-ASP. Therefore, $\mathcal{X}_{\mathcal{H}, \mathbf{e}_v}^{\gamma(g_{j-1})}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_j}$. Then, by applying Corollary 12.2 to the geodesic ray $\langle \gamma(g_i g_{j-1}^{-1}) \rangle_{i \geq j-1}$ we see that

$$\Upsilon_{\mathcal{G}}^{\gamma(g_i g_j^{-1})}(\mathbf{y}_j) \in \hat{\mathcal{Q}}_{s(i, j)},$$

where $s(i, j) = \Sigma^{\gamma(g_i g_{j-1}^{-1})}(s'_{j-1})$. We observe $s(i, j) = s'_i$, and therefore we have proved equation 14.13 as desired. $\square$

Lemma 14.23 (Covers and v-ASP). *Let $v \in \mathcal{V}$, and assume that there is a finite cover $\tilde{\mathcal{G}}$ of $\mathcal{G}$ and a lift $\tilde{v}$ of v for so that $\tilde{\mathcal{G}}$ has $\tilde{v}$ -ASP. Then, $\mathcal{G}$ has v-ASP.*

Recall our notation for dealing with covers given in section 12. When discussing a covering graph $\tilde{G}$ of $\mathcal{G}$, we will use tildes to denote functions in $\mathbb{R}^{\tilde{\mathcal{V}}}$ and operators on this space. For instance, $\tilde{\mathcal{X}}_{\tilde{\mathbf{y}}}$ denotes an action of G on $\mathbb{R}^{\tilde{\mathcal{V}}}$. Also recall that $\pi : \tilde{\mathcal{G}} \rightarrow \mathcal{G}$ denotes the covering map, and $\pi_* : \mathbb{R}^{\tilde{\mathcal{V}}} \rightarrow \mathbb{R}^{\mathcal{V}}$ is the induced map given in equation 12.2.

The key to the lemma is the following observation.

Proposition 14.24. *Let $\tilde{\mathcal{G}}$ be a finite cover of $\mathcal{G}$. Let $\tilde{\mathbf{x}}, \tilde{\mathbf{y}} \in \mathbb{R}^{\tilde{\mathcal{V}}}$. Then*

- (1) $\pi_* \circ \tilde{\pi}_{\tilde{\mathcal{A}}}(\tilde{\mathbf{y}}) = \pi_{\mathcal{A}} \circ \pi_*(\tilde{\mathbf{y}})$,
- (2) $\pi_* \circ \tilde{\pi}_{\tilde{\mathcal{B}}}(\tilde{\mathbf{y}}) = \pi_{\mathcal{B}} \circ \pi_*(\tilde{\mathbf{y}})$, and

(3) for all $g \in G$, $\pi_* \circ \tilde{\mathcal{X}}_{\tilde{\mathbf{y}}}^g(\tilde{\mathbf{x}}) = \mathcal{X}_{\pi_*(\tilde{\mathbf{y}})}^g \circ \pi_*(\tilde{\mathbf{x}})$.

Proof. The first two statements follow trivially from the definitions. It is enough to check statement (3) on powers of the generators. We will consider the case of $g = h^k$. The case of $g = v^k$ follows similarly. By definition of $\tilde{\mathcal{X}}^{h^k}$, by statement (1) and by Proposition 12.4 we have

$$\pi_* \circ \tilde{\mathcal{X}}_{\tilde{\mathbf{y}}}^{h^k}(\tilde{\mathbf{x}}) = \pi_*(\tilde{\Upsilon}^{h^k}(\tilde{\mathbf{x}}) + k\tilde{\pi}_{\mathcal{A}}(\tilde{\mathbf{y}})) = \Upsilon^{h^k} \circ \pi_*(\tilde{\mathbf{x}}) + k\pi_{\mathcal{A}} \circ \pi_*(\tilde{\mathbf{y}})$$

So, by definition of $\mathcal{X}^{h^k}$, we see $\pi_* \circ \tilde{\mathcal{X}}_{\tilde{\mathbf{y}}}^{h^k}(\tilde{\mathbf{x}}) = \mathcal{X}_{\pi_*(\tilde{\mathbf{y}})}^{h^k} \circ \pi_*(\tilde{\mathbf{x}})$ as desired. $\square$

Proof of Lemma 14.23. First note that $\pi_*(\mathbf{e}_{\tilde{\mathbf{v}}}) = \mathbf{e}_v$. Since $\tilde{\mathcal{G}}$ has $\tilde{\mathbf{v}}$ -ASP, we know $\tilde{\mathcal{X}}_{\tilde{\mathbf{e}}_{\tilde{\mathbf{v}}}}^{\gamma(g_i)}(\tilde{\mathbf{0}}) \in \hat{\mathcal{Q}}_{s'_i}$ for all i . By Proposition 14.24,

$$\mathcal{X}_{\mathbf{e}_v}^{\gamma(g_i)}(\mathbf{0}) = \pi_* \circ \tilde{\mathcal{X}}_{\tilde{\mathbf{e}}_{\tilde{\mathbf{v}}}}^{\gamma(g_i)}(\tilde{\mathbf{0}}),$$

and therefore $\mathcal{X}_{\mathbf{e}_v}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}$ for all i . $\square$

14.6. The adjacency sign property for $\mathbb{Z}$. In this subsection, we complete the proof that graphs with no vertices of valance one have the adjacency sign property.

Theorem 14.25. *The graph $\mathcal{G}_{\mathbb{Z}}$ has $\mathbf{v}$ -ASP for all $\mathbf{v} \in \mathbb{Z}$.*

From the previous subsection, we have the following corollary.

Corollary 14.26. *Suppose $\mathcal{G}$ be any infinite graph without vertices of valance one. Then $\mathcal{G}$ has $\mathbf{v}$ -ASP for all $\mathbf{v} \in \mathcal{V}$. Thus, $\mathcal{G}$ has the adjacency sign property.*

Proof. Lemma 12.5 guarantees that given any $\mathbf{v} \in \mathcal{V}$, we can find an embedding of $\mathcal{G}_{\mathbb{Z}}$ into $\mathcal{G}$ such that $\mathbf{v}$ lies in the image, or we can find a double cover $\tilde{\mathcal{G}}$ of $\mathcal{G}$ and a lift $\tilde{\mathbf{v}}$ of $\mathbf{v}$ and a embedding of $\mathcal{G}_{\mathbb{Z}}$ into $\tilde{\mathcal{G}}$ which passes through $\tilde{\mathbf{v}}$. In the first case, $\mathcal{G}$ has $\mathbf{v}$ -ASP by Lemma 14.22. In the second, $\tilde{\mathcal{G}}$ has $\tilde{\mathbf{v}}$ -ASP by Lemma 14.22 and $\mathcal{G}$ has $\mathbf{v}$ -ASP by Lemma 14.23. The adjacency sign property follows from Proposition 14.21. $\square$

Proof of Theorem 14.25. Applying dihedral symmetries we may assume that $\mathbf{v} \in \mathcal{A}$. By further applying translational symmetries of the graph, we may assume that $\mathbf{v} = 0$. Thus, $\mathcal{A}$ consists of the even integers and $\mathcal{B}$ consists of the odd integers. To simplify notation let $\mathbf{y} = \mathbf{e}_0$.

We now recall what we must prove. Let $\lambda \geq 2$ and suppose $\boldsymbol{\theta} \in \mathcal{R}_{\lambda} \cap \mathcal{Q}_{++}$ has shrinking sequence $\langle g_i \rangle$ and sign sequence $\langle s_i \rangle$. Let $s'_i = \Sigma^{\gamma(g_i)}(++)$. Then, we must show that for all $i \geq 0$, we have $\mathcal{X}_{\mathbf{y}}^{\gamma(g_i)}(\mathbf{0}) \in \hat{\mathcal{Q}}_{s'_i}$.

This is lemma is a combinatorial statement, however we will find it useful to use some geometric tricks. Therefore, we define $\mathbf{1} \in \mathbb{R}^{\mathbb{Z}}$ to be the function such that $\mathbf{1}(x) = 1$ for all $x \in \mathbb{Z}$. This is an eigenvector for the adjacency operator. Namely, $\mathbf{A}(\mathbf{1}) = 2 \cdot \mathbf{1}$. We can therefore build a surface $S_1 = S(\mathcal{G}_{\mathbb{Z}}, \mathbf{1})$. We will find it useful to compute the holonomies of homology classes on this surface. We use

$$\text{hol}_1 : H_1(S_1, V; \mathbb{Z}) \rightarrow \mathbb{R}^2$$

to denote the holonomy map for this surface.

The idea of the proof is to find $\mathbf{z}_i, \mathbf{w}_i \in \mathbb{R}_c^{\mathcal{V}} \cap \hat{\mathcal{Q}}_{s'_i}$ for integers $i \geq 0$ such that the following statements hold.




FIGURE 7. Diagram for the proof of Theorem 14.25. Arrows labeled $\gamma(g_i g_{i-1}^{-1})$ join $\mathbf{z}_{i-1}$ to $\mathbf{z}_i$.

- (1) For all $i \geq 0$ we have $\mathcal{X}_y^{\gamma(g_i)}(\mathbf{0}) = \mathbf{z}_i + \mathbf{w}_i$. By Proposition 14.18, this ensures that for all $k \geq 0$

$$\mathcal{X}_y^{\gamma(g_{i+k})}(\mathbf{0}) = \mathcal{X}_y^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{z}_i) + \Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{w}_i).$$

- (2) $\mathbf{z}_i \in \hat{Q}_{s'_i}$ for all $i \geq 0$.

- (3) For all i and all $k \geq 0$ we have $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{w}_i) \in \hat{Q}_{s'_{i+k}}$. In particular, $\mathbf{w}_i \in \hat{Q}_{s'_i}$ for all i .

These statements guarantee $\mathcal{X}_y^{\gamma(g_i)}(\mathbf{0}) \in \hat{Q}_{s'_i}$ for all i as desired. Therefore, they imply the lemma.

We will now explain our choice of $\mathbf{z}_i$. Our choice determined inductively. We set $\mathbf{z}_0 = \mathbf{0}$. Note that $\mathbf{z}_0 \in \hat{Q}_{s'_0} = \hat{Q}_{++}$. Subsequent choices are determined by following the arrows in Figure 7. We begin at the node labeled $\mathbf{0}$ in the quadrant $++$ of the diagram. Then to find $\mathbf{z}_1$ we follow the arrow labeled $\gamma(g_1)$. To find $\mathbf{z}_i$ we follow the arrows labeled $\gamma(g_1)$, then $\gamma(g_2 g_1^{-1})$, then $\gamma(g_3 g_2^{-1})$, and continue until we follow the arrow labeled $\gamma(g_i g_{i-1}^{-1})$. By comparison with the diagram in Definition 10.1 of the expanding sign action, we see that $\mathbf{z}_i \in \hat{Q}_{s'_i}$ for all i . This verifies statement (2) above.

The formula in statement (1) above inductively determines $\mathbf{w}_i$ from $\mathbf{z}_i$. We have $\mathbf{w}_0 = \mathbf{0}$. For $i \geq 1$ we have

$$\begin{aligned} \mathbf{w}_i &= \mathcal{X}_y^{\gamma(g_i)}(\mathbf{0}) - \mathbf{z}_i = \mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})} \circ \mathcal{X}_y^{\gamma(g_{i-1})}(\mathbf{0}) - \mathbf{z}_i \\ (14.14) \quad &= \mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{z}_{i-1} + \mathbf{w}_{i-1}) - \mathbf{z}_i \\ &= \mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{z}_{i-1}) - \mathbf{z}_i + \Upsilon^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{w}_{i-1}). \end{aligned}$$

Because of choice of $\mathbf{w}_i$, we have automatically verified statement (1) above.

It remains to verify statement (3). We prove this statement by induction in i . Clearly, statement (3) is true for $\mathbf{w}_0 = \mathbf{0}$ for all k . Now assume the statement is true for $i-1$.

In particular, we assume that $\Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})}(\mathbf{w}_{i-1}) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$. We wish to show statement (3) holds for i . By equation 14.14, we have

$$\Upsilon^{\gamma(g_{i+k}g_i^{-1})}(\mathbf{w}_i) = \Upsilon^{\gamma(g_{i+k}g_i^{-1})}(\mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{z}_{i-1}) - \mathbf{z}_i) + \Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})}(\mathbf{w}_{i-1}).$$

By our inductive hypothesis, it is sufficient to show that for all $k \geq 0$,

$$(14.15) \quad \Upsilon^{\gamma(g_{i+k}g_i^{-1})}(\mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{z}_{i-1}) - \mathbf{z}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}.$$

Despite the fact that we need equation 14.15 for all $k \geq 0$, this is really a finite check. We will use the Quadrant Tracking Proposition 11.4 to verify this for all k in one step. The number of checks is therefore equal to the number (30) of arrows in figure 7 above.

In order to cut down on the number of checks, we consider the element of the dihedral group action on the plane $\Delta : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $\Delta(x, y) = (-x, y)$. Note that it satisfies the equations

$$\Delta \circ \rho_\lambda^h = \rho_\lambda^{h^{-1}} \circ \Delta \quad \text{and} \quad \Delta \circ \rho_\lambda^v = \rho_\lambda^{v^{-1}} \circ \Delta.$$

We define $\delta : G \rightarrow G$ to be the automorphism induced by the action on generators $\delta(h) = h^{-1}$ and $\delta(v) = v^{-1}$. The action satisfies $\Delta \circ \rho_\lambda^g = \rho_\lambda^{\delta(g)} \circ \Delta$ for all $g \in G$. There is an orientation reversing affine automorphism of the surface S_1 which preserves each rectangle which appears as an intersection of a horizontal and vertical cylinder and which has derivative given by Δ . This affine automorphism preserves $\Xi(\mathbb{R}^\mathbb{Z})$. The action therefore lifts to an action on $\mathbb{R}^\mathbb{Z}$. This action $\Delta_* : \mathbb{R}^\mathbb{Z} \rightarrow \mathbb{R}^\mathbb{Z}$ is given by

$$\Delta_*(\mathbf{f})(v) = \begin{cases} -\mathbf{f}(v) & \text{if } v \in \mathcal{A} \\ \mathbf{f}(v) & \text{if } v \in \mathcal{B}. \end{cases}$$

We can use this action to conjugate our group of operators Υ^G .

$$\Upsilon^g \circ \Delta_* = \Delta_* \circ \Upsilon^{\delta(g)} \quad \text{for all } g \in G.$$

For our specific choice of $\mathbf{y} = \mathbf{e}_0$ (or more generally for any $\mathbf{y}$ supported on $\mathcal{A}$) we have

$$\Delta_* \circ \mathbf{H}_y = \mathbf{H}_y^{-1} \circ \Delta_* \quad \text{and} \quad \Delta_* \circ \mathbf{V}_y = \mathbf{V}_y^{-1} \circ \Delta_*.$$

Thus for all $g \in G$, we have $\mathcal{X}_y^g \circ \Delta_* = \Delta_* \circ \mathcal{X}_y^{\delta(g)}$. In particular, we can apply Δ_* to both sides of equation 14.15 to obtain

$$\Upsilon^{\delta \circ \gamma(g_{i+k}g_i^{-1})}(\mathcal{X}_y^{\delta \circ \gamma(g_i g_{i-1}^{-1})} \circ \Delta_*(\mathbf{z}_{i-1}) - \Delta_*(\mathbf{z}_i)) \in \hat{\mathcal{Q}}_{\Delta(s'_{i+k})}.$$

Therefore by the invariance of figure 7 under Δ_* , we only need to consider the case when $\mathbf{z}_{i-1} \in \hat{\mathcal{Q}}_{++} \cup \mathcal{Q}_{+-}$. We will show that equation 14.15 holds in each of these cases below.

For all i let $\mathbf{x}_i = \mathcal{X}_y^{\gamma(g_i g_{i-1}^{-1})}(\mathbf{z}_{i-1}) - \mathbf{z}_i$.

(1) Suppose $\mathbf{z}_{i-1} = \mathbf{0} \in \hat{\mathcal{Q}}_{++}$.

(a) If $\gamma(g_i g_{i-1}^{-1}) = h$, then $\mathbf{z}_i = \mathbf{e}_0 \in \hat{\mathcal{Q}}_{++}$. We compute

$$\mathbf{x}_i = \mathbf{H}_y(\mathbf{0}) - \mathbf{e}_0 = \mathbf{0}.$$

Thus, in this case $\Upsilon^{\gamma(g_{i+k}g_i^{-1})}(\mathbf{x}_i) = \mathbf{0}$ for all k . The conclusion follows trivially, because $\mathbf{0} \in \hat{\mathcal{Q}}_s$ for all $s \in SP$.

(b) If $\gamma(g_i g_{i-1}^{-1}) = v$, then $\mathbf{z}_i = \mathbf{0} \in \hat{\mathcal{Q}}_{++}$. We compute $\mathbf{x}_i = \mathbf{0}$. The conclusion follows as in case (1a).

- (c) If $\gamma(g_i g_{i-1}^{-1}) = h^{-1}$, then $\mathbf{z}_i = -\mathbf{e}_0 \in \hat{\mathcal{Q}}_{-+}$. We compute $\mathbf{x}_i = \mathbf{0}$. The conclusion follows as in case (1a).
- (2) Suppose $\mathbf{z}_{i-1} = \mathbf{e}_0 \in \hat{\mathcal{Q}}_{++}$.
- (a) If $\gamma(g_i g_{i-1}^{-1}) = h$, then $\mathbf{z}_i = \mathbf{e}_0 \in \hat{\mathcal{Q}}_{++}$. We compute

$$\mathbf{x}_i = \mathbf{H}_y(\mathbf{z}_{i-1}) - \mathbf{z}_i = 2\mathbf{e}_0 - \mathbf{e}_0 = \mathbf{e}_0.$$

We see that $\mathbf{x}_i = \mathbf{H}(\mathbf{e}_0) \in \hat{\mathcal{Q}}_{++}$. We have $\mathbf{x}_i, \mathbf{e}_0 \in \mathcal{Q}_{++}$, while $\rho_\lambda^h(\mathcal{Q}_{++}) \subset \mathcal{Q}_{++}$. Thus Proposition 11.4 entails that the quadrant containing $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i)$ is governed by the expanding sign action. Namely, we consider the geodesic ray $\langle \gamma(g_{l+i-1} g_{i-1}^{-1}) \rangle_{l \geq 0}$. Proposition 11.4 implies that $\Upsilon^{\gamma(g_{l+i-1} g_{i-1}^{-1})}(\mathbf{e}_0) \in \hat{\mathcal{Q}}_{\tilde{s}_l}$ where $\tilde{s}_l = \Sigma^{\gamma(g_{l+i-1} g_{i-1}^{-1})}(++)$. Note by induction that $\tilde{s}_{k+1} = s'_{i+k}$ for all $k \geq 0$, since $s'_{i-1} = \tilde{s}_0 = ++$ and $s'_{i+k} = \Sigma^{g_{i+k} g_{i-1}^{-1}}(s'_{i-1}) = \tilde{s}_{k+1}$. Therefore, for all $k \geq 0$,

$$\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) = \Upsilon^{\gamma(g_{i+k} g_{i-1}^{-1})}(\mathbf{e}_0) \in \hat{\mathcal{Q}}_{s'_{i+k}}.$$

- (b) If $\gamma(g_i g_{i-1}^{-1}) = v$, then $\mathbf{z}_i = \mathbf{0} \in \hat{\mathcal{Q}}_{++}$. We compute

$$\mathbf{x}_i = \mathbf{V}_y(\mathbf{z}_{i-1}) - \mathbf{z}_i = \mathbf{e}_{-1} + \mathbf{e}_0 + \mathbf{e}_1.$$

We have $\mathbf{x}_i = \mathbf{V}(\mathbf{e}_0)$ with both $\mathbf{x}_i, \mathbf{e}_0 \in \hat{\mathcal{Q}}_{++}$, while $\rho_\lambda^v(\mathcal{Q}_{++}) \subset \mathcal{Q}_{++}$. Thus, Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$. See case (2a).

- (c) If $\gamma(g_i g_{i-1}^{-1}) = v^{-1}$, then $\mathbf{z}_i = -\mathbf{e}_{-1} + \mathbf{e}_0 - \mathbf{e}_1 \in \hat{\mathcal{Q}}_{-+}$. We compute $\mathbf{x}_i = \mathbf{0}$. The conclusion follows as in case (1a).
- (3) Suppose $\mathbf{z}_{i-1} = -\mathbf{e}_{-1} + \mathbf{e}_0 - \mathbf{e}_1 \in \hat{\mathcal{Q}}_{+-}$.
- (a) If $\gamma(g_i g_{i-1}^{-1}) = h$, then $\mathbf{z}_i = -\mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 - \mathbf{e}_2 \in \hat{\mathcal{Q}}_{--}$. We compute

$$\mathbf{x}_i = \mathbf{H}_y(\mathbf{z}_{i-1}) - \mathbf{z}_i = (-\mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 - \mathbf{e}_2) - \mathbf{z}_i = \mathbf{0}.$$

The conclusion follows as in case (1a).

- (b) If $\gamma(g_i g_{i-1}^{-1}) = h^{-1}$, then $\mathbf{z}_i = \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\mathbf{x}_i = \mathbf{H}_y^{-1}(\mathbf{z}_{i-1}) - \mathbf{z}_i = (\mathbf{e}_{-2} - \mathbf{e}_{-1} + 2\mathbf{e}_0 - \mathbf{e}_1 + \mathbf{e}_2) - \mathbf{z}_i = 2\mathbf{e}_0.$$

Therefore $\mathbf{x}_i = \mathbf{H}^{-1}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{+-}$ while $\rho_\lambda^{h^{-1}}(\mathcal{Q}_{+-}) \subset \mathcal{Q}_{+-}$. Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$. See case (2a).

- (c) If $\gamma(g_i g_{i-1}^{-1}) = v^{-1}$, then $\mathbf{z}_i = -\mathbf{e}_{-1} + \mathbf{e}_0 - \mathbf{e}_1 \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\mathbf{x}_i = \mathbf{V}_y^{-1}(\mathbf{z}_{i-1}) - \mathbf{z}_i = (-2\mathbf{e}_{-1} + \mathbf{e}_0 - 2\mathbf{e}_1) - \mathbf{z}_i = -\mathbf{e}_{-1} - \mathbf{e}_1.$$

Therefore $\mathbf{x}_i = \mathbf{V}^{-1}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{+-}$. Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$ as in case (2a).

- (4) Suppose $\mathbf{z}_{i-1} = \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 \in \hat{\mathcal{Q}}_{+-}$.

- (a) If $\gamma(g_i g_{i-1}^{-1}) = v$, then $\mathbf{z}_i = \mathbf{0} \in \hat{\mathcal{Q}}_{++}$. We compute

$$\mathbf{x}_i = \mathbf{V}_y(\mathbf{z}_{i-1}) = \mathbf{V}(\mathbf{z}_{i-1}) = \mathbf{e}_{-3} + \mathbf{e}_{-2} + \mathbf{e}_2 + \mathbf{e}_3.$$

In this case $\mathbf{x}_i = \mathbf{V}(\mathbf{z}_{i-1})$. There exist saddle connections σ_1 and σ_2 on the surface S_1 such that $\mathcal{Z}(\llbracket \sigma_1 \rrbracket) = \mathbf{e}_{-2} - \mathbf{e}_{-1}$ and $\mathcal{Z}(\llbracket \sigma_2 \rrbracket) = -\mathbf{e}_1 + \mathbf{e}_2$. These saddle connections are pictured on the left side of Figure 8. For $j \in \{1, 2\}$, we have




FIGURE 8. Both sides of this figure illustrate the surface $S_1 = S(\mathcal{G}_{\mathbb{Z}}, \mathbf{1})$. To obtain the surface, identify opposite horizontal and vertical edges by vertical and horizontal translations (respectively). This figure illustrates the saddle connections $\sigma_1, \dots, \sigma_5$ used in the proof of Theorem 14.25. From left to right, the dotted arrows represent the saddle connections $\Phi^{h^{-1}}(\sigma_1)$, $\Phi^{h^{-1}}(\sigma_2)$, $\Phi^{v^{-1}}(\sigma_3)$ and $\Phi^{v^{-1}}(\sigma_5)$. Also, $\Phi^{v^{-1}}(\sigma_4) = \sigma_4$.

$\mathbf{V} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket) = \mathcal{Z} \circ \Phi^{h^{-1}}(\llbracket \sigma_j \rrbracket)$ by Corollary 11.1. We note that the length of the holonomy of these saddle connections is

$$\|hol_1 \circ \Phi^{h^{-1}}(\sigma_j)\| = \|hol_1(\sigma_j)\| = \sqrt{2}.$$

Thus, $hol \sigma_j \in cl(Exp_2(h^{-1}) \cap \mathcal{Q}_{--})$. Therefore, by considering the geodesic ray $\langle \gamma(g_{i-1+l}g_{i-1}^{-1}) \rangle_{l \geq 0}$, Corollary 11.3 implies that $\Upsilon^{\gamma(g_{i-1+l}g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket) \in \hat{\mathcal{Q}}_{\tilde{s}_l}$ where $\tilde{s}_l = r \circ \Sigma^{g_{i-1+l}g_{i-1}^{-1}}(--)$. By equation 11.3, we have

$$\tilde{s}_l = \Sigma^{\gamma(g_{i-1+l}g_{i-1}^{-1})} \circ r(--) = \Sigma^{\gamma(g_{i-1+l}g_{i-1}^{-1})}(+-).$$

Thus, we have $s'_{i+k} = \tilde{s}_{k+1}$ for all $k \geq 0$, since $s'_{i-1} = \tilde{s}_0 = +-$ and $s'_{i+k} = \Sigma^{\gamma(g_{i+k}g_{i-1}^{-1})}(s'_{i-1})$ while $\tilde{s}_{k+1} = \Sigma^{\gamma(g_{k+i}g_{i-1}^{-1})}(\tilde{s}_0)$. Therefore, we have

$$\Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})}(\mathbf{x}_i) = \Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_1 \rrbracket) + \Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_2 \rrbracket) \in \hat{\mathcal{Q}}_{s'_{i+k}}$$

for all $k \geq 0$.

(b) If $\gamma(g_i g_{i-1}^{-1}) = h^{-1}$, then $\mathbf{z}_i = \mathbf{z}_{i-1} = \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\mathbf{x}_i = \mathbf{H}_y^{-1}(\mathbf{z}_{i-1}) - \mathbf{z}_i = (2\mathbf{e}_{-2} - \mathbf{e}_{-1} + \mathbf{e}_0 - \mathbf{e}_1 + 2\mathbf{e}_2) - \mathbf{z}_i = \mathbf{e}_{-2} + \mathbf{e}_0 + \mathbf{e}_2.$$

Thus $\mathbf{x}_i = \mathbf{H}^{-1}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{+-}$ while $\rho_{\lambda}^{h^{-1}}(\mathcal{Q}_{+-}) \subset \mathcal{Q}_{+-}$. Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$ as in case (2a).

(c) If $\gamma(g_i g_{i-1}^{-1}) = v^{-1}$, then $\mathbf{z}_i = -\mathbf{e}_{-3} + \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 - \mathbf{e}_3 \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\mathbf{x}_i = (-\mathbf{e}_{-3} + \mathbf{e}_{-2} - 2\mathbf{e}_{-1} - 2\mathbf{e}_1 + \mathbf{e}_2 - \mathbf{e}_3) - \mathbf{z}_i = -\mathbf{e}_{-1} - \mathbf{e}_1.$$

In particular, we have that $\mathbf{x}_i = \mathbf{V}^{-1}(\mathbf{x}_i)$. So, Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k}g_{i-1}^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$ as in case (2a).

(5) Suppose $\mathbf{z}_{i-1} = -\mathbf{e}_{-3} + \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 - \mathbf{e}_3 \in \hat{\mathcal{Q}}_{+-}$.

(a) If $\gamma(g_i g_{i-1}^{-1}) = h$, then $\mathbf{z}_i = -\mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 - \mathbf{e}_2 \in \hat{\mathcal{Q}}_{--}$. We compute

$$\begin{aligned} \mathbf{x}_i &= \mathbf{H}_y(\mathbf{z}_{i-1}) - \mathbf{z}_i \\ &= (-\mathbf{e}_{-4} - \mathbf{e}_{-3} - \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_0 - \mathbf{e}_1 - \mathbf{e}_2 - \mathbf{e}_3 - \mathbf{e}_4) - \mathbf{z}_i \\ &= -\mathbf{e}_{-4} - \mathbf{e}_{-3} - \mathbf{e}_0 - \mathbf{e}_3 - \mathbf{e}_4. \end{aligned}$$

In this case $\mathbf{x}_i = \mathbf{H}(-\mathbf{e}_{-3} + \mathbf{e}_{-2} - \mathbf{e}_0 + \mathbf{e}_2 - \mathbf{e}_3)$. We can find saddle connections σ_3, σ_4 and σ_5 so that $\mathcal{Z}(\llbracket \sigma_3 \rrbracket) = -\mathbf{e}_{-3} + \mathbf{e}_{-2}$, $\mathcal{Z}(\llbracket \sigma_4 \rrbracket) = -\mathbf{e}_0$ and $\mathcal{Z}(\llbracket \sigma_5 \rrbracket) = \mathbf{e}_2 - \mathbf{e}_3$. These saddle connections are depicted on the right side of Figure 8. For $j \in \{3, 4, 5\}$, we have $\mathbf{H} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket) = \mathcal{Z} \circ \Phi^{v^{-1}}(\llbracket \sigma_j \rrbracket)$ by Corollary 11.1. For each such j we have

$$\|hol_1(\sigma_j)\| = \|hol_1 \circ \Phi^{v^{-1}}(\llbracket \sigma_j \rrbracket)\|.$$

In particular, $hol_1(\sigma_j) \in cl(Exp_2(v^{-1}) \cap (\mathcal{Q}_{-+} \cup \mathcal{Q}_{--}))$. Corollary 11.3 implies that the quadrant containing $\Upsilon^{\gamma(g_{i+k} g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket)$ is given by may determined by following the expanding sign action. As in case (4a), we consider the geodesic ray $\langle \gamma(g_{i-1+l} g_{i-1}^{-1}) \rangle_{l \geq 0}$, and $\Upsilon^{\gamma(g_{i+l} g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket) \in \hat{\mathcal{Q}}_{\tilde{s}_l}$ where

$$\tilde{s}_l = r \circ \Sigma^{g_{i+l} g_{i-1}^{-1}}(s) = \Sigma^{\gamma(g_{i+l} g_{i-1}^{-1})} \circ r(s),$$

where $s = --$ or $s = -+$ depending on $j \in \{3, 4, 5\}$. In these cases $r(s) = +-$ or $r(s) = --$. We have $\gamma(g_i g_{i-1}^{-1}) = h$, and

$$\Sigma^h(+-) = \Sigma^h(--) = --.$$

Therefore in either case, we have $\tilde{s}_l = \Sigma^{\gamma(g_{i-1+l} g_{i-1}^{-1})}(+-)$ for $l \geq 1$. We also observe $s'_{i+k} = \tilde{s}_{1+k}$ for all $k \geq 0$. We conclude that for all $k \geq 0$,

$$\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) = \sum_{j=3}^5 \Upsilon^{\gamma(g_{i+k} g_{i-1}^{-1})} \circ \mathcal{Z}(\llbracket \sigma_j \rrbracket) \in \hat{\mathcal{Q}}_{s'_{i+k}}.$$

(b) If $\gamma(g_i g_{i-1}^{-1}) = h^{-1}$, then $\mathbf{z}_i = \mathbf{e}_{-2} - \mathbf{e}_{-1} - \mathbf{e}_1 + \mathbf{e}_2 \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\begin{aligned} \mathbf{x}_i &= \mathbf{H}_y^{-1}(\mathbf{z}_{i-1}) - \mathbf{z}_i \\ &= (\mathbf{e}_{-4} - \mathbf{e}_{-3} + 3\mathbf{e}_{-2} - \mathbf{e}_{-1} + \mathbf{e}_0 - \mathbf{e}_1 + 3\mathbf{e}_2 - \mathbf{e}_3 + \mathbf{e}_4) - \mathbf{z}_i \\ &= \mathbf{e}_{-4} - \mathbf{e}_{-3} + 2\mathbf{e}_{-2} + \mathbf{e}_0 + 2\mathbf{e}_2 - \mathbf{e}_3 + \mathbf{e}_4. \end{aligned}$$

In particular, $\mathbf{x}_i = \mathbf{H}^{-1}(-\mathbf{e}_{-3} + \mathbf{e}_{-2} + \mathbf{e}_0 + \mathbf{e}_2 - \mathbf{e}_3)$. Both $\mathbf{x}_i$ and $-\mathbf{e}_{-3} + \mathbf{e}_{-2} + \mathbf{e}_0 + \mathbf{e}_2 - \mathbf{e}_3$ lie in $\hat{\mathcal{Q}}_{+-}$, while $\rho_\lambda^{h^{-1}}(\mathcal{Q}_{+-}) \subset \mathcal{Q}_{+-}$. Therefore, Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$ as in case (2a).

(c) If $\gamma(g_i g_{i-1}^{-1}) = v^{-1}$, then $\mathbf{z}_i = \mathbf{z}_{i-1} \in \hat{\mathcal{Q}}_{+-}$. We compute

$$\mathbf{x}_i = \mathbf{V}_y^{-1}(\mathbf{z}_{i-1}) - \mathbf{z}_i = -\mathbf{e}_{-3} - \mathbf{e}_{-1} - \mathbf{e}_1 - \mathbf{e}_3.$$

Therefore, $\mathbf{x}_i = \mathbf{V}^{-1}(\mathbf{x}_i)$ while $\rho_\lambda^{v^{-1}}(\mathcal{Q}_{+-}) \subset \mathcal{Q}_{+-}$. So, Proposition 11.4 guarantees that $\Upsilon^{\gamma(g_{i+k} g_i^{-1})}(\mathbf{x}_i) \in \hat{\mathcal{Q}}_{s'_{i+k}}$ for all $k \geq 0$ as in case (2a).

□

15. THE MARTIN BOUNDARIES

15.1. The Martin Boundaries of a graph. In this section, we will briefly review some relevant facts about the Martin boundaries of the adjacency operator of an infinite connected graph $\mathcal{G}$ with bounded valance. We will follow the surveys [MW89] and [Woe00]. The reader is especially encouraged to refer to [MW89], because it specifically discusses the adjacency operator. The book [Woe00] deals exclusively with stochastic matrices. However, our discussion of the adjacency operator can be reduced to the discussion of stochastic matrices; see remark 15.8.

We may view the adjacency operator as an infinite matrix. For $v, w \in \mathcal{V}$ and for n a non-negative integer, we define the non-negative number

$$\mathbf{A}_{v,w}^{(n)} = \langle \mathbf{A}^n(\mathbf{e}_w), \mathbf{e}_v \rangle \in \mathbb{R}$$

where $\mathbf{e}_v$ denote the function $\mathcal{V} \rightarrow \mathbb{R}$ which is one at v and zero elsewhere. This means that we can write

$$\mathbf{A}^n(\mathbf{f}) = \sum_{v \in \mathcal{V}} \sum_{w \in \mathcal{V}} \mathbf{A}_{v,w}^{(n)} \mathbf{f}(w) \mathbf{e}_v.$$

Given a $\lambda > 0$, we define the matrix $\mathbf{R}_\lambda = \frac{1}{\lambda} \sum_{n=0}^{\infty} (\frac{1}{\lambda} \mathbf{A})^n$. For λ small this sum will diverge. But there is a real constant $r > 0$ for which $\mathbf{R}_\lambda$ has all finite entries for all $\lambda > r$, and $\mathbf{R}_\lambda$ has all infinite entries for $\lambda < r$. This constant r coincides with the spectral radius of the action of $\mathbf{A}$ on $\ell_2(\mathcal{V})$.

Note that the matrix $\mathbf{R}_\lambda$ satisfies the equation

$$(15.1) \quad \lambda \mathbf{R}_\lambda = \mathbf{A} \mathbf{R}_\lambda + \mathbf{I}.$$

From this point of view, the columns $\mathbf{R}_\lambda(\mathbf{e}_v)$ for $v \in \mathcal{V}$ are nearly positive eigenfunctions with eigenvalue λ .

We have not explained what happens for $\lambda = r$. $\mathbf{A}$ is called *r-transient* if $\mathbf{R}_r$ has all finite entries, and $\mathbf{A}$ is called *r-recurrent* if $\mathbf{R}_r$ has all infinite entries. No other possibilities can occur. In the *r-recurrent* case, we might try to compute the matrix

$$\mathbf{L} = \lim_{n \rightarrow \infty} \frac{1}{r^n} \mathbf{A}^n.$$

This limit always exists. We say that $\mathbf{A}$ is *r-null* if all entries of $\mathbf{L}$ are zero, and *r-positive* if all entries are non-zero. Again, no other possibilities can occur. In the *r-positive* case, the columns of $\mathbf{L}$ are positive eigenfunctions. Moreover, since $\mathbf{L}^2 = \mathbf{L}$, these eigenfunctions lie in $\ell_2(\mathcal{V})$. It turns out that all columns are multiples of one another. For the following see [MW89, Theorem 6.2].

Theorem 15.1 ([Pru64] and [VJ67]). *If $\mathbf{A}$ is r-recurrent, then there is a positive solution to the equation $\mathbf{A}(\mathbf{f}) = r\mathbf{f}$. This solution is unique up to scaling.*

Definition 15.2 (Martin kernel). Suppose that $\lambda > r$ or that $\lambda = r$ and $\mathbf{A}$ is *r-transient*. Choose a *root* vertex $o \in \mathcal{V}$. The λ -*Martin kernel* is the matrix satisfying

$$\mathbf{K}_\lambda : \mathcal{V}^2 \rightarrow \mathbb{R}; (v, w) \mapsto \frac{\mathbf{R}_\lambda(\mathbf{e}_w)(v)}{\mathbf{R}_\lambda(\mathbf{e}_w)(o)}.$$

We may view $\mathbf{K}_\lambda$ as a modification of the matrix $\mathbf{R}_\lambda$ where all columns have been rescaled so that the row associated to the root consists of all ones. Any non-trivial pointwise limit of the columns $\mathbf{K}_\lambda(\mathbf{e}_w)$ of $\mathbf{K}_\lambda$ produces a positive eigenfunction by equation 15.1.

Definition 15.3 (Martin boundary). The λ -Martin compactification $\widehat{\mathcal{V}}_\lambda$ of the vertex set $\mathcal{V}$ is the smallest compactification of $\mathcal{V}$ to which the function

$$\mathbf{k}_\lambda : \mathcal{V} \rightarrow \mathbb{R}^\mathcal{V}; w \mapsto \mathbf{K}_\lambda(\mathbf{e}_w)$$

extends continuously. The λ -Martin boundary is $\mathcal{M}_\lambda = \widehat{\mathcal{V}}_\lambda \setminus \mathcal{V}$.

It follows from the above discussion that if $\zeta \in \mathcal{M}_\lambda$, we have $\mathbf{A}\mathbf{k}_\lambda(\zeta) = \lambda\mathbf{k}_\lambda(\zeta)$.

Remark 15.4. Sometimes the Martin compactification is defined as the smallest compactification to which the maps which send a vertex to the associated row of $\mathbf{K}_\lambda$ are continuous. In our case, these two definitions are equivalent. See [Woe00, §IV.24].

Definition 15.5 (Minimal Martin boundary). A point $\zeta \in \mathcal{M}_\lambda$ is called *minimal* if whenever $\eta_1, \eta_2 \in \mathcal{M}_\lambda$ and $0 < t < 1$ satisfy

$$t\mathbf{k}_\lambda(\eta_1) + (1-t)\mathbf{k}_\lambda(\eta_2) = \mathbf{k}_\lambda(\zeta),$$

we have $\zeta = \eta_1 = \eta_2$. We use $\mathcal{M}_\lambda^{\min} \subset \mathcal{M}_\lambda$ to denote the set of all minimal $\zeta \in \mathcal{M}_\lambda$. The subset $\mathcal{M}_\lambda^{\min} \subset \mathcal{M}_\lambda$ is a Borel subset.

From the Poisson-Martin Representation Theorem (Theorem 6.45), we have the following.

Corollary 15.6. *The function $\mathbf{k}_\lambda$ restricts to a bijection between the minimal Martin boundary and the extremal eigenfunctions of $\mathbf{A}$ with eigenvalue λ .*

Since an infinite set with the discrete topology can not be compact, we have the following.

Corollary 15.7. *There exists a positive function satisfying $\mathbf{A}(\mathbf{f}) = \lambda\mathbf{f}$ for all $\lambda \geq r$.*

Remark 15.8 (Reduction to the stochastic case [DSW60]). Much of the literature on this subject is concerned with *stochastic matrices* $\mathbf{P}$, which are defined to have the property that $\mathbf{P}(\mathbf{1}) = \mathbf{1}$. In this case, functions $\mathbf{w}$ satisfying $\mathbf{P}(\mathbf{w}) = \mathbf{w}$ are called *harmonic*. Suppose we have a positive function $\mathbf{f} \in \mathbb{R}^\mathcal{V}$ satisfying $\mathbf{A}(\mathbf{f}) = \lambda\mathbf{f}$. Define $\mathbf{D}$ to be diagonal matrix with $\mathbf{f}(v)$ in the diagonal entry associated to $v \in \mathcal{V}$. Consider the matrix $\mathbf{P} = \frac{1}{\lambda}\mathbf{D}^{-1}\mathbf{A}\mathbf{D}$. The matrix $\mathbf{P}$ is easily seen stochastic. Moreover, if $\mathbf{g}$ is another function satisfying $\mathbf{A}(\mathbf{g}) = \lambda\mathbf{g}$, then $\mathbf{P}(\mathbf{D}^{-1}\mathbf{g}) = \mathbf{D}^{-1}\mathbf{g}$. And conversely, if $\mathbf{h}$ satisfies $\mathbf{P}(\mathbf{h}) = \mathbf{h}$ then $\mathbf{A}(\mathbf{D}\mathbf{h}) = \lambda\mathbf{D}\mathbf{h}$. Therefore, $\mathbf{D}$ induces a linear bijection between the harmonic functions of $\mathbf{P}$ and the eigenfunctions of $\mathbf{A}$ with eigenvalue λ . This map also respects the definitions above.

15.2. Hyperbolic graphs. A graph is called *hyperbolic* if it is δ -hyperbolic as a metric space equipped with the edge metric for some $\delta > 0$. We refer the reader to [BBI01, §8.4] for the definition of and background for δ -hyperbolicity. If $\mathcal{G}$ is hyperbolic, we can compactify $\mathcal{G}$ with the δ -hyperbolic boundary $\partial_{\text{hyp}}\mathcal{G}$. For a discussion of the following see [Woe00, Theorem IV.27.1].

Theorem 15.9 ([Anc88]). *Suppose the graph $\mathcal{G}$ is hyperbolic, and let $\lambda > r$. Then, every point in $\mathcal{M}_\lambda$ is minimal and $\mathcal{M}_\lambda$ is homeomorphic to $\partial_{\text{hyp}}\mathcal{G}$.*

Example 15.10. Consider the graph $\mathcal{G}$ given in figure 2. The eigenfunction $\mathbf{w}$ given in the figure lies in $\ell_2(\mathcal{V})$. We conclude that $\mathbf{A}$ is r -positive and that the spectral radius is the associated eigenvalue $r = \frac{3\sqrt{2}}{2}$. By Theorem 15.1, we conclude that the only positive functions satisfying $\mathbf{A}(\mathbf{f}) = r\mathbf{f}$ are the multiples of the function $\mathbf{w}$. For $\lambda > r$, we may apply Theorem 15.9. In the case of a tree, the hyperbolic boundary of $\mathcal{G}$ is homeomorphic to the

space of ends of $\mathcal{G}$. In this case, we have exactly 3 ends. We conclude the space of positive functions satisfying $\mathbf{A}(\mathbf{f}) = \lambda \mathbf{f}$ is linearly isomorphic to the cone on a triangle; we have 3 such extremal eigenfunctions up to scaling. These functions are given in figure 9. The space of all extremal positive eigenfunctions is homeomorphic to a cone on the graph itself.




FIGURE 9. The extremal eigenfunctions of the graph $\mathcal{G}$ of are shown (up to scaling and the symmetries of the graph). Each $t \geq \sqrt{2}$ corresponds to an extremal eigenfunction with eigenvalue $\lambda = t + t^{-1}$ if we set $a = \frac{t-2t^{-1}}{t-t^{-1}}$.

The theorem above leaves open the question of what the Martin boundary is in the case that $\lambda = r$ and $\mathbf{A}$ is r -transient. In the special case of trees, we have an answer.

Theorem 15.11 ([PW88]). *If $\mathcal{G}$ is a tree and $\mathbf{A}$ is r -transient with $\lambda \geq r$ or if $\mathbf{A}$ is r -recurrent and $\lambda > r$, then every point in $\mathcal{M}_\lambda$ is minimal and $\mathcal{M}_\lambda$ is homeomorphic to the space of ends of $\mathcal{G}$ (which is also homeomorphic to $\partial_{\text{hyp}} \mathcal{G}$).*

15.3. Minimal Martin boundaries of Cayley graphs. In this section, we let $\mathcal{G}$ be the Cayley graph of a finitely generated infinite group $\mathbb{G}$. In this case, we have a characterization of r -recurrence.

Theorem 15.12 ([Var86]). *The adjacency operator on a Cayley graph of the group $\mathbb{G}$ is r -recurrent if and only if $\mathbb{G}$ has a finite index subgroup isomorphic to $\mathbb{Z}$ or $\mathbb{Z}^2$.*

Now let $\chi : \mathcal{G} \rightarrow \mathbb{R}_\times$ be any group homomorphism to the multiplicative group of positive real numbers. Since the Cayley graph is invariant under the left action by $\mathbb{G}$, χ is a positive eigenfunction for the adjacency operator. Sometimes, these eigenfunctions constitute the minimal Martin boundary. Building on work in [DSW60] and [DM61], Margulis proved a theorem which implies the following [Mar66]. See also theorem 7.3 of [Woe94].

Theorem 15.13 (Margulis). *The collection of extremal positive eigenfunctions for the adjacency operator on the Cayley graph of a nilpotent group $\mathbb{G}$ is the collection of scalar multiples of group homomorphisms $\mathbb{G} \rightarrow \mathbb{R}_\times$.*

16. INFINITE INTERVAL EXCHANGE TRANSFORMATIONS

In this section, we discuss the infinite interval exchange maps that arise from return maps of straight line flows on $S(\mathcal{G}, \mathbf{w})$ to the collection of horizontal edges of rectangles. In subsection 16.1 we describe the interval exchange maps that arise in the general case. In subsection 16.2 we restrict attention to skew rotations and describe the proofs of statements made in section 5. And, in subsection 16.3 we define and discuss Maharam measures for skew rotations.

16.1. Return maps of the straight-line flows. Let $\mathcal{G}$ be an infinite, connected, bipartite, ribbon graph with bounded valance, and let $\mathbf{w} \in \mathbb{R}^{\mathcal{V}}$ be a positive function satisfying $\mathbf{A}(\mathbf{w}) = \lambda \mathbf{w}$. We let $H(\mathcal{G}, \mathbf{w})$ denote the union of horizontal edges of the rectangles making up the surface $S(\mathcal{G}, \mathbf{w})$. In this section we will describe the return maps $T_{\theta} : H(\mathcal{G}, \mathbf{w}) \rightarrow H(\mathcal{G}, \mathbf{w})$ of the flows $F_{\theta}^t : S(\mathcal{G}, \mathbf{w}) \rightarrow S(\mathcal{G}, \mathbf{w})$.

Recall, there is a bijection between the edges $e \in \mathcal{E}$ of $\mathcal{G}$ and the rectangles R_e making up $S(\mathcal{G}, \mathbf{w})$. Let I_e denote the top horizontal edge of R_e . Since this edge is identified with the bottom edge of $R_{\mathcal{N}(e)}$, it is useful to think of the points in I_e as infinitesimally below this top edge.

The rightmost endpoint of each interval I_e coincides with the left endpoint of $I_{\mathcal{E}(e)}$. Each orbit under $\mathcal{E}$ is finite, so $H(\mathcal{G}, \mathbf{w})$ is a disjoint union of the circles

$$(16.1) \quad H_a = \bigcup_{e \in \alpha^{-1}(a)} I_e.$$

(See the Definition 2.2(4) of α .) The length of the circle H_a is given by

$$(16.2) \quad \sum_{e \in \alpha^{-1}(a)} \mathbf{w} \circ \beta(e) = \mathbf{A}(\mathbf{w})(a) = \lambda \mathbf{w}(a),$$

so we may identify H_a with the circle $\mathbb{R}/\lambda \mathbf{w}(a)\mathbb{Z}$.

Now fix a unit vector $\theta = (x, y)$ with $y > 0$. As a point $p \in I_e$ flows vertically it immediately crosses into the rectangle $R_{\mathcal{N}(e)}$, which is a subset of the horizontal cylinder $cyl_{\alpha \circ \mathcal{N}(e)}$. To simplify notation let $e' = \mathcal{N}(e)$. This horizontal cylinder has vertical width $w = \mathbf{w} \circ \alpha(e')$. So the first return of p to $H(\mathcal{G}, \mathbf{w})$ occurs when $t = w/y$, and in a local chart p has moved along the vector $(\frac{wx}{y}, w)$. From our point of view it is equivalent to travel first along the vector $(0, w)$ and then along the vector $(\frac{wx}{y}, 0)$. (These three vectors form a triangle immersed inside the cylinder $cyl_{\alpha(e')}$.) In other words, we can obtain the return map T_{θ} , by first traveling straight along a vertical vector until you return to $H(\mathcal{G}, \mathbf{w})$. That is, we follow the interval exchange

$$(16.3) \quad S : H(\mathcal{G}, \mathbf{w}) \rightarrow H(\mathcal{G}, \mathbf{w}) : I_e \mapsto I_{\mathcal{N}(e)} \text{ for all } e \in \mathcal{E}.$$

This transformation sends $p \in I_e$ to the corresponding point in $I_{e'}$. We need to follow this by moving horizontally along the vector $(\frac{wx}{y}, 0)$. Thus, we follow S with a rotation in the cylinder $\alpha(e')$ by $\frac{wx}{y}$. This rotation is proportional to the circumference of the cylinder (λw). Therefore we consider the rotational flow $R : \mathbb{R} \times H(\mathcal{G}, \mathbf{w}) \rightarrow H(\mathcal{G}, \mathbf{w})$ described by the actions on the circles

$$R^u|_{H_a} : t \mapsto t + u \mathbf{w}(\alpha) \pmod{\lambda \mathbf{w}(a)}.$$

Our second transformation is therefore $R^{x/y}$. We have shown the following.

Proposition 16.1. *The first return T_{θ} of the flow F_{θ}^t to the horizontal edges $H(\mathcal{G}, \mathbf{w})$ is given by $T_{\theta} = R^{x/y} \circ S$.*

16.2. Skew Rotations. As in the introduction and in section 5, let $\mathbb{G}$ be a group with generators $\gamma_1, \dots, \gamma_n$ so that $\gamma_n \dots \gamma_1 = e$. Then given $\alpha \in \mathbb{R}/\mathbb{Z}$, we may define the skew rotation $T : \mathbb{R}/\mathbb{Z} \times \mathbb{G} \rightarrow \mathbb{R}/\mathbb{Z} \times \mathbb{G}$ as in equation 1.1.

As mentioned in section 5, we will describe a surface of the form $S(\mathcal{G}, \mathbf{w})$ so that T arises as a return map of a straight-line flow to the horizontal edges of the decomposition of $S(\mathcal{G}, \mathbf{w})$ into rectangles. Let C denote the cylinder $\mathbb{R}/\mathbb{Z} \times [0, 1/n]$ with a decomposition into n squares

and n top and bottom edges labeled $t_1, \dots, t_n$ and $b_1, \dots, b_n$. See figure 10. We apply the label t_i to the segment $[\frac{i-1}{n}, \frac{i}{n}] \times \{\frac{1}{n}\} \subset C$ and b_i to the segment $[\frac{i-1}{n}, \frac{i}{n}] \times \{0\} \subset C$. We let S be the translation surface $\mathbb{G} \times C / \sim$ where $\sim$ is a gluing of edges. For each $i \in \{1, \dots, n\}$ and $g \in \mathbb{G}$, we glue edge t_i of cylinder $\{g\} \times C$ to edge $b_{i+1 \pmod n}$ of cylinder $\{\gamma_i g\} \times C$ by parallel translation. Our surface has a decomposition into vertical cylinders as well, because a flow in the vertical direction results in passing through edges in the order $t_i, t_{i+1}, \dots, t_{i+n-1}$ with subscripts written modulo n . This results in visiting a list of horizontal cylinders of the form

$$(16.4) \quad \{g\} \times C, \{\gamma_i g\} \times C, \{\gamma_{i+1} \gamma_i g\} \times C, \dots, \{\gamma_{i+n-1} \dots \gamma_i g\} \times C.$$

By our assumed group relation, $\gamma_{i+n-1} \dots \gamma_{i+1} \gamma_i g = g$ and the vertical trajectory returns to its starting point after crossing n horizontal edges. That is, we have a vertical decomposition into cylinders of inverse modulus n as well. In particular, S is of the form $S(\mathcal{G}, \mathbf{w}_{\frac{1}{n}})$, where $\mathcal{G}$ is the valance n cylinder intersection graph and $\mathbf{w}_{\frac{1}{n}} \in \mathbb{R}^{\mathcal{V}}$ is the constant function with value $\frac{1}{n}$. This is an eigenfunction of eigenvalue $\lambda = n$. Note that, if we choose the trajectory to start on the edge labeled t_1 of $g \times C$, then it passes through the cylinders

$$(16.5) \quad \{\eta_1 g\} \times C, \{\eta_2 g\} \times C, \dots, \{\eta_n g\} \times C,$$

where the group elements η_i are defined to be

$$(16.6) \quad \eta_1 = e, \quad \eta_2 = \gamma_1, \quad \text{and} \quad \eta_i = \gamma_{i-1} \dots \gamma_2 \gamma_1 \text{ for } 2 \leq i \leq n.$$




FIGURE 10. The horizontal cylinder C when $n = 5$.

We will briefly describe the graph $\mathcal{G}$ obtained by this construction. The horizontal cylinders and hence the nodes of the graph in $\mathcal{A}$ are identified with the group $\mathcal{G}$. We use $a_g \in \mathcal{A}$ to denote the node associated to $g \in \mathcal{G}$. If x is a point of t_1 the vertical straight line flow crosses the cylinders as described by equation 16.5. Every vertical cylinder passes through some t_1 . Thus the vertical cylinders can be identified by an ordered n -tuple of the form $(\eta_1 g, \eta_2 g, \dots, \eta_n g) \in \mathbb{G}^n$. We use $[g]$ to abbreviate this n -tuple. We denote the vertex associated to this cylinder by $b_g \in \mathcal{B}$. Finally the collection of edges is given by the condition

$$(16.7) \quad a_g \sim b_{g'} \text{ if and only if } g = \eta_i g' \text{ for some } 1 \leq i \leq n.$$

Hence edges correspond to the choice of a n -tuple $[g] = (\eta_1 g, \eta_2 g, \dots, \eta_n g)$ and an element $\eta_i g$ of the n -tuple. This choice refers to the edge $\overline{a_{\eta_i g} b_g}$. The ribbon graph structure is given by

$$\mathcal{N}(\overline{a_{\eta_i g} b_g}) = \overline{a_{\eta_{i+1} g} b_g} \quad \text{and} \quad \mathcal{E}(\overline{a_{\eta_i g} b_g}) = \overline{a_{\eta_i g} b_{\eta_{i+1}^{-1} \eta_i g}},$$

where subscript addition is taken modulo n .

As in the previous section, we may identify the circle $\mathbb{R}/\mathbb{Z} \times \{g\}$ with the top edge of $\{g\} \times C$ so that edge t_i is identified with the interval $[\frac{i-1}{n}, \frac{i}{n}] \times \{g\}$. In this case, the skew rotation T with rotation constant α becomes the return map of the straight-line flow F_θ^t in

direction $\boldsymbol{\theta} = (\alpha - \frac{1}{n}, \frac{1}{n}) / \|(\alpha - \frac{1}{n}, \frac{1}{n})\|$. Recall that Proposition 5.1 stated that $\alpha \in \Omega_n$ if and only if this $\boldsymbol{\theta}$ is n -renormalizable.

Proof of Proposition 5.1. By Proposition 7.14, the n -renormalizable directions $\mathcal{R}_n$ consist of unit vectors $\boldsymbol{\theta}$ whose projectivizations $[\boldsymbol{\theta}]$ lie in the limit set of $\Lambda(\rho_n^G)$, but are not endpoints of intervals in $\mathbb{RP}^1 \setminus \Lambda(\rho_n^G)$ and are not fixed by parabolics in ρ_n^G . Define $\phi : \mathbb{R} \rightarrow \mathbb{R}^2$ by $\phi(\alpha) = (\alpha - \frac{1}{n}, \frac{1}{n})$. Then the correspondence between $\boldsymbol{\theta}$ and α described above satisfies $[\phi(\alpha)] = [\boldsymbol{\theta}]$ where brackets denote projectivization as above. We observe

$$\begin{aligned} [\rho_n^h \circ \phi(z)] &= [(z + 1 - \frac{1}{n}, \frac{1}{n})] = [\phi(z + 1)], \quad \text{and} \\ [\rho_n^v \circ \phi(z)] &= [(z - \frac{1}{n}, nz - 1 + \frac{1}{n})] = [(\frac{z - \frac{1}{n}}{n^2z - n + 1}, \frac{1}{n})] = [\phi(\frac{(n+1)z - 1}{n^2z - n + 1})]. \end{aligned}$$

Thus, the action of ρ_n^G on $\boldsymbol{\theta}$ induces the action of the transformations in equation 1.2 on z . That is, the above action is $\phi^{-1} \circ \rho_n^G \circ \phi$ up to scaling. From the description of $\mathcal{R}_n$ in 7.14 and the definition of Ω_n in the introduction, we see $[\phi(\Omega_n)] = [\mathcal{R}_n]$. $\square$

As mentioned in section 5, the graph $\mathcal{G}$ is not a Cayley graph of $\mathbb{G}$. However, the graph $\mathcal{G}_{\mathcal{A}}$ is. Recall $\mathcal{G}_{\mathcal{A}}$ is the graph with vertex set $\mathcal{A}$ and with an edge between $a_g, a_{g'} \in \mathcal{A}$ for every path of length 2 in $\mathcal{G}$ from a_g to $a_{g'}$ in $\mathcal{G}$. In particular, we allow backtracking and therefore loops in $\mathcal{G}_{\mathcal{A}}$. This definition makes $\mathcal{G}_{\mathcal{A}}$ valance n^2 , since $\mathcal{G}$ is valance n . Let $a_g \in \mathcal{A}$. Based on equation 16.7, such each path of length 2 is of the form

$$a_g \sim b_{\eta_i^{-1}g} \sim a_{\eta_j\eta_i^{-1}g}$$

for some $g \in \mathbb{G}$, $1 \leq i \leq n$ and $1 \leq j \leq n$. In particular, this shows that $\mathcal{G}_{\mathcal{A}}$ is a generalized Cayley graph with generating set $\{\eta_j\eta_i^{-1}\}$, proving the first half of Proposition 5.2. To complete the proof, we need to discuss how the positive eigenfunctions for the adjacency operators on $\mathcal{G}_{\mathcal{A}}$ and $\mathcal{G}$ relate. This is the content of the following proposition.

Proposition 16.2. *Let $\mathcal{G}$ be bipartite, and define $\mathcal{G}_{\mathcal{A}}$ as above. Then the restriction map $\mathbb{R}^{\mathcal{V}} \rightarrow \mathbb{R}^{\mathcal{A}}$ given by $\mathbf{f} \mapsto \mathbf{f}|_{\mathcal{A}}$ induces a bijection between the positive eigenfunctions of $\mathcal{G}$ and those of $\mathcal{G}_{\mathcal{A}}$.*

Proof. It is clear that if $\mathbf{f}$ is a positive eigenfunctions of $\mathcal{G}$ with eigenvalue λ then $\mathbf{f}|_{\mathcal{A}}$ is a positive eigenfunction with eigenvalue λ^2 . We will provide an inverse map which depends on λ . Fix $\lambda > 0$. Given $\mathbf{g} \in \mathbb{R}^{\mathcal{A}}$, define $\tilde{\mathbf{g}} \in \mathbb{R}^{\mathcal{V}}$ according to the rule that $\tilde{\mathbf{g}}(a) = \mathbf{g}(a)$ for $a \in \mathcal{A}$ and

$$\tilde{\mathbf{g}}(b) = \frac{1}{\lambda} \sum_{a \sim b} \mathbf{g}(a) \quad \text{for } b \in \mathcal{B}.$$

If $\mathbf{g}$ is an positive eigenfunction of $\mathcal{G}_{\mathcal{A}}$ with eigenvalue λ^2 , then $\tilde{\mathbf{g}}$ is a positive eigenfunction of $\mathcal{G}$ with eigenvalue λ . In this case we have $\mathbf{g} = \tilde{\mathbf{g}}|_{\mathcal{A}}$. And, if $\mathbf{f}$ is a positive eigenfunction of $\mathcal{G}$ then $\mathbf{f}|_{\mathcal{A}} = \mathbf{f}$. $\square$

16.3. Maharam measures. Maharam measures were introduced in [Mah64]. We will follow the treatment provided in the introduction of [ANSS02], but we restrict to the case of skew products with discrete groups. Maharam measures arise in the more general setting of skew products. If $\tau : X \rightarrow X$ is a measurable transformation and $\psi : X \rightarrow \mathbb{G}$ is any measurable map to a group, we define the *skew product transformation*

$$(16.8) \quad T : X \times \mathbb{G} \rightarrow X \times \mathbb{G}; (x, g) \mapsto (\tau(x), \psi(x)g).$$

In this paper $\mathbb{G}$ is always a discrete group.

Let $\mathcal{B}$ denote the Borel σ -algebra on a space X . If $h : X \rightarrow \mathbb{R}_+$ is measurable, we call the probability measure μ a (h, τ) -conformal measure if $\mu \circ \tau \sim \mu$ and

$$(16.9) \quad \frac{d\mu \circ \tau}{d\mu} = h \quad \mu\text{-a.e.}$$

In many cases, this choice of μ is known to exist and be unique. See for instance [Con09].

Now let $\chi : \mathbb{G} \rightarrow \mathbb{R}_+$ be any group homomorphism to the multiplicative group of positive real numbers. Let μ be an $(\chi \circ \psi, \tau)$ -conformal measure. For each $g \in \mathbb{G}$, we can consider the maps $\pi_g : X \rightarrow X \times \mathbb{G}$ given by $\pi_g(x) = (x, g)$. The Maharam measure associated to μ and χ is the measure $\tilde{\mu}_\chi$ on $X \times \mathbb{G}$ defined so that

$$(16.10) \quad \tilde{\mu}_\chi \circ \pi_g = \frac{1}{\chi(g)} \mu.$$

Such a measure is always invariant under the skew product transformation T . The scaling factor of χ from $(\chi \circ \psi, \tau)$ -conformality cancels with the $\frac{1}{\chi}$ factor that arises from the change in the $\mathbb{G}$ -coordinate under T . (Maharam measures for skew products with non-discrete groups can also be considered [ANSS02] [Con09].)

We now return to the case of skew rotations. The skew rotation T of equation 1.1 is a skew product with the irrational rotation $\tau : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z}$ given by $\tau(x) = x + \alpha \pmod{1}$. Via the identification of $\mathbb{G}$ with the vertices of the Cayley graph $\mathcal{G}_A$, the homomorphism χ is an example of a positive eigenfunction for the adjacency operator of $\mathcal{G}_A$. This yields a connection between the Maharam measures and the pullback measures of the topological conjugacy provided by Corollary 5.3 and Theorem 1.2.

In particular, we have the following.

Proposition 16.3. *Suppose $\alpha \in \Omega_n$. Consider the skew rotation T as in equation 1.1, and let $\psi : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{G}$ be as in this equation. Let $\chi : \mathbb{G} \rightarrow \mathbb{R}_+$ be any group homomorphism. Then,*

- (1) *There exists a unique $(\chi \circ \psi, T)$ -conformal measure, and therefore a unique Maharam measure $\tilde{\mu}_\chi$ on $\mathbb{R}/\mathbb{Z} \times \mathbb{G}$.*
- (2) *Let $\chi^{-1} : \mathbb{G} \rightarrow \mathbb{R}_+$ be the map $g \mapsto \frac{1}{\chi(g)}$. The function $\chi^{-1} \in \mathbb{R}^A$ is also a positive eigenfunction of the adjacency operator of $\mathcal{G}_A$. Let $\tilde{\chi}^{-1} \in \mathbb{R}^\mathcal{V}$ be the eigenfunction of the adjacency operator of $\mathcal{G}$ associated to χ^{-1} by restriction as in Proposition 16.2, and let λ be the associated eigenvalue. Then the Maharam measure $\tilde{\mu}_\chi$ is the pull back of Lebesgue measure on $H(\mathcal{G}, \frac{1}{\lambda} \tilde{\chi}^{-1})$ under the topological conjugacy provided in Theorem 1.2.*

We remark that statement (1) in the case of $\mathbb{G}$ abelian and for T any irrational rotation is studied in [Con09].

Proof. We first concentrate on proving statement (2). Let ν be the pull back of Lebesgue measure on $H(\mathcal{G}, \tilde{\chi}^{-1})$ to $\mathbb{R}/\mathbb{Z} \times \mathbb{G}$. We observe that the transformation $\phi_g : (t, x) \mapsto (t, gx)$ has the property that

$$(16.11) \quad \frac{d\nu \circ \phi_g}{d\nu} = \frac{1}{\chi(g)}.$$

This follows from uniqueness of the pullback measure given an eigenfunction. See Theorem 4.1. Recall $\pi_g : \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}/\mathbb{Z} \times \mathbb{G}$ is defined by $\pi_g(x) = (x, g)$, and consider the measure

$\mu = \nu \circ \pi_e$ on $\mathbb{R}/\mathbb{Z}$. For $t \in \mathbb{R}/\mathbb{Z}$ we have

$$\frac{d\mu \circ \tau(t)}{d\mu(t)} = \frac{d\mu(t + \alpha)}{d\mu(t)} = \frac{d\nu \circ T(t, \psi(t)^{-1})}{d\nu(t, e)},$$

because $T(t, \psi(t)^{-1}) = (t + \alpha, e)$. Therefore, by T invariance of ν we have

$$\frac{d\mu \circ \tau(t)}{d\mu(t)} = \frac{d\nu \circ \phi_{\psi^{-1}(t)}(t, e)}{d\nu(t, e)} = \frac{1}{\chi(\psi^{-1}(t))} = \chi \circ \psi(t).$$

Thus μ satisfies equation 16.9 in the definition of $(\chi \circ \psi, \tau)$ -conformal. However, note that μ is not a probability measure. By equation 16.2, we have $\mu(\mathbb{R}/\mathbb{Z}) = \lambda$, where $\mathbf{A}(\tilde{\chi}^{-1}) = \lambda \tilde{\chi}^{-1}$. Therefore $\frac{1}{\lambda} \mu$ is a $(\chi \circ \psi, \tau)$ -conformal measure, and $\frac{1}{\lambda} \nu$ is the associated Maharam measure. This measure $\frac{1}{\lambda} \mu$ arises from pulling back Lebesgue measure on $H(\mathcal{G}, \frac{1}{\lambda} \tilde{\chi}^{-1})$ using the topological conjugacy.

Now we will prove the first statement by showing uniqueness of the Maharam measure. Now suppose that $\tilde{\mu}_\chi$ is a Maharam measure associated to the homomorphism χ , and μ is the associated $(\chi \circ \psi, \tau)$ -conformal measure on $\mathbb{R}/\mathbb{Z}$. Note that $\tilde{\mu}_\chi$ is T invariant. We will use $\tilde{\mu}_\chi^T$ to denote the transverse measure associated to $\tilde{\mu}_\chi$. There is a natural $\mathbb{G}$ action $\Delta^\mathbb{G}$ on $\mathcal{G}$ given by

$$\Delta^g(a_{g'}) = a_{g'g^{-1}} \quad \text{and} \quad \Delta^g(b_{g'}) = b_{g'g^{-1}}.$$

This induces an action $\tilde{\Delta}^\mathbb{G}$ on $S(\mathcal{G}, \mathbf{w}_{\frac{1}{n}})$ which moves rectangles around according to the action of $\Delta^\mathbb{G}$. The property of being Maharam given in equation 16.10 extends to the transverse measure as

$$(16.12) \quad \tilde{\mu}_\chi^T \circ \tilde{\Delta}^g(\tau) = \chi(g) \tilde{\mu}_\chi^T(\tau),$$

By corollary 6.30, $\Psi_\theta(\tilde{\mu}_\chi^T) = \Xi(\mathbf{f})$ for some $\mathbf{f}$. (Recall, $\Psi_\theta(\tilde{\mu}_\chi^T)$ is the associated cohomology class, and Ξ promotes an element of $\mathbb{R}^\nu$ to such a class as in equation 6.5.) By Lemma 6.2 and Theorem 6.3, $\mathbf{f}$ uniquely determines the measure $\tilde{\mu}_\chi^T$. Therefore, to show uniqueness it is sufficient to show the uniqueness of $\mathbf{f}$ given χ .

Consider $b_e \in \mathcal{B}$, and let σ_e be a horizontal saddle connection crossing the vertical cylinder cyl_{b_e} . For an arbitrary $b_g \in \mathcal{B}$ set $\sigma_g = \tilde{\Delta}^{g^{-1}}(\sigma_e)$. This is a horizontal saddle connection crossing cyl_{b_g} . Then by equation 16.12,

$$(16.13) \quad \mathbf{f}(b_g) = \tilde{\mu}_\chi^T(\sigma_g) = \tilde{\mu}_\chi^T \circ \tilde{\Delta}^{g^{-1}}(\sigma_e) = \frac{1}{\chi(g)} \tilde{\mu}_\chi^T(\sigma_e) = \frac{1}{\chi(g)} \mathbf{f}(b_e).$$

A similar argument with vertical saddle connections shows that $\mathbf{f}(a_g) = \frac{1}{\chi(g)} \mathbf{f}(a_e)$. This determines $\mathbf{f}$ up to the two parameters $\mathbf{f}(a_e)$ and $\mathbf{f}(b_e)$. Recall the definitions of η_i given in equation 16.6. We can compute

$$\begin{aligned} \mathbf{A}(\mathbf{f})(a_g) &= \sum_{b \sim a} \mathbf{f}(b) = \sum_{i=1}^n \mathbf{f}(b_{\eta_i^{-1}g}) = \mathbf{f}(b_g) \sum_{i=1}^n \chi(\eta_i) \\ \mathbf{A}(\mathbf{f})(b_g) &= \sum_{a \sim b} \mathbf{f}(a) = \sum_{i=1}^n \mathbf{f}(a_{\eta_i g}) = \mathbf{f}(a_g) \sum_{i=1}^n \frac{1}{\chi(\eta_i)}. \end{aligned}$$

It will be useful to define the constants $\delta = \sum_{i=1}^n \frac{1}{\chi(\eta_i)}$ and $\epsilon = \sum_{i=1}^n \chi(\eta_i)$. It follows from the above equations that $\mathbf{f}|_{\mathcal{A}}$ is an eigenfunction for the adjacency operator acting on $\mathcal{G}_{\mathcal{A}}$ with eigenvalue $\delta\epsilon$. In fact, $\mathbf{f}|_{\mathcal{A}}$ is a scalar multiple of the eigenfunction $\chi^{-1} : \mathbb{G} \rightarrow \mathbb{R}_\times$. We define

a modified version of $\mathbf{f}$ according to the rule $\widehat{\mathbf{f}}(a_g) = \mathbf{f}(a_g)$ and $\widehat{\mathbf{f}}(b_g) = \mathbf{f}(a_g)\sqrt{\delta/\epsilon}$. It can be checked that $\widehat{\mathbf{f}}$ is an eigenvector of $\mathbf{A}$ with eigenvalue $\lambda = \sqrt{\delta\epsilon}$. Therefore, the eigenfunction $\widehat{\mathbf{f}} \in \mathbb{R}^\nu$ is the canonical extension of the eigenfunction $\mathbf{f}|_{\mathcal{A}} \in \mathbb{R}^A$ given by Proposition 16.2. Therefore $\widehat{\mathbf{f}}$ is a multiple of $\widetilde{\chi}^{-1}$. We will show that $\mathbf{f}$ is in the plane $P_{\widehat{\mathbf{f}}}$ associated to $\widehat{\mathbf{f}}$ by section 6.4. Note that the ratio $q = \frac{\mathbf{f}(b_g)}{\mathbf{f}(a_g)}$ is independent of the choice of $g \in \mathbb{G}$. Therefore,

$$\mathbf{f}(a_g) = \widehat{\mathbf{f}}(a_g) \quad \text{and} \quad \mathbf{f}(b_g) = q\widehat{\mathbf{f}}(b_g)\sqrt{\epsilon/\delta}.$$

This shows that $\mathbf{f}$ is in the plane associated to $\widehat{\mathbf{f}}$. Since $\widetilde{\mu}_\chi^T$ is a transverse measure, $\mathbf{f}$ must be a θ -survivor. By Proposition 6.23, there is a unique ray in this plane of θ -survivors. This determines $\mathbf{f}$ up to a positive scalar constant. Now note that by equation 16.10, the Maharam measure $\widetilde{\mu}_\chi$ must satisfy $\widetilde{\mu}_\chi(\mathbb{R}/\mathbb{Z} \times \{e\}) = 1$. Therefore, we know that $\mathbf{A}(\mathbf{f})(a_e) = 1$. This fixes the scalar constant, and determines $\mathbf{f}$ as desired. $\square$

Using the notion of Maharam measure, we have a stronger version of Corollary 5.4.

Corollary 16.4. *Suppose $\mathcal{G}$ is nilpotent and $\alpha \in \Omega_n$. Then the locally finite ergodic invariant measures of the skew rotation T of equation 1.1 are the Maharam measures, which exist and are unique for every homomorphism $\chi : \mathbb{G} \rightarrow \mathbb{R}_\times$.*

REFERENCES

- [Aar97] Jon Aaronson. *An introduction to infinite ergodic theory*, volume 50 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 1997.
- [Anc88] Alano Ancona. Positive harmonic functions and hyperbolicity. In *Potential theory—surveys and problems (Prague, 1987)*, volume 1344 of *Lecture Notes in Math.*, pages 1–23. Springer, Berlin, 1988.
- [ANSS02] Jon Aaronson, Hitoshi Nakada, Omri Sarig, and Rita Solomyak. Invariant measures and asymptotics for some skew products. *Israel J. Math.*, 128:93–134, 2002.
- [AT07] Charalambos D. Aliprantis and Rabee Tourky. *Cones and duality*. Graduate Studies in Mathematics 84. Providence, RI: American Mathematical Society (AMS). xiv, 279 p., 2007.
- [BBi01] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001.
- [BM74] Alan F. Beardon and Bernard Maskit. Limit points of Kleinian groups and finite sided fundamental polyhedra. *Acta Math.*, 132:1–12, 1974.
- [CGL06] R. Chamanara, F. P. Gardiner, and N. Lakic. A hyperelliptic realization of the horseshoe and baker maps. *Ergodic Theory Dynam. Systems*, 26(6):1749–1768, 2006.
- [CK76] J.-P. Conze and M. Keane. Ergodicité d’un flot cylindrique. In *Séminaire de Probabilités, I (Univ. Rennes, Rennes, 1976)*, Exp. No. 5, page 7. Dépt. Math. Informat., Univ. Rennes, Rennes, 1976.
- [CK77] B. D. Craven and J. J. Koliha. Generalizations of Farkas’ theorem. *SIAM J. Math. Anal.*, 8(6):983–997, 1977.
- [Con09] Jean-Pierre Conze. Recurrence, ergodicity and invariant measures for cocycles over a rotation. Assani, Idris (ed.), *Ergodic theory. Probability and ergodic theory workshops*, Chapel Hill, NC, USA, February 15–18, 2007 and February 14–17, 2008. Providence, RI: American Mathematical Society (AMS). Contemporary Mathematics 485, 45–70 (2009)., 2009.
- [DM61] E. B. Dynkin and M. B. Maljutov. Random walk on groups with a finite number of generators. *Dokl. Akad. Nauk SSSR*, 137:1042–1045, 1961.
- [DSW60] Joseph L. Doob, J.L. Snell, and R.E. Williamson. Application of boundary theory to sums of independent random variables. *Contrib. Probab. Stat., Essays in Honor of H. Hotelling*, 182–197 (1960)., 1960.
- [Far02] J. Farkas. Über die theorie der einfachen ungleichungen. *J. Reine Angew. Math.*, 124:1–24, 1902.
- [Hoo08a] W. Patrick Hooper. The Bouw-Möller lattice surfaces and eigenvectors of grid graphs. preprint, 2008.

- [Hoo08b] W. Patrick Hooper. Dynamics on an infinite surface with the lattice property. preprint, 2008.
- [HS09] Pascal Hubert and Gabriela Schmithüsen. Infinite translation surfaces with infinitely generated Veech groups. preprint, 2009.
- [HW08] Pascal Hubert and Barak Weiss. Dynamics on the infinite staircase. preprint, 2008.
- [HW10] W. Patrick Hooper and Barak Weiss. Generalized staircases: recurrence and symmetry authors: W. patrick hooper, barak weiss. *Ann. Inst. Fourier*, 2010. to appear.
- [KH95] Anatole Katok and Boris Hasselblatt. *Introduction to the modern theory of dynamical systems*, volume 54 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1995. With a supplementary chapter by Katok and Leonardo Mendoza.
- [Mah64] D. Maharam. Incompressible transformations. *Fund. Math.*, 56:35–50, 1964.
- [Mar66] G. A. Margulis. Positive harmonic functions on nilpotent groups. *Soviet Math. Dokl.*, 7:241–244, 1966.
- [Mas82] Howard Masur. Interval exchange transformations and measured foliations. *Ann. of Math. (2)*, 115(1):169–200, 1982.
- [McM06] Curtis T. McMullen. Prym varieties and Teichmüller curves. *Duke Math. J.*, 133(3):569–590, 2006.
- [MT98] Katsuhiko Matsuzaki and Masahiko Taniguchi. *Hyperbolic manifolds and Kleinian groups*. Oxford Mathematical Monographs. The Clarendon Press Oxford University Press, New York, 1998. Oxford Science Publications.
- [MW89] Bojan Mohar and Wolfgang Woess. A survey on spectra of infinite graphs. *Bull. London Math. Soc.*, 21(3):209–234, 1989.
- [Pru64] William E. Pruitt. Eigenvalues of non-negative matrices. *Ann. Math. Statist.*, 35:1797–1800, 1964.
- [PSV09] Piotr Przytycki, Gabriela Schmithuesen, and Ferran Valdez. Veech groups of loch ness monsters. preprint <http://arxiv.org/abs/0906.5268>, 2009.
- [PW88] Massimo A. Picardello and Wolfgang Woess. Harmonic functions and ends of graphs. *Proc. Edinburgh Math. Soc. (2)*, 31(3):457–461, 1988.
- [Sch78] K. Schmidt. A cylinder flow arising from irregularity of distribution. *Compositio Math.*, 36(3):225–232, 1978.
- [Sen81] E. Seneta. *Nonnegative matrices and Markov chains*. Springer Series in Statistics. Springer-Verlag, New York, second edition, 1981.
- [SU09] John Smillie and Corinna Ulcigrai. Symbolic coding for linear trajectories in the regular octagon. preprint <http://arxiv.org/abs/0905.0871>, 2009.
- [Thu88] William P. Thurston. On the geometry and dynamics of diffeomorphisms of surfaces. *Bull. Amer. Math. Soc. (N.S.)*, 19(2):417–431, 1988.
- [Tro04] S. Troubetzkoy. Recurrence and periodic billiard orbits in polygons. *Regul. Chaotic Dyn.*, 9(1):1–12, 2004.
- [Var86] Nicholas Th. Varopoulos. Théorie du potentiel sur des groupes et des variétés. *C. R. Acad. Sci. Paris Sér. I Math.*, 302(6):203–205, 1986.
- [Vee89] W. A. Veech. Teichmüller curves in moduli space, Eisenstein series and an application to triangular billiards. *Invent. Math.*, 97(3):553–583, 1989.
- [VJ67] D. Vere-Jones. Ergodic properties of nonnegative matrices. I. *Pacific J. Math.*, 22:361–386, 1967.
- [VJ68] D. Vere-Jones. Ergodic properties of nonnegative matrices. II. *Pacific J. Math.*, 26:601–620, 1968.
- [Woe94] Wolfgang Woess. Random walks on infinite graphs and groups—a survey on selected topics. *Bull. London Math. Soc.*, 26(1):1–60, 1994.
- [Woe00] Wolfgang Woess. *Random walks on infinite graphs and groups*, volume 138 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, 2000.
- [ZK75] A. N. Zemljakov and A. B. Katok. Topological transitivity of billiards in polygons. *Mat. Zametki*, 18(2):291–300, 1975.